



HAL
open science

Recommendations for the analysis of hemocyte-related immunocompetent oxidative activity in the freshwater snail *Lymnaea stagnalis*

P. Boisseaux, P. Noury, Marie Laure Delignette-Muller, H. Thomas, J. Garric

► To cite this version:

P. Boisseaux, P. Noury, Marie Laure Delignette-Muller, H. Thomas, J. Garric. Recommendations for the analysis of hemocyte-related immunocompetent oxidative activity in the freshwater snail *Lymnaea stagnalis*. *Journal of Xenobiotics*, 2016, 6, pp.25-27. 10.4081/xeno.2016.6585 . hal-02016380

HAL Id: hal-02016380

<https://univ-lyon1.hal.science/hal-02016380>

Submitted on 1 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Recommendations for the analysis of hemocyte-related immunocompetent oxidative activity in the freshwater snail *Lymnaea stagnalis*

P. Boisseaux,¹ P. Noury,¹

M.-L. Delignette-Muller,^{2,3}

H. Thomas,⁴ J. Garric¹

¹Irstea, UR MALY, Centre de Lyon-Villeurbanne, Villeurbanne; ²Université de Lyon, Université Lyon 1, CNRS, UMR 5558, Laboratoire de Biométrie et Biologie Evolutive, Villeurbanne; ³Université de Lyon, VetAgro Sup Campus Vétérinaire de Lyon, Marcy l'Etoile; ⁴Littoral ENvironnement et Sociétés (LIENSs) - UMR 7266, La Rochelle, France

Introduction

One of the most remarkable features of immune systems through many species is the capacity of phagocytes to produce diverse form of oxidizing agents in order to destruct invaders. Reactive oxygen species (ROS) represent a family of various reactive molecules that play pivotal roles in a myriad of physiological and pathophysiological events. The NADPH oxidase catalytic complexes (also called *respiratory burst oxidases*) are most often admitted to be responsible for the hemocyte immunocompetent oxidative activity.¹⁻³ These enzymes are activated when the cell is exposed to appropriate stimuli such as *pathogen-associated-molecular patterns* (PAMPs): the phenomenon is inducible.^{4,5} In several mollusks, the phagocytosis of microorganisms promotes the activation of the NADPH oxidases from hemocytes,⁶⁻⁹ in particular in *Lymnaea stagnalis*.¹⁰ In *L. stagnalis*, it has been observed that repeated hemolymph samplings at a 48 h frequency did not affect the snails immunocapacity defined as the combination of hemocyte density and hemocyte viability.¹¹ However, other endpoints may still be affected such as functional markers for example, as it was reported for lysosomal enzymatic activity in *L. stagnalis*.¹² The aims of this study was: i) to develop a method in order to follow kinetics of hemocyte-related oxidative activities under basal and immunostimulated *ex vivo* conditions; and ii) to assess possible effects of repeated hemolymph collections on hemocyte oxidative activities.

Materials and Methods

A total of 24 calibrated snails (32±2 mm shell-length), free of parasites, were taken from the breeding stock of our laboratory and randomly assigned into three aquaria. Snails were maintained in a flow-through system of clean groundwater as for breeding stocks (20±1°C, 16 h/8 h of light/dark cycles, *ad libitum* feeding with Tetraphyll©). The experiment duration lasted for four days and the hemolymph of individual snails was collected mechanically as described by Sminia¹³ and at three times: at t=0 h, t=48 h and t=96 h. When tickling the foot sole of the snail with the tip of a micropipette, the snail retracts into its shell and extrudes hemolymph through the hemal pore. The methodology of ROS measurement was adapted from Adema, van Deutekom-Mulder,¹⁰ Moss and Allam¹⁴ and Hahn, Bender.⁹ For each snail, 25 µL of hemolymph were gently deposited into 12 wells of a 96-well plate (treated for cell culture, Greiner, Cat. No.

655180). The plates were centrifuged (100 g, 20 min), the lymph was removed and then replaced by snail saline buffer solution (SSB),¹⁵ pre-warmed at room temperature, to achieve 200 µL.well⁻¹. An inhibitor of the NADPH-oxidase, namely the *protocatechuic acid* (PCA),^{9,10,16} was used for the half of the wells (160 µM). After 30 min of slow agitation on a shaker, the half of these two groups of wells (N=24 triplicates per organism per *ex vivo* condition) was supplemented with zymosan particles (50 µL of a 2 mg.mL⁻¹ of SSB working solution) (Sigma Z4250), which are PAMPs from yeasts (*Saccharomyces cerevisiae*). The non-stimulated cells received 50 µL of SSB. A stock solution of the cell-permeant probe 2',7'-dichlorodihydrofluorescein diacetate (H₂DCFDA) (D6883 Sigma) was previously prepared in ethanol (20 mM) and conserved at -20°C. Straightforwardly after zymosan and SSB additions, the H₂DCFDA stock solution was diluted at 2 mM using SSB and then 10 µL were added into each well. Samples were homogenized and then the plate was read using a spectrofluorimeter (TECAN, Safire). The fluorescence at 485/530 nm (EX/EM) wavelengths was measured for 1 h every 2 min. First, the raw data of relative fluorescence obtained from kinetics were converted into amount of converted product DCF (*Dichlorofluorescein*) by using a standard curve. Second, the activity was calculated as the mean of triplicates for each *ex vivo* condition from each snail, according to the formula:

$$\text{Oxidative activity (fmol of DCF /min/}\mu\text{L)} = \frac{\text{slope } X_{ij} (20-40 \text{ min})}{25 \mu\text{L}} \quad (1)$$

Correspondence: Jeanne Garric, Irstea, UR MALY, centre de Lyon-Villeurbanne, 5 rue de la Doua, BP 32108, 69616 Villeurbanne Cedex, France. E-mail: jeanne.garric@irstea.fr

Key words: *Lymnaea stagnalis*; hemocyte-related oxidative activity; immunocompetence.

Funding: the Rhône-Alpes region provided financial support to Paul Boisseaux (IRC3 Environnement).

Conference presentation: part of this paper was presented at *ECOBIM meeting*, 2016 May-June, Le Havre, France.

This work is licensed under a Creative Commons Attribution NonCommercial 4.0 License (CC BY-NC 4.0).

©Copyright P. Boisseaux et al., 2016
Licensee PAGEPress, Italy
Journal of Xenobiotics 2016; 6:6585
doi:10.4081/xeno.2016.6585

where X is the amount of DCF (fmol) at the minute i for the snail j .

The datasets were analyzed by fitting linear mixed-effects models (LMEM) using the *lmer* function from the *lme4* package¹⁷ in the R software. Beforehand, we applied a log₁₀ transformation to the data to normalize the distributions. LMEM were first built with three fixed factors, the *zymosan* (*i.e.*, presence/absence), the *PCA* (*i.e.*, presence/absence), and the day, and two nested random factors, the snail and the replicate (aquarium). The initial model also included interaction terms between *zymosan* and *PCA*, *zymosan* and *day*, *PCA* and *day*. The model was simplified by removing the effects that did not appear significant. An effect was considered as statistically significant when the 95 % confidence interval of any of its corresponding coefficients did not contain the value 0.

Results and Discussion

Baseline activities

In the present study, the data are expressed in fmol of DCF.min⁻¹.µL⁻¹. It corresponds to the oxidative activity from hemocytes (*i.e.*, the lymph was removed) per microliter of hemolymph. The transformation of relative fluorescent units in amount of substance converted has been made in order to facilitate the data comparisons between independent experiments and the use of different spectrofluorimeter settings (*e.g.*, gains). By using the corresponding standard curve, the same data calculation is possible. The underlying question that justified such a methodology was: what is

the hemocyte-related immunocompetent oxidizing ability in the circulating system of the snail? Indeed, interpretation of ROS productions are tricky in immunopathology because they encompass many origins (*e.g.*, mitochondrial, endoplasmic reticulum, peroxisome activities, *etc.*),³ NADPH-activity,¹⁰ mechanical-stress¹⁸ and xenobiotic-oxidative stress^{15,19,20} among others. They may constitute indication of toxicity or of immunocompetence (oxidative stress *vs* immunocompetent oxidative activity) and so basal together with stimulated activities are both complementary. Last, the lymph was removed in order to reduce ROS background coming from other pathways than hemocytes.

ROS production showed a strong increase when the samples were stimulated with zymosan and a strong decrease when they were incubated with PCA (Figure 1). A transient reduction phase occurred at approximately 25 min (Figure 2). Afterwards, ROS production increased linearly. The mean (in decimal logarithm) at day 0 for samples without zymosan and without PCA was estimated to 2.39 [with 95% confidence interval (CI) of (2.26-2.51)]. This \log_{10} value corresponds to a hemocyte-related oxidative activity of 245.47 fmol of DCF.min⁻¹.μL⁻¹. In average, the immunostimulation with zymosan increased significantly the \log_{10} value of the basal activity by 0.82 (95% CI of 0.74-0.90), which corresponds to an increase by a factor 6.61 of the activity ($10^{0.82}=6.61$). The presence of the inhibitor PCA decreased significantly the \log_{10} value of the basal activity by 0.23 (95% CI of -0.19 - -0.25). A significant interaction was found between zymosan and PCA. Hence, the presence of PCA diminished by 0.53 (-0.53 = -0.23 - 0.30) the \log_{10} values of the zymosan-stimulated oxidative activity instead of 0.23 in the absence of zymosan. Addition of PCA did not lead to a proportional inhibitory effect whether the samples were immunostimulated with zymosan or not.

Repeated collections

Data analysis from the model revealed a significant effect of the day 4 but not the day 2 compared to the day 0. The oxidative activities at day 4 were estimated to multiply by 0.64 the ones from day 0. In other words, the third hemolymph collection led to a significant decrease in the oxidative activities (*i.e.*, basal and immunostimulated) of approximately 36% from the oxidative activities measured at day 0. No significant interactions were found neither between the day and the zymosan nor between the day and the PCA. The repeated hemolymph collections decreased both basal and zymosan-stimulated oxidative activity from hemocytes. The decreases are particularly important for the third hemolymph collection (*i.e.*, day 4) on Figure 1. Such phenome-




Figure 1. Hemocyte-related oxidative activities under repeated hemolymph collections. The graph represents the oxidative activity from snail's hemocytes - being sampled every 48 h - under different *ex vivo* conditions. The *ex vivo* conditions correspond to samples with zymosan (right) and without zymosan (left); both groups were supplemented with PCA (blue borders) or not (black borders).




Figure 2. Example of kinetics of hemocyte-related oxidative activities from 25 μL of hemolymph [gain=130]. The graph represents an example of the ROS kinetics obtained from raw data with the spectrofluorimeter into the different *ex vivo* conditions, for one snail, with corresponding controls. The black arrow indicates a kinetic transition (*i.e.*, unknown phenomenon) being observed in most of the snails. Note that the scale from graph B is ten times bigger than the one from the graph A (values are much higher with zymosan particles and hemolymph).

non could be the result of a shift in hemocytes maturity (linked to hemocyte renewal turnover) provoked by repeated collections and so young hemocytes would not have the cell machinery - or to a lesser extent - to carry out the immunocompetent oxidative activity. Further studies should be conducted to confirm or infirm this hypothesis. In parallel, at each sampling time the snails' hemocyte density and viability have been quantified by flow cytometry but no *sampling effect* has been detected on these endpoints.

Conclusions

The presented method has been optimized because: i) only 25 μ L of hemolymph per well replicate are necessary and so it is possible to set up several well replicates per individual; ii) spectrofluorimetry allows assessing real kinetics of a broad number of samples, what is tricky in flow cytometry; iii) lymph is removed and so the ROS production originates mainly from hemocytes; iv) zymosan particles are strong inductors of the immunocompetent oxidative activity; v) linear mixed-effect models used allow quantifying the effect of all factors, including time.

We recommend increasing the time interval (*i.e.*, more than 48 h) if more than two consecutive hemolymph collections are performed on one snail, otherwise the effect of *mechanical hemolymph collection* will likely impact the hemocyte oxidative activities and could lead to a bias in results interpretation.

References

- Bedard K, Krause K-H. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev* 2007;87:245-313.
- Buggé DM, Hégaret H, Wikfors GH, Allam B. Oxidative burst in hard clam (*Mercenaria mercenaria*) haemocytes. *Fish Shellfish Immun* 2007;23:188-96.
- Donaghy L, Hong H-K, Jauzein C, Choi K-S. The known and unknown sources of reactive oxygen and nitrogen species in haemocytes of marine bivalve molluscs. *Fish Shellfish Immun* 2015;42:91-7.
- Babior BM, Benna JE, Chanock SJ, Smith RM. The NADPH oxidase of leukocytes: the respiratory burst oxidase. *Cold Spring Harbor Monogr Archiv* 1997;34:737-83.
- Donaghy L, Kraffe E, Le Goïc N, Lambert C, Volety AK, Soudant P. Reactive oxygen species in unstimulated hemocytes of the Pacific oyster *Crassostrea gigas*: a mitochondrial involvement. *PLoS One* 2012;7:e46594.
- Lambert C, Soudant P, Jegaden M, Delaporte M, Labreuche Y, Moal J, et al. *In vitro* modulation of reactive oxygen and nitrogen intermediate (ROI/RNI) production in *Crassostrea gigas* hemocytes. *Aquaculture* 2007;270:413-21.
- Manduzio H, Rocher B, Durand F, Galap C, Leboulenger F. The point about oxidative stress in molluscs. *ISJ* 2005;2:91-104.
- Bugge DM, Hégaret H, Wikfors GH, Allam B. Oxidative burst in hard clam (*Mercenaria mercenaria*) haemocytes. *Fish Shellfish Immunol* 2007;23:188-96.
- Hahn UK, Bender RC, Bayne CJ. Production of reactive oxygen species by hemocytes of *Biomphalaria glabrata*: carbohydrate-specific stimulation. *Develop Comparat Immunol* 2000;24:531-41.
- Adema C, van Deutekom-Mulder E, van der Knaap W, Sminia T. NADPH-oxidase activity: the probable source of reactive oxygen intermediate generation in hemocytes of the gastropod *Lymnaea stagnalis*. *J Leuk Biol* 1993;54:379-83.
- Boisseaux P, Delignette-Muller M-L, Abbaci K, Thomas H, Garric J. Analysis of hemocytes in *Lymnaea stagnalis*: characterization and effects of repeated hemolymph collections. *Fish Shellfish Immun* 2016 [Epub ahead of print].
- Mohandas A, Adema C, Van der Knaap W, Sminia T. The effect of haemolymph extraction on distribution of lysosomal enzymes in *Lymnaea stagnalis* haemocytes: a cytochemical study. *Comparat Haematol Int* 1992;2:61-7.
- Sminia T. Structure and function of blood and connective tissue cells of the fresh water pulmonate *Lymnaea stagnalis* studied by electron microscopy and enzyme histochemistry. *Zeitschr Zellforsch mikroskop Anat* 1972;130:497-526.
- Moss B, Allam B. Fluorometric measurement of oxidative burst in lobster hemocytes and inhibiting effect of pathogenic bacteria and hypoxia. *J Shellfish Res* 2006;25:1051-7.
- Russo J, Lefeuvre-Orfila L, Lagadic L. Hemocyte-specific responses to the peroxidizing herbicide fomesafen in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata). *Environ Pollut* 2007;146:420-7.
- A't Hart B, Simons JM, Rijkers GT, Hoogvliet JC, Van Dijk H, Labadie RP. Reaction products of 1-naphthol with reactive oxygen species prevent NADPH oxidase activation in activated human neutrophils, but leave phagocytosis intact. *Free Radical Biol Med* 1990;8:241-9.
- Bates D, Maechler M, Bolker B, Walker S. Fitting linear mixed-effects models using {lme4}. *J Stat Softw* 2015;67:1-48.
- Lacoste A, Malham SK, Gélébart F, Cueff A, Poulet SA. Stress-induced immune changes in the oyster *Crassostrea gigas*. *Dev Comparat Immunol* 2002;26:1-9.
- Winston GW, Moore MN, Kirchin MA, Soverchia C. Production of reactive oxygen species by hemocytes from the marine mussel, *Mytilus edulis*: lysosomal localization and effect of xenobiotics. *Comparat Biochem Physiol Part C* 1996;113:221-9.
- Koutsogiannaki S, Franzellitti S, Fabbri E, Kaloyianni M. Oxidative stress parameters induced by exposure to either cadmium or 17- β -estradiol on *Mytilus galloprovincialis* hemocytes. The role of signaling molecules. *Aquat Toxicol* 2014;146:186-95.