



HAL
open science

Effects of Action Observation and Action Observation Combined with Motor Imagery on Maximal Isometric Strength

Franck Di Rienzo, Pierric Joassy, Thiago Kanthack, Tadhg Eoghan Macintyre, Ursula Debarnot, Yoann Blache, Christophe Hautier, Christian Collet, Aymeric Guillot

► **To cite this version:**

Franck Di Rienzo, Pierric Joassy, Thiago Kanthack, Tadhg Eoghan Macintyre, Ursula Debarnot, et al.. Effects of Action Observation and Action Observation Combined with Motor Imagery on Maximal Isometric Strength. *Neuroscience*, 2019, 418, pp.82-95. 10.1016/j.neuroscience.2019.08.025 . hal-02307121

HAL Id: hal-02307121

<https://univ-lyon1.hal.science/hal-02307121>

Submitted on 20 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Effects of action observation and action observation combined with motor imagery on maximal isometric strength

Franck Di Rienzo^{1*}, Pierric Joassy¹, Thiago Kanthack¹, Tadhg Eoghan MacIntyre², Ursula Debarnot¹, Yoann Blache¹, Christophe Hautier¹, Christian Collet¹ & Aymeric Guillot^{1,3}

¹ Univ Lyon, University Claude Bernard Lyon 1, Inter-university Laboratory of Human Movement Sciences, EA 7424, F-69622 Villeurbanne Cedex, France

²Health Research Institute, University of Limerick, Limerick, Ireland

² Institut Universitaire de France, Paris, France.

Short title: Mental training of isometric strength

Corresponding author:

Franck Di Rienzo, Laboratoire Interuniversitaire de Biologie de la Motricité, EA 7424, Université de Lyon, Université Claude Bernard Lyon 1, 27-29 boulevard du 11 Novembre 1918, 69622 Villeurbanne Cedex, France. Tel : +33 472432837. E-mail: franck.di-rienzo@univ-lyon1.fr

Abstract

Action observation (AO) alone or combined with motor imagery (AO+MI) has been shown to engage the motor system. While recent findings support the potential relevance of both techniques to enhance muscle function, this issue has received limited scientific scrutiny. In the present study, we implemented a counterbalanced conditions design where 21 participants performed 10 maximal isometric contractions (12s duration) of elbow flexor muscles against a force platform. During the inter-trial rest periods, participants completed i) AO of the same task performed by an expert athlete, ii) AO+MI, i.e. observation of an expert athlete while concurrently imagining oneself performing the same task, and iii) watching passively a video documentary about basketball shooting (CONTROL). During force trials, we recorded the total force and integrated electromyograms from the *biceps brachii* and *anterior deltoideus*. We also measured skin conductance from two finger electrodes as an index of sympathetic nervous system activity. Both AO and AO+MI outperformed the CONTROL condition in terms of total force (2.79-3.68%, $p < 0.001$). For all conditions, we recorded a positive relationship between the *biceps brachii* activation and the total force developed during the task. However, only during AO was a positive relationship between the activation of the *anterior deltoideus* and the total force. We interpreted the results with reference to the statements of the psychoneuromuscular theory of mental practice. Present findings extend current knowledge regarding the priming effects of AO and AO+MI on muscle function, and may contribute to the optimization of training programs in sports and rehabilitation.

Keywords: neural plasticity, neural priming, muscle function, motor cognition, conditioning

Introduction

Dose-response relationships in the development of muscle strength have been the focus of a large body of research since the middle of the 20th century (Rhea et al., 2003; Peterson et al., 2004). Strength development originates from both structural and functional adaptations (Komi, 1986). Strength development is classically associated with hypertrophy, i.e. an increase in muscle mass (Damas et al., 2015). Muscle hypertrophy occurs as a *long-term* result of resistance training (programs scheduled within a span of several weeks), and stems from an adaptive response to the mechanical and metabolic demands elicited by training (Schoenfeld, 2010). Functional adaptations also play a key role in strength development, particularly during the early stages where *short-term* improvements occur in the absence of muscle hypertrophy (Sale, 1988; Moritani, 1993). Indeed, strength produced during maximal voluntary contractions mirrors the capacity of the central nervous system to synchronise the recruitment of motor units (Clark et al., 2014). Weighted resistance training above 90% of the maximal voluntary contraction threshold emphasises neurophysiological adaptations yielding improved motor units recruitment (Kraemer et al., 1996).

Resistance training programs typically consist of physical exercise with progressive poundage. Nonetheless, the potential relevance of training with mental practice is gaining increased attention (Tod et al., 2003, 2015; Paravlic et al., 2018). Recent evidence supports the positive effects of motor imagery (MI), i.e. the voluntary process of mentally simulating a movement without engaging in its physical execution, on strength and its inclusion within resistance training programs (Tod et al., 2015; Paravlic et al., 2018). Functional brain imaging findings support the hypothesis of a functional equivalence between MI and the physical execution of the same action. MI activates brain regions controlling the actual motor preparation and execution, albeit with reduced magnitude (Lotze and Halsband, 2006; Munzert and Zentgraf, 2009; Héту et al., 2013). There is evidence of premotor, parietal and primary sensorimotor cortex activation during MI (Gerardin et al., 2000; Ehrsson et al., 2003; Solodkin et al., 2004; Gao et al., 2011). MI also recruits subcortical regions such as the basal ganglia and the cerebellum (Hanakawa et al., 2003; Szameitat et al., 2007; Wagner et al., 2008; Burianová et al., 2013). At the peripheral level, the effector-specific facilitation of corticospinal

pathways during MI is well-established (Stinear, 2010; Grosprêtre et al., 2016, for reviews). Furthermore, both Jowdy and Harris (1990) and Gandevia et al. (1997) reported low-threshold muscle activity during MI. The subliminal muscle activation reproduced the intensity and regimen of the imagined contractions (Bakker et al., 1996; Guillot et al., 2007; Lebon et al., 2008). Henceforth, rather than a purely mental state, MI should overall better be considered a “(...) *special form of motor behaviour*” (Stephan and Frackowiak, 1996).

From an applied viewpoint, MI increases the cognitive demand on brain motor regions, which then could prompt experience-based plasticity and improve motor performance (Di Rienzo et al., 2016). Pioneering investigations of the effects of MI on strength focused on behavioural measures, sometimes associated with recordings of muscle activation (Cornwall et al., 1991; Yue and Cole, 1992). There is an emerging consensus that embedding MI within resistance training programs of several weeks yields additional strength benefits. Despite methodological differences across studies (e.g. the MI to physical practice ratio administered during the training sessions), strength typically increases from 10 to 30% (Reiser et al., 2011). Improvements occur in the absence of muscle hypertrophy, hence suggesting neural adaptation (Yue and Cole, 1992). More recently, we observed strength gains after a single-session of MI (Di Rienzo et al., 2015). These findings corroborate the proposition of the psycho-neuromuscular theory of mental practice, which posits that motor simulation can be effective to improve the cortical gain over motor units (Jacobson, 1932). While short-term changes in the cortical gain over motor units might drive most of the short-term effects (Di Rienzo et al., 2015; Grosprêtre et al., 2018), Hebbian-type plasticity comparable to that elicited by physical training may be the core process mediating improvements of motor performance through MI (Di Rienzo et al., 2016, for a review). This applies to strength development paradigms, where resistance training programs with MI extend across several weeks. For instance, Ranganathan et al. (2004) demonstrated that strength improvements consecutive to MI training were associated with increased amplitudes of the cortical motor potentials generated during high-intensity strength trials (see also Yao et al., 2013; Grosprêtre et al., 2018).

Action observation (AO) is another cognitive motor process that shares common neural substrates with both MI and the actual motor preparation/execution (Tkach et al., 2007; Hardwick et al., 2018). AO and MI remain functionally dissociable and hierarchized with regards to the degree of involvement of the brain motor regions (Macuga and Frey, 2012). Nonetheless, AO consistently engages premotor, parietal and cerebellar networks. Brain networks recruited during AO have been described as part of a “*mirror neuron*” system (Rizzolatti and Craighero, 2004; Casile et al., 2011). Research in both humans and primates demonstrated that the mirror neuron system encodes relevant behavioural features of the observed actions (Agnew et al., 2007; Vogt and Thomaschke, 2007; Alaerts et al., 2012; Catmur, 2015). AO thus involves inputs to the brain motor system under the form of “(...) *Externally guided motor simulation*” (Vogt et al., 2013, p. 3). AO improves motor learning, specifically when the observer has a previous experience of the task (Rizzolatti et al., 1996; Bird et al., 2005; Mattar and Gribble, 2005; Calvo-Merino et al., 2006). However, few studies have investigated the efficacy of AO to improve strength. This is rather surprising considering the currently growing interest in mental training with AO to potentiate the recovery of muscle function (Pomeroy et al., 2005; Sale and Franceschini, 2012; Buccino, 2014). Porro et al. (2007) pioneered that daily training sessions with AO scheduled over the course of two consecutive weeks improved index finger abduction strength by 30%. The gains were specific to the movement observed during training. Noteworthy, the authors recorded increased excitability of corticospinal pathways targeting the first dorsal interosseous after training completion. These data support earlier findings by Ranganathan et al. (2004) suggesting that the statements of the psycho-neuromuscular theory of MI could also apply to training with AO.

MI and AO have been broadly conceptualized within the field of motor cognition as motor simulation processes due to their capacity to engage the brain motor system in the absence of physical execution (Jeannerod, 1994, 2001). Previously, Shepard (1984) suggested that MI and AO belonged to a continuum extending from passive/external (bottom-up) to active/internal (top-down) motor simulation (Vogt et al., 2013 for a more recent discussion). Consequently, the overlaps between MI and AO are readily exemplified in sporting situations. For instance, observing a teammate or an

opponent's performance concomitantly elicits the simulation of an adapted motor response. An emergent body of research investigates training with AO and MI concomitantly (AO+MI). AO+MI was referred to as “(...) *video guided imagery*” or “(...) *imagining imitation*” in the scientific literature. Eaves et al. (2016b) underlined that combining bottom-up and top-down processing through AO+MI yields synchronized visual and kinaesthetic inputs to the brain. Performing MI while observing the same movement on a video might thus facilitate the build-up of accurate motor representations by prompting attentional focus on the relevant spatial and temporal features of the coordination. Similarly, video cues provide direct inputs to the visual system and reduce the need for top-down processing of MI. The top-down processing of kinaesthetic information might consequently benefit from increased allocation of attentional resources. Transcranial magnetic stimulation provided evidence of higher corticospinal excitability during AO+MI compared to either AO or MI alone (Sakamoto et al., 2009; Ohno et al., 2011; Wright et al., 2014; Mouthon et al., 2015). Functional brain imaging recordings confirmed that AO+MI was associated with increased activity in brain motor regions compared to AO and MI alone (Nedelko et al., 2012). Due to greater involvement of motor regions in the brain, AO+MI might represent a more efficient way of enhancing the motor performance compared the isolated practice of AO and MI (Eaves et al., 2014; Taube et al., 2014; Bek et al., 2016).

Research on the effects of AO+MI on motor performance remains sparse. A limited number of experiments compared the effects of AO+MI to AO or MI *alone*. This is particularly valid when considering strength development, although recent studies have specifically compared the efficacy of AO+MI with that of MI practice alone. In a seminal study, Scott et al. (2018) reported greater effectiveness of AO+MI compared to MI alone to increase hamstring strength. The study design also included a MI-control condition (i.e. MI focused on the upper-limbs). The three-weeks of training involved 3 sessions of 20 min of MI training incorporating the “*Physical, Environmental, Task, Timing, Learning, Emotion Perspective*” (PETTLEP) guidelines (Holmes and Collins, 2001). The AO+MI and MI alone session focused on Nordic hamstring exercises, while the AO component of the AO+MI condition involved an external visual perspective. Eccentric hamstring peak torque increased

in the right leg in the AO+MI group (i.e. ~6% of increase compared to the baseline), whereas the ~3.5% increase in the MI alone group did not reach significance. Further analyses (magnitude-based inference) confirmed the potential superior relevance of AO+MI, specifically from a clinical perspective. Similarly, Smith et al. (2019) addressed whether the adjunct of AO to MI training improved the maximal strength on a biceps curl machine. The authors implemented a multiple-case counterbalanced design, where AO+MI and MI alone were each administered during four weeks after a three-week baseline period. Surprisingly, the authors found that MI and AO+MI were both equally efficient. Notably, AO did not focus on an external model, but was built from a video capture of participants' own physical performance, filmed from the first person perspective during a set of repetitions to failure during the baseline period. Although their findings conflicted with those reported by Scott et al. (2018), the authors did not rule out the hypothesis that AO+MI might outperform MI alone. This is primarily due to the trajectory of performance changes. When AO+MI took place before MI alone the authors systematically observed a performance increase along the course of the four weeks, whereas MI alone appeared to maintain the strength benefits. They also considered the possibility of delayed gains, where AO+MI would ultimately outperform MI alone if the intervention was prolonged up to 15 weeks (e.g. Wakefield and Smith, 2011). In keeping with these exploratory findings, we found no experiments that compared AO with AO+MI in a strength paradigm. Considering the limited amount of research on the development of maximal isometric strength and the growing interest in AO+MI, we designed the present study to investigate the short-term effects of AO and AO+MI on maximal isometric strength. Based on the psycho-neuromuscular theory, we hypothesized that AO+MI should outperform AO, while AO would outperform the control condition.

Material and Methods

Participants

Twenty-one participants were recruited from the Faculty of Sports Sciences of the University Claude Bernard Lyon 1 (F-69100, Villeurbanne; 17 males; mean age = 22.25 ± 2.18 years). All were athletes of regional to national level in terrestrial sports (judo, karate, climbing, basket and soccer),

and had a background of at least 2 years in resistance training (i.e. >1 year of practice, 2-4 sessions of 45 min per week). Participants were also screened based on their ability to engage in MI practice using a standardized MI questionnaire, i.e. average score >5 on the Movement Imagery Questionnaire-3 (Williams et al., 2012). The study was a relative strength contest with reference to body weight. This was expected to encourage the commitment to a maximal effort throughout experimental sessions. We did not provide any information regarding the purpose of the study until after completion of the design. The local Ethics Committee approved the study. Each participant provided written informed consent according to the statements of the Declaration of Helsinki (2013).

Experimental design

The repeated-measures design comprised three experimental sessions of 30 minutes, separated from each other by 48 hours. We interposed a no-training day between experimental sessions, which were scheduled at 12 am before lunch to control for the potential influence of circadian rhythms. Each experimental session consisted of 10 maximal elbow flexions against a fixated force platform. Participants attempted to lift the fixated force platform from a seated position with the elbow at 90° and the hand against the platform (Figure 1). For each trial, the voluntary maximal isometric contraction was sustained for 12 s. Each trial was separated from each other by a 60 s period allocated to recovery.

*** *Insert Figure 1 about here* ***

Experimental settings

Participants warmed-up for 10 min in a quiet room. They first completed 6 minutes of body weight exercises (i.e. lunges and squat jumps). Then, they seated on a bench equipped with a reclining seatback. They were instructed to keep permanent contact between the reclining seatback and the back of their head, the posterior apex of the thoracic spine, and the pelvis in order to standardize the trunk position (Figure 1A). Participants constantly fixed a cross mark on the wall at eye level. From this standardized position, each participant performed 5 consecutive isometric contractions of 12 s of incremental intensity against the fixated force platform. They were instructed to achieve a self-paced

increase in the intensity of the isometric contractions across the 5 trials, but below their maximal isometric strength threshold. Each trial was separated by recovery periods of 50 s. Participants then finished the warm-up by two maximal isometric contractions sustained for 12 s, separated by 90 s of passive recovery. Finally, the experimenter provided oral instructions corresponding to the upcoming experimental condition. While sitting on the bench, they faced a force platform under which they placed their dominant hand in supine position, with the elbow at 90°, as controlled by goniometers. We rigorously controlled the hand position at each trial, checking that the wrist, palm and fingers remained in flat contact with the force platform (Figure 1A).

Experimental conditions

Across the repeated measures of the design, we administered the following experimental conditions during the inter-trial recovery periods: i) observation of a bodybuilder athlete performing the task (AO), ii) observation of a bodybuilder athlete performing the task combined with MI of oneself performing the task (AO+MI) and iii) watching passively a video documentary about basketball shooting (CONTROL). Experimental settings were identical to those during the warm-up. During each 60 s inter-trial recovery period, we administered three blocks (15 s each) of mental practice corresponding to the different experimental conditions. The onset and the offset of each trial and recovery periods was externally cued by audio stimuli (Presentation, Neurobehavioral systems®, 240 Hz, 50 Db). To prevent carryover effects, e.g. residual muscle fatigue from one experimental session to another, experimental conditions were administered in counterbalanced order (block randomization).

During inter-trial recovery periods, participants watched a 15 s video displayed on a screen in front of them. During AO and AO+MI, the video displayed a bodybuilder athlete (79 kg, 7.63% of body fat, 22 years old) completing the maximal isometric strength task in identical experimental settings. The video showed the bodybuilder from a lateral view (Figure 1A). During AO, the experimenter gave participants the following instructions: *“Watch carefully the athlete performing the maximal voluntary contraction. Focus on the effort produced by the athlete trying to lift the platform. Focus your attention on his upper limb and observe the intense muscles contractions all along with the*

maximal effort intensity". The experimenter explicitly instructed participants not to engage in MI during the AO condition. This method is comparable to that described by Eaves et al. (2016a), where the authors requested: "*Please refrain from undertaking any MI during (...) observation in this pure AO condition*" (p. 94). During AO+MI, participants performed the same action observation task as during AO while simultaneously engaging in MI. When watching the video, the participants were instructed to: "*Imagine yourself in an attempt to lift the platform. Feel the intense contraction of your biceps and your shoulder muscles during the maximal effort. Focus on the total recruitment of muscle fibres throughout the duration of the effort*". Participants were prompted by such instructions to engage in the kinaesthetic modality of MI. During CONTROL, participants remained motionless and watched the video documentary for 15 s.

Dependent variables

Psychometric recordings

Functions of Observational Learning Questionnaire

Before the first experimental session, participants completed the Functions of Observational Learning Questionnaire (FOLQ; Cumming et al., 2005). This is a qualitative tool investigating participants' uses of action observation with the aim to improve performance in their sporting activities (Cumming et al., 2005). The FOLQ investigates the uses of AO with reference to the cognitive and motivational functions of MI earlier described by Paivio (1985) to improve sporting performance. Participants reported each item on a Likert-type scale ranging from 1 ("*I never use*") to 7 ("*I often use*"). The FOLQ specifically investigates two cognitive (Strategy and Skills) and one motivational (Performance) functions of AO. STRATEGY refers to uses of AO to improve cognitive processing related to tactical aspects mediating performance (e.g. "*I use observational learning to determine how a strategy will work in an event/game*"). SKILLS refers to uses of AO to improve cognitive processing related to technical execution mediating performance (e.g. "*I use observational learning to understand how to perfectly perform a skill*"). Finally, PERFORMANCE refers to uses of AO to achieve optimal arousal and anxiety appraisal in sporting situations (e.g. "*I use observational*

learning to know how to respond to the excitement associated with performing well”). The FOLQ showed high internal consistency with Cronbach’s alphas ranging from 0.84-0.90 (Cumming et al., 2005).

Likert ratings

After each experimental session, participants rated their perceived strength throughout the maximal isometric strength trials on a Likert-type scale ranging from 1 (“*Very low level of strength output*”) to 10 (“*Very high level of strength output*”). Participants also rated their motivation to complete the experimental session on a Likert-type scale ranging from 1 (“*Very low level of motivation*”) to 10 (“*Very high level of motivation*”). We also evaluated the perceived difficulty to complete the experimental session with self-reports using a Likert-type scale ranging from 1 (“*Very low level of difficulty*”) to 10 (“*Very high level of difficulty*”). After completion of the AO+MI condition, we finally evaluated MI vividness with self-reports with a Likert-type scale ranging from 1 (i.e. “*Absence of visual/kinaesthetic information during imagination*”) to 10 (i.e. “*Comparable visual/kinaesthetic information during imagination as during physical practice*”).

Behavioural recordings

We measured the elbow flexion strength with a force platform (AMTI, model 0R6-7-2000, Watertown, Massachusetts, USA). Data were continuously recorded and synchronized by LabChart Pro V8[®] (ADInstruments Pty Ltd, 2014) at 1000 Hz. Data were smoothed with a zero-lag 4th-order low-pass Butterworth filter with a 20 Hz cut-off frequency. During each trial, the sudden force increase in response to auditory stimulus was detected using a threshold detection function (Matlab[®]). We then calculated the total force by integrating the force slope with respect to the duration of each trial (12 s, trapezoid rules). We finally normalized the total force in percentage of the total force value recorded during the maximal isometric force trial of the warm-up:

$$Total\ Force_{(Normalized)} = \left(\frac{Total\ Force_{(Maximal\ isometric\ strength\ trial\ 1-10)}}{Total\ Force_{(Best\ maximal\ isometric\ strength\ during\ warm-up)}} \right) \times 100$$

The total force was used as the behavioural performance variable quantifying strength.

Physiological recordings

For each trial, we recorded the surface electromyograms (EMG) of the main agonists. After shaving and cleaning the skin with alcohol, we recorded EMG from the *biceps brachii* using pairs of surface electrodes (1 cm EMG Triode, nickel-plated brass, inter-electrode distance 2 cm, Thought Technology, Montreal, Canada). We also recorded EMG from the *anterior deltoideus*, a synergist muscle of the experimental task (Di Rienzo et al., 2015). Electrodes positioning was determined according to the recommendations of the “*Surface Electromyography for the Non-Invasive Assessment of Muscles*” (SENIAM) project (Hermens et al., 2000). Electrodes location was marked with a pen and photographed to ensure reproducible positioning across experimental sessions. We continuously recorded and synchronized EMG data by LabChart ProV8[®] (ADInstruments Pty Ltd, 2014). We processed the raw signal using the TrignoTM Wireless EMG[®] system (2014, Delsys Incorporated). We then rectified and smoothed the raw EMG signal with a 20-500 Hz pass-band filter (Butterworth 4th). For each trial, we calculated the integrated EMG (iEMG) with respect to the 12 s of each voluntary contraction (Figure 1B). We considered iEMG a global measurement of motor units activation related to isometric strength output (Moritani and deVries, 1978). We finally normalized iEMG data in percentage of the maximal iEMG value recorded during the maximal isometric force trial of the warm-up:

$$iEMG_{(Normalized)} = \left(\frac{iEMG_{(Maximal\ isometric\ strength\ trial\ 1-10)}}{iEMG_{(Best\ maximal\ isometric\ strength\ during\ warm-up)}} \right) \times 100$$

We continuously recorded electrodermal activity through skin conductance measures (micro Siemens, μS , i.e. $1 \cdot 10^{-6} \text{ m}^{-2} \cdot \text{kg} \cdot 1 \cdot \text{s}^3 \cdot \text{A}^2$), using two 50 mm² unpolarizable bipolar electrodes placed on the second phalanx of the second and third digits of the non-dominant hand (Figure 1; MLT116f GSR Finger Electrodes, ADInstruments, Dunedin, New Zealand). Electrodermal activity reflects the activity of eccrine sweats glands, which are under the unique control of the sympathetic branch of the autonomic nervous system (Shields et al., 1987). Increased skin conductance attests increased sympathetic activity, and inversely. After checking that electrodermal activity exhibited a similar pattern across experimental sessions, we collected the electrodermal response (EDR) for each maximal

isometric strength trial. We then calculated the EDR amplitude as the delta separating the sudden increase and the onset of the prolonged and regular return to the pre-stimulus baseline (for further development and experimental illustration, see Vernet-Maury et al., 1995; Kanthack et al., 2017). EDR amplitude is typically associated with the mental workload elicited by cognitive processing during both physical and mental tasks (Collet et al., 2013). Accordingly, we used EDR amplitudes as an objective measure of participants' commitment to a maximal effort during the isometric strength trials.

Statistical analysis

Linear mixed effects analysis

We used R (2018) and *nlme* (Pinheiro et al., 2014) to run a linear mixed-effects analysis with a by-subject random intercept of the psychometric, behavioural and physiological dependent variables. For FOLQ scores, we entered DIMENSION (i.e. SKILL, STRATEGY and PERFORMANCE) as fixed effect. We investigated self-reports of motivation and perceived performance, as well as self-reports of perceived difficulty to complete the cognitive tasks administered during the inter-trial recovery periods (i.e. AO, AO+MI), using CONDITION (i.e. CONTROL, AO, AO+MI) as fixed effect. For EDR amplitude, we built a linear mixed effect model with the fixed effect of CONDITION and TRIAL (with interaction term). For total force, we entered the interaction between CONDITION and TRIAL, $iEMG_{\text{BICEPS BRACHII}}$, $iEMG_{\text{ANTERIOR DELTOIDEUS}}$ as fixed effects. For all analyses, we included TRIAL as numeric regressor (i.e. ranging from 1 to 10). This procedure enabled qualitative investigation of linear trends (Di Rienzo et al., 2015, 2019; Kanthack et al., 2017 for comparable statistical analysis frameworks). We implemented a backward stepwise procedure to fit the random-coefficient regression model formulae (Hocking, 1976; Draper and Smith, 2014). Inspection of the residual plots did not reveal any obvious deviation from the hypotheses of homoscedasticity or normality. We set up the statistical significance threshold for a type 1 error rate of $\alpha = 5\%$. As effect sizes, we reported partial coefficients of determination (R_p^2), using the procedure for linear mixed effects models implemented in the *r2glmm* package (Edwards et al., 2008; Jaeger et al., 2017). We finally investigated main effects and interactions using general linear hypotheses testing of planned contrasts from the *multcomp*

package (Hothorn et al., 2008; Bretz et al., 2016). We applied Holm's sequential corrections to control the false discovery rate (Holm, 1979).

Power analysis

We carried out power calculations using the *pwr* package implemented in R (Champely et al., 2018). Due to the nature of the experimental hypotheses, sample size should afford a reliable statistical power to detect a main effect of the EXPERIMENTAL CONDITION on total force. Considering the repeated measures nature of the design, a sample size of $n = 21$ yielded a statistical power of $p_{(1-\beta)} = 0.80$ to detect small effect sizes, i.e. corresponding to 1-5% of explained variance with a type 1 error rate of 5%. The statistical power was superior to 0.90 for medium and large effect sizes, i.e. corresponding to 10% of explained variation and higher.

Results

Monitoring compliance to the experimental design

Psychometric data

FOLQ scores

The linear mixed effects analysis revealed a statistically significant effect of the DIMENSION ($\chi^2(2) = 85.66, p < 0.001, R_p^2 = 0.47$). FOLQ for PERFORMANCE scores ($M \pm SE; 2.85 \pm 0.25$) were lower than those recorded for both STRATEGY (4.33 ± 0.33) and SKILLS (5.41 ± 0.31) dimensions (fitted difference: $1.46 \pm 0.28, p < 0.001; 2.57 \pm 0.28, p < 0.001$; respectively). FOLQ scores for SKILL were also higher than those for STRATEGY (fitted difference: $1.08 \pm 0.28, p < 0.001$).

Self-reports on Likert-type scales

The linear mixed effects analysis on motivation and perceived performance yielded no CONDITION effect ($\chi^2(2) = 0.53, p = 0.77; \chi^2(2) = 1.35, p = 0.51$, respectively), as shown by Figure 2. The COGNITIVE TASK affected Likert ratings of perceived difficulty ($\chi^2(1) = 3.90, p = 0.04, R_p^2 = 0.06$ - Figure 2). Participants rated that it was more difficult to complete AO+MI (4.38 ± 0.51) than

AO alone (3.23 ± 0.65 , fitted difference: 1.14 ± 0.59 , $p = 0.04$). Finally, vividness ratings after AO+MI were 6.66 ± 1.83 on the 10-point Likert scale, i.e. corresponding to the “*Quite good (...)*” vividness item (Figure 2).

*** *Insert Figure 2 about here* ***

Skin conductance data

The analysis of EDR amplitude revealed that the CONDITION \times TRIAL interaction approached the statistical significance threshold ($\chi^2(2)$: 4.56, $p = 0.10$, $R_p^2 = 0.01$). As shown in Figure 3, there was a marginally negative TRIAL effect on EDR amplitudes during AO+MI (fitted estimate: $-0.05 \mu\text{S} \pm 0.06$, $p = 0.09$). By contrast, there was no TRIAL effect on EDR amplitudes recorded during both AO ($p = 0.74$) and CONTROL conditions ($p = 0.73$). The linear mixed effects analysis revealed no main effect of CONDITION ($\chi^2(2) = 3.79$, $p = 0.15$) or TRIAL ($\chi^2(1) = 1.36$, $p = 0.24$) (Figure 3).

*** *Insert Figure 3 about here* ***

Analysis of the maximal isometric strength

Raw group values used for normalization (see Materials and methods) of the total force, $i\text{EMG}_{\text{BICEPS BRACHII}}$ and $i\text{EMG}_{\text{ANTERIOR DELTOIDEUS}}$, are provided in Table 1.

The CONDITION \times TRIAL and CONDITION \times $i\text{EMG}_{\text{BICEPS BRACHII}}$ interactions were removed during the backward stepwise model selection ($\chi^2(2) = 1.91$, $p = 0.38$; $\chi^2(2) = 3.85$, $p = 0.15$, respectively). However, the total force was affected by the TRIAL \times $i\text{EMG}_{\text{DELTOIDEUS}}$ interaction ($\chi^2(2) = 7.90$, $p = 0.02$, $R_p^2 = 0.01$). As shown in Figure 4, there was a positive relationship between $i\text{EMG}_{\text{DELTOIDEUS}}$ and strength during AO (fitted estimate: $0.09\% \pm 0.02$, $p < 0.01$), but not during CONTROL (fitted estimate: $0.02\% \pm 0.02$, $p = 0.66$) nor AO+MI (fitted estimate: $0.00\% \pm 0.02$, $p = 0.99$).

*** *Insert Table 1 about here* ***

The total force analysis revealed a main effect of TRIAL ($\chi^2(1) = 90.42$, $p < 0.001$, $R_p^2 = 0.10$), $i\text{EMG}_{\text{BICEPS BRACHII}}$ and $i\text{EMG}_{\text{ANTERIOR DELTOIDEUS}}$ ($\chi^2(1) = 53.39$, $p < 0.001$, $R_p^2 = 0.12$; $\chi^2(1) = 3.66$, $p = 0.05$,

$R_p^2 = 0.001$; respectively). As shown in Figure 4, there was a negative relationship between TRIAL and strength (fitted estimate: $-0.94\% \pm 0.01$, $p < 0.001$). Conversely, the relationship between $iEMG_{\text{BICEPS BRACHII}}$, $iEMG_{\text{ANTERIOR DELTOIDEUS}}$ and the total force was positive (fitted estimate: $0.17\% \pm 0.02$, $p < 0.001$; fitted estimate: $0.04\% \pm 0.01$, $p = 0.03$, respectively). There was also a main CONDITION effect ($\chi^2(2) = 32.43$, $p < 0.001$, $R_p^2 = 0.02$). Total force recorded during AO+MI ($79.07\% \pm 19.58$) outperformed that recorded during CONTROL ($76.47\% \pm 20.59$; fitted difference: $3.68\% \pm 0.69$, $p < 0.001$). Likewise, the total force recorded during AO ($78.72\% \pm 23.17$) outperformed that recorded during CONTROL ($2.79\% \pm 0.75$ of fitted difference, $p < 0.001$). There was no difference between AO+MI and AO ($p = 0.44$, see Figure 4).

*** *Insert Figure 4 about here* ***

Discussion

The present experiment investigated the short-term effects of AO and AO+MI on maximal isometric strength. In a previous study, we demonstrated the efficacy of MI practice during the inter-trial recovery periods (Di Rienzo et al., 2015). The performance analysis first revealed a positive relationship between the maximal isometric strength and the activation of primary and secondary agonists of the elbow flexion. This corroborates the well-established relationship between isometric strength and the integrated surface electromyogram (e.g. Moritani and deVries, 1978). The *biceps brachii* activation accounted for 12% of the variance of the maximal isometric strength. This was expected considering the nature of the strength task (Serrau et al., 2012). We also found a positive relationship between the *anterior deltoideus* activation and strength, which also confirmed its assistance function in the present elbow flexion force task (Di Rienzo et al., 2015). Repeated maximal isometric strength efforts lead to a corollary increase in the difficulty to recruit motor units (Kroll, 1968; Babault et al., 2006; Camic et al., 2013). A decrease in strength occurred along with trials repetition, yielding to a 9.46% decrease in maximal isometric strength at the session level. This measure is congruent with the 10% of maximal isometric strength decrease reported by Di Rienzo et al. (2015) in a similar paradigm, and attests participant's commitment to a maximal isometric effort.

The nature of the experimental condition did not impact the decrease in strength across trials. Performing AO and AO+MI during inter-trial recovery periods did not attenuate the performance decrease across trials compared to CONTROL. This argues for the relative independence between the mental load associated with mental training and the accumulation of muscle fatigue (Rozand et al., 2014a, 2014b). However, both AO and AO+MI outperformed CONTROL in terms of maximal isometric strength output across all trials of the session. These results confirm a previous report from Di Rienzo et al. (2015), where MI practice yielded no effect on the trial-to-trial decrease in strength but enhanced the overall strength output by 2.1-3.5% at the session level (i.e. irrespective of the trial number). Here, short-term improvements during AO and AO+MI ranged from 2.8-3.7% compared to CONTROL. Albeit modest, these gains are of practical relevance considering ceiling effects associated with the voluntary repetition of maximal isometric contractions. The literature reported 10-30% improvements in strength after training programs scheduled within a span of several weeks including mental training (Ranganathan et al., 2004; Porro et al., 2007; Lebon et al., 2010; Yao et al., 2013). The levels of perceived performance and motivation to engage in the strength paradigm were similar across experimental conditions. Hence, the superiority of AO and AO+MI compared to CONTROL may not account for psychological factors. We also ruled out the effects of increased attentional focus on the dominant upper limb during inter-trial recovery periods. Indeed, the CONTROL condition also involved a visual exposure to upper limb movements. Skin conductance measures confirmed that the sympathetic activity during maximal isometric strength trials was comparable across conditions. We therefore postulate that there is neurophysiological origin to the short-term beneficial effects of AO and AO+MI on maximal isometric strength.

The psycho-neuromuscular theory hypothesizes that mental training improves muscle function by prompting neural excitability within the neural pathways mediating the forthcoming physical performance. This was advanced as a primary underlying mechanism to the beneficial effects of training with AO and MI on strength (Porro et al., 2007; Di Rienzo et al., 2015; Grosprêtre et al., 2018). Indeed, AO and AO+MI both involve low-threshold facilitation of the corticospinal pathways controlling the corresponding physical performance (Naish et al., 2014; Wright et al., 2014, 2016,

2018). We predicted corticospinal facilitation during the inter-trial recovery periods during AO and AO+MI, but not during CONTROL. In light of the psycho-neuromuscular theory, we suggest that the preliminary activation of the corticospinal pathways controlling the elbow flexion task during AO and AO+MI facilitated the forthcoming maximal voluntary isometric contractions. We thus posit that a priming effect would explain the short-term impact of AO and AO+MI on strength (for illustrations in AO experiments, see Obhi and Hogeveen, 2010; Salama et al., 2011). Priming refers to “[a] change in behaviour based on previous stimuli” (Stoykov and Madhavan, 2015, p. 33). At a neurophysiological level, priming originates from short-term plasticity yielding increased cortical gain over motor units (Stoykov and Madhavan, 2015; Stoykov et al., 2017). Priming-related plasticity may first facilitate *intramuscular* coordination, i.e. the capacity of the central nervous system to recruit motor units targeting agonist muscles (Fallentin et al., 1993; Farina et al., 2002; Taylor et al., 2003). Priming-related plasticity can also facilitate *intermuscular* coordination. It involves muscle synergies, e.g. between the *biceps brachii* and *anterior deltoideus*. Reciprocal inhibition represents another central component of intermuscular coordination (Baratta et al., 1988; Hautier et al., 2000; Simoneau et al., 2006). Although necessary for joint stabilization, residual activity in antagonist muscles can affect the development of strength (Hautier et al., 2000).

The strength task involved a primarily quantitative effort. There was thus a potentially confounding factor with regards to participants’ uses of AO, which rarely involved achieving physiological arousal goals, as reflected by the low FOLQ performance scores. AO of a bodybuilder could have placed too low demand on the motor system to elicit priming effects. The performance analysis did not confirm this statement since AO outperformed CONTROL. It can be objected that characterizing an effect of AO alone would require a control condition without upper-limb movements, e.g. watching a bodybuilder sitting passively in a chair. Such control condition would be relevant although it would not discriminate the reallocation of attentional resources to the upper-limb, which is in itself sufficient to elicit a facilitation of corticospinal pathways (Hiraoka et al., 2013), from task-specific motor simulation. Also, since we did not independently evaluate the effect of AO and MI it may be objected that the experimental design does not make possible to identify what was driving

the effects on performance during AO+MI. Considering the recent evidence demonstrating the beneficial effect of MI alone in a comparable design (Di Rienzo et al., 2015), the absence of difference between AO and AO+MI could indicate that MI has in fact no effect on performance. Rigorously disentangling this issue would require including a MI alone condition in the design. Albeit valid, this limitation is grounded in the theoretical claim that AO+MI represents a form of combination of AO and MI in the sense of summed cognitive processes. Theoretical framework rather emphasize a theory of AO+MI where AO+MI should be considered a unitary form of motor cognition that is relatively independent from AO and MI in the sense of:

“(...) *quasi-encapsulated sensorimotor streams, which could either merge or compete depending on their contents and potential usefulness for on-going action plans*” (Eaves et al., 2016b, p. 4).

An important implication is that AO+MI might leverage distinct neurophysiological mechanisms than the practice of AO or MI alone, specifically with regards to priming effects. AO+MI required a voluntary state of attentional focus on the proprioceptive information associated with *biceps brachii* and *anterior deltoideus* contractions, while concomitantly receiving external video cues. Participants were explicitly instructed to rehearse proprioceptive information related to the targeted muscles. This is known to elicit an effector-specific facilitation of the corresponding corticospinal pathways (Wright et al., 2014). In addition, in the video of the bodybuilder, the contraction of the *biceps brachii* was prevalent (Figure 1). This possibly placed a greater emphasis on this muscle during AO+MI (Calatayud et al., 2016; Eaves et al., 2016b). Accordingly, priming effects during AO+MI possibly favoured intramuscular rather than intermuscular coordination. By contrast, AO involved an attentional focus on the whole-limb coordination (Flanagan and Johansson, 2003; Stefan et al., 2008; Richardson et al., 2009), thus suggesting that AO primarily primed intermuscular coordination, and more specifically the synergy between the *biceps brachii* and *anterior deltoideus*. This postulate is congruent with the positive relationship recorded between maximal isometric strength and the integrated electromyograms from both the *biceps brachii* and *anterior deltoideus* during AO, whereas only the integrated electromyogram from the *biceps brachii* predicted the maximal isometric strength during AO+MI.

Contrary to Scott et al. (2018), who emphasized the clinical relevance of AO+MI compared to MI, we did not find any superiority of AO+MI compared to the training condition involving a single method of motor simulation (here AO). The present design examined the performance outcome at the single-session level, and involved the dominant upper-limb. Both aspects represent major differences with the methodological framework adopted by Scott et al. (2018). Also, it is possible that differences between AO+MI and AO would emerge in the case of longer intervention periods targeting the lower-limbs. Another qualitative aspect that might account for differences in the pattern of results is that Scott et al. (2018) trained their participants in the mental rehearsal of a functional task involving the targeted effectors (i.e. Nordic hamstring). By contrast, the mental training intervention embedded within the inter-trial recovery periods in the present study focused on the same task from which we measured the behavioural performance. Overall, the present experimental design was closer to the methodological framework adopted by Smith et al. (2019). These authors reported comparable strength improvements on an elbow flexion task after either AO+MI or MI training (administered in a counterbalanced order across several weeks). We observed comparable results at the single-session level, particularly regarding the benefits of AO+MI (albeit we used an external visual perspective for the AO component of the AO+MI intervention). The present findings provide further original insights to Smith et al. (2019)'s conclusions with regards to equivalent gains following AO+MI and MI alone. If AO+MI represent a unitary form of motor cognition, it is plausible that the effects on strength operate through qualitatively distinct neurophysiological processes than AO or MI alone. Albeit somehow speculative, this could explain why strength gains may not be dissociable from a quantitative analysis of performance. Finally, present strength gains after AO+MI training had reduced magnitude compared to the findings by Scott et al. (2018) and Smith et al. (2019), i.e. 2.5-3.5% increase compared to the control condition versus ~6% and ~15% increase, respectively. Aside from considerations related to the cortical representation of the somatic effectors targeted by mental training (see Yue and Cole, 1992 for pioneering insights), we suggest that the magnitude of strength gains could be primarily related to mental training duration, specifically when the sessions are separately administered from physical training across a span of several weeks.

Conceptually, AO might involve MI to some degree, hence making difficult to control for a strict practice of AO during the AO condition (Vogt et al., 2013; Scott et al., 2018). At this point, no method can provide objective means to control the absence of MI during AO, perhaps with the exception of fMRI. Unfortunately, fMRI is not adapted to ecological situations and revealed largely overlapping neural networks (e.g., Macuga and Frey, 2012). This limitation is inherent to any research on AO. Yet, an interesting methodological approach to address this issue in future designs would be asking participants to rate from a Likert-type scale their degree of MI experience during AO. Obviously, such scores would not represent objective proof that MI did not occur in the AO condition, but would provide some evidence of the amount of MI during AO. Here, the benefits reported for AO are unlikely to represent the concomitant effects of MI practice, for several reasons. First, participants received clear instructions not to engage MI during AO, and just focus attention on the upper limb of the bodybuilder. Therefore, if some MI occurred, it was probably not an explicit process. On the contrary, participants were explicitly instructed to engage in MI during the AO+MI condition. In other words, if some MI occurred during AO, it remains a different form of motor cognition than the deliberate MI practice instructed during AO+MI. In addition, the results advocate for distinct mechanisms underlying the benefits of AO and AO+MI condition, specifically about increased synergies under the AO condition. Eventually, the fact that participants reported that AO+MI was more difficult than AO potentially evidenced a different cognitive strategy under the two conditions. The priming effects of AO and AO+MI occurred without triggering awareness of increased efficacy compared to CONTROL, as attested by the Likert ratings. This finding somehow challenges data by Di Rienzo et al. (2015), where the participants perceived a greater strength during MI practice. MI involves a voluntary process of mental simulation, whereas AO emulates the motor system by engaging the mirror neuron system (Shepard, 1984; Vogt et al., 2013). In AO+MI, the participant primarily engaged in top-down processing of kinesthetic information since the video alleviates for the need of top-down visual processing by providing relevant afferent visual information. Yet, AO+MI required coordinating an internal and an external state of attentional focus. In addition, AO+MI required synchronising MI with the timing of the video. Overall, this could increase the mental load, presumably eliciting a conflict between the top-down and bottom-up processing of motor information

yielding to a negative bias in performance perception. This postulate is congruent with participants' higher perceived difficulty to perform AO+MI compared to AO as well as with the marginally negative relationship between the number of trials and the EDR amplitudes present during the AO+MI condition only. EDR amplitudes are strongly linked with attentional processes, and more specifically the allocation of mental resources to face the cognitive workload of the task (Collet et al., 2011, 2013). Overall, the increased perceived difficulty of AO+MI compared to AO and CONTROL was associated with autonomic nervous system response patterns attesting an increased mental fatigue state.

To the best of our knowledge, this is the first study investigating the short-term effectiveness of mental training including AO on maximal isometric strength. Previous research demonstrated the potential relevance of AO+MI to improve strength over the course of several weeks of training (Sun et al., 2016; Scott et al., 2018; Smith et al., 2019). The present study extends these findings and demonstrates that both AO and AO+MI increased strength at the single-session level. Increased training intensity at the session level contributes to enhance the workout load of longer training programs. Considering the importance of training intensity in strength development (Kraemer et al., 1996; Fleck and Kraemer, 2014), this may be an underlying process to the benefits of resistance training programs combining physical training with AO and AO+MI (Porro et al., 2007; Scott et al., 2018). Interestingly, AO and AO+MI facilitated strength without increasing participants' own performance perception. They even experienced a greater difficulty to perform AO+MI than AO, which was associated with decreased EDR amplitudes across trials. This result has important practical implications. Implementation of AO+MI during training might increase the mental workload for athletes while concomitantly contributing to increase their performances. While we assumed a neurophysiological origin to the beneficial effects of AO and AO+MI on maximal isometric strength through priming effects on corticospinal facilitation, this remains a working hypothesis considering the nature of the study design. Confirming this interpretation of the results would require replicating the design with direct measures of neural excitability within pathways targeting agonist and antagonist muscles. This could be implemented in future designs with transcranial magnetic stimulation and electroencephalographic measures.

Acknowledgments

This work was conducted within the facilities of the sport sciences of the University Claude Bernard Lyon 1. There was no funding from organizations for this work. The authors declare no competing interests.

References

- Agnew ZK, Bhakoo KK, Puri BK (2007) The human mirror system: a motor resonance theory of mind-reading. *Brain Res Rev* 54:286–293.
- Alaerts K, de Beukelaar TT, Swinnen SP, Wenderoth N (2012) Observing how others lift light or heavy objects: time-dependent encoding of grip force in the primary motor cortex. *Psychol Res* 76:503–513.
- Babault N, Desbrosses K, Fabre M-S, Michaut A, Pousson M (2006) Neuromuscular fatigue development during maximal concentric and isometric knee extensions. *J Appl Physiol Bethesda Md* 100:780–785.
- Bakker FC, Boschker MS, Chung T (1996) Changes in muscular activity while imagining weight lifting using stimulus or response propositions. *J Sport Exerc Psychol* 18:313–324.
- Baratta R, Solomonow M, Zhou BH, Letson D, Chuinard R, D'Ambrosia R (1988) Muscular coactivation. The role of the antagonist musculature in maintaining knee stability. *Am J Sports Med* 16:113–122.
- Bek J, Poliakoff E, Marshall H, Trueman S, Gowen E (2016) Enhancing voluntary imitation through attention and motor imagery. *Exp Brain Res* 234:1819–1828.
- Bird G, Osman M, Saggerson A, Heyes C (2005) Sequence learning by action, observation and action observation. *Br J Psychol Lond Engl* 96:371–388.
- Bretz F, Westfall P, Hothorn T (2016) Multiple comparisons using R. Chapman and Hall/CRC.
- Buccino G (2014) Action observation treatment: a novel tool in neurorehabilitation. *Philos Trans R Soc B Biol Sci* 369 Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4006186/> [Accessed September 12, 2018].
- Burianová H, Marstaller L, Sowman P, Tesan G, Rich AN, Williams M, Savage G, Johnson BW (2013) Multimodal functional imaging of motor imagery using a novel paradigm. *NeuroImage* 71:50–58.
- Calatayud J, Vinstrup J, Jakobsen MD, Sundstrup E, Brandt M, Jay K, Colado JC, Andersen LL (2016) Importance of mind-muscle connection during progressive resistance training. *Eur J Appl Physiol* 116:527–533.
- Calvo-Merino B, Grèzes J, Glaser DE, Passingham RE, Haggard P (2006) Seeing or doing? Influence of visual and motor familiarity in action observation. *Curr Biol* 16:1905–1910.
- Camic CL, Housh TJ, Zuniga JM, Russell Hendrix C, Bergstrom HC, Traylor DA, Schmidt RJ, Johnson GO (2013) Electromyographic and mechanomyographic responses across repeated maximal isometric and concentric muscle actions of the leg extensors. *J Electromyogr Kinesiol Off J Int Soc Electrophysiol Kinesiol* 23:342–348.

- Casile A, Caggiano V, Ferrari PF (2011) The mirror neuron system: a fresh view. *Neurosci Rev J Bringing Neurobiol Neurol Psychiatry* 17:524–538.
- Catmur C (2015) Understanding intentions from actions: Direct perception, inference, and the roles of mirror and mentalizing systems. *Conscious Cogn* 36:426–433.
- Champely S, Ekstrom C, Dalgaard P, Gill J, Weibelzahl S, Anandkumar A, Ford C, Volcic R, De Rosario H, De Rosario MH (2018) Package ‘pwr.’
- Clark BC, Mahato NK, Nakazawa M, Law TD, Thomas JS (2014) The power of the mind: the cortex as a critical determinant of muscle strength/weakness. *J Neurophysiol* 112:3219–3226.
- Collet C, Di Rienzo F, El Hoyek N, Guillot A (2013) Autonomic nervous system correlates in movement observation and motor imagery. *Front Hum Neurosci* 7 Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3726866/> [Accessed September 13, 2018].
- Collet C, Guillot A, Lebon F, MacIntyre T, Moran A (2011) Measuring motor imagery using psychometric, behavioral, and psychophysiological tools. *Exerc Sport Sci Rev* 39:85–92.
- Cornwall MW, Bruscatto MP, Barry S (1991) Effect of mental practice on isometric muscular strength. *J Orthop Sports Phys Ther* 13:231–234.
- Cumming J, Clark SE, Ste-Marie DM, McCullagh P, Hall C (2005) The functions of observational learning questionnaire (FOLQ). *Psychol Sport Exerc* 6:517–537.
- Damas F, Phillips S, Vechin FC, Ugrinowitsch C (2015) A review of resistance training-induced changes in skeletal muscle protein synthesis and their contribution to hypertrophy. *Sports Med Auckl NZ* 45:801–807.
- Di Rienzo F, Barlaam F, Daligault S, Delpuech C, Roy AC, Bertrand O, Jerbi K, Schmitz C (2019) Tracking the acquisition of anticipatory postural adjustments during a bimanual load-lifting task: A MEG study. *Hum Brain Mapp*.
- Di Rienzo F, Blache Y, Kanthack TFD, Monteil K, Collet C, Guillot A (2015) Short-term effects of integrated motor imagery practice on muscle activation and force performance. *Neuroscience* 305:146–156.
- Di Rienzo F, Debarnot U, Daligault S, Saruco E, Delpuech C, Doyon J, Collet C, Guillot A (2016) Online and Offline Performance Gains Following Motor Imagery Practice: A Comprehensive Review of Behavioral and Neuroimaging Studies. *Front Hum Neurosci* 10 Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4923126/> [Accessed September 11, 2018].
- Draper NR, Smith H (2014) *Applied regression analysis*. John Wiley & Sons.
- Eaves DL, Behmer LP, Vogt S (2016a) EEG and behavioural correlates of different forms of motor imagery during action observation in rhythmical actions. *Brain Cogn* 106:90–103.

- Eaves DL, Haythornthwaite L, Vogt S (2014) Motor imagery during action observation modulates automatic imitation effects in rhythmical actions. *Front Hum Neurosci* 8:28.
- Eaves DL, Riach M, Holmes PS, Wright DJ (2016b) Motor Imagery during Action Observation: A Brief Review of Evidence, Theory and Future Research Opportunities. *Front Neurosci* 10:514.
- Edwards LJ, Muller KE, Wolfinger RD, Qaqish BF, Schabenberger O (2008) An R2 Statistic for Fixed Effects in the Linear Mixed Model. *Stat Med* 27:6137–6157.
- Ehrsson HH, Geyer S, Naito E (2003) Imagery of voluntary movement of fingers, toes, and tongue activates corresponding body-part-specific motor representations. *J Neurophysiol* 90:3304–3316.
- Fallentin N, Jørgensen K, Simonsen EB (1993) Motor unit recruitment during prolonged isometric contractions. *Eur J Appl Physiol* 67:335–341.
- Farina D, Fosci M, Merletti R (2002) Motor unit recruitment strategies investigated by surface EMG variables. *J Appl Physiol Bethesda Md* 1985 92:235–247.
- Flanagan JR, Johansson RS (2003) Action plans used in action observation. *Nature* 424:769–771.
- Fleck SJ, Kraemer W (2014) *Designing Resistance Training Programs*, 4E. Human Kinetics.
- Gandevia SC, Wilson LR, Inglis JT, Burke D (1997) Mental rehearsal of motor tasks recruits alpha-motoneurons but fails to recruit human fusimotor neurons selectively. *J Physiol* 505 (Pt 1):259–266.
- Gao Q, Duan X, Chen H (2011) Evaluation of effective connectivity of motor areas during motor imagery and execution using conditional Granger causality. *NeuroImage* 54:1280–1288.
- Gerardin E, Sirigu A, Lehericy S, Poline JB, Gaymard B, Marsault C, Agid Y, Le Bihan D (2000) Partially overlapping neural networks for real and imagined hand movements. *Cereb Cortex N Y N* 1991 10:1093–1104.
- Grosprêtre S, Jacquet T, Lebon F, Papaxanthis C, Martin A (2018) Neural mechanisms of strength increase after one-week motor imagery training. *Eur J Sport Sci* 18:209–218.
- Grosprêtre S, Ruffino C, Lebon F (2016) Motor imagery and cortico-spinal excitability: A review. *Eur J Sport Sci* 16:317–324.
- Guillot A, Lebon F, Rouffet D, Champely S, Doyon J, Collet C (2007) Muscular responses during motor imagery as a function of muscle contraction types. *Int J Psychophysiol Off J Int Organ Psychophysiol* 66:18–27.
- Hanakawa T, Immisch I, Toma K, Dimyan MA, Van Gelderen P, Hallett M (2003) Functional properties of brain areas associated with motor execution and imagery. *J Neurophysiol* 89:989–1002.

- Hardwick RM, Caspers S, Eickhoff SB, Swinnen SP (2018) Neural correlates of action: Comparing meta-analyses of imagery, observation, and execution. *Neurosci Biobehav Rev* 94:31–44.
- Hautier CA, Arsac LM, Deghdegh K, Souquet J, Belli A, Lacour JR (2000) Influence of fatigue on EMG/force ratio and cocontraction in cycling. *Med Sci Sports Exerc* 32:839–843.
- Hermens HJ, Freriks B, Disselhorst-Klug C, Rau G (2000) Development of recommendations for SEMG sensors and sensor placement procedures. *J Electromyogr Kinesiol Off J Int Soc Electrophysiol Kinesiol* 10:361–374.
- Héту S, Grégoire M, Saimpont A, Coll M-P, Eugène F, Michon P-E, Jackson PL (2013) The neural network of motor imagery: an ALE meta-analysis. *Neurosci Biobehav Rev* 37:930–949.
- Hiraoka K, Mori N, Horino H (2013) Immediate effect of visual attention on corticospinal excitability in the upper trapezius muscle. *Percept Mot Skills* 117:211–214.
- Hocking RR (1976) A Biometrics invited paper. The analysis and selection of variables in linear regression. *Biometrics* 32:1–49.
- Holm S (1979) A simple sequentially rejective multiple test procedure. *Scand J Stat*:65–70.
- Holmes PS, Collins DJ (2001) The PETTLEP approach to motor imagery: A functional equivalence model for sport psychologists. *J Appl Sport Psychol* 13:60–83.
- Hothorn T, Bretz F, Westfall P (2008) Simultaneous inference in general parametric models. *Biom J Biom Z* 50:346–363.
- Jacobson E (1932) Electrophysiology of mental activities. *Am J Psychol* 44:677–694.
- Jaeger BC, Edwards LJ, Das K, Sen PK (2017) An R2 statistic for fixed effects in the generalized linear mixed model. *J Appl Stat* 44:1086–1105.
- Jeannerod M (1994) The representing brain: Neural correlates of motor intention and imagery. *Behav Brain Sci* 17:187–202.
- Jeannerod M (2001) Neural simulation of action: a unifying mechanism for motor cognition. *NeuroImage* 14:S103-109.
- Jowdy DP, Harris DV (1990) Muscular responses during mental imagery as a function of motor skill level. *J Sport Exerc Psychol* 12:191–201.
- Kanthack TFD, Guillot A, Papaxanthis C, Guizard T, Collet C, Di Rienzo F (2017) Neurophysiological insights on flexibility improvements through motor imagery. *Behav Brain Res* 331:159–168.
- Komi PV (1986) Training of muscle strength and power: interaction of neuromotoric, hypertrophic, and mechanical factors. *Int J Sports Med* 7 Suppl 1:10–15.

- Kraemer WJ, Fleck SJ, Evans WJ (1996) Strength and power training: physiological mechanisms of adaptation. *Exerc Sport Sci Rev* 24:363–397.
- Kroll W (1968) Isometric fatigue curves under varied intertrial recuperation periods. *Res Q* 39:106–115.
- Lebon F, Collet C, Guillot A (2010) Benefits of motor imagery training on muscle strength. *J Strength Cond Res* 24:1680–1687.
- Lebon F, Rouffet D, Collet C, Guillot A (2008) Modulation of EMG power spectrum frequency during motor imagery. *Neurosci Lett* 435:181–185.
- Lotze M, Halsband U (2006) Motor imagery. *J Physiol Paris* 99:386–395.
- Macuga KL, Frey SH (2012) Neural representations involved in observed, imagined, and imitated actions are dissociable and hierarchically organized. *NeuroImage* 59:2798–2807.
- Mattar AAG, Gribble PL (2005) Motor learning by observing. *Neuron* 46:153–160.
- Moritani T (1993) Neuromuscular adaptations during the acquisition of muscle strength, power and motor tasks. *J Biomech* 26 Suppl 1:95–107.
- Moritani T, deVries HA (1978) Reexamination of the relationship between the surface integrated electromyogram (IEMG) and force of isometric contraction. *Am J Phys Med* 57:263–277.
- Mouthon A, Ruffieux J, Wälchli M, Keller M, Taube W (2015) Task-dependent changes of corticospinal excitability during observation and motor imagery of balance tasks. *Neuroscience* 303:535–543.
- Munzert J, Zentgraf K (2009) Motor imagery and its implications for understanding the motor system. *Prog Brain Res* 174:219–229.
- Naish KR, Houston-Price C, Bremner AJ, Holmes NP (2014) Effects of action observation on corticospinal excitability: Muscle specificity, direction, and timing of the mirror response. *Neuropsychologia* 64:331–348.
- Nedelko V, Hassa T, Hamzei F, Schoenfeld MA, Dettmers C (2012) Action imagery combined with action observation activates more corticomotor regions than action observation alone. *J Neurol Phys Ther JNPT* 36:182–188.
- Obhi SS, Hogeveen J (2010) Incidental action observation modulates muscle activity. *Exp Brain Res* 203:427–435.
- Ohno K, Higashi T, Sugawara K, Ogahara K, Funase K, Kasai T (2011) Excitability changes in the human primary motor cortex during observation with motor imagery of chopstick use. *J Phys Ther Sci* 23:703–706.
- Paivio A (1985) Cognitive and motivational functions of imagery in human performance. *Can J Appl Sport Sci J Can Sci Appl Au Sport* 10:22S–28S.

- Paravlic AH, Slimani M, Tod D, Marusic U, Milanovic Z, Pisot R (2018) Effects and Dose-Response Relationships of Motor Imagery Practice on Strength Development in Healthy Adult Populations: a Systematic Review and Meta-analysis. *Sports Med Auckl NZ* 48:1165–1187.
- Peterson MD, Rhea MR, Alvar BA (2004) Maximizing strength development in athletes: a meta-analysis to determine the dose-response relationship. *J Strength Cond Res* 18:377–382.
- Pinheiro J, Bates D, DebRoy S, Sarkar D (2014) Linear and nonlinear mixed effects models. R Package Version 3.
- Pomeroy VM, Clark CA, Miller JSG, Baron J-C, Markus HS, Tallis RC (2005) The potential for utilizing the “mirror neurone system” to enhance recovery of the severely affected upper limb early after stroke: a review and hypothesis. *Neurorehabil Neural Repair* 19:4–13.
- Porro CA, Facchin P, Fusi S, Dri G, Fadiga L (2007) Enhancement of force after action observation: behavioural and neurophysiological studies. *Neuropsychologia* 45:3114–3121.
- Ranganathan VK, Siemionow V, Liu JZ, Sahgal V, Yue GH (2004) From mental power to muscle power--gaining strength by using the mind. *Neuropsychologia* 42:944–956.
- Reiser M, Büsch D, Munzert J (2011) Strength gains by motor imagery with different ratios of physical to mental practice. *Front Psychol* 2:194.
- Rhea MR, Alvar BA, Burkett LN, Ball SD (2003) A meta-analysis to determine the dose response for strength development. *Med Sci Sports Exerc* 35:456–464.
- Richardson MJ, Campbell WL, Schmidt RC (2009) Movement interference during action observation as emergent coordination. *Neurosci Lett* 449:117–122.
- Rizzolatti G, Craighero L (2004) The mirror-neuron system. *Annu Rev Neurosci* 27:169–192.
- Rizzolatti G, Fadiga L, Gallese V, Fogassi L (1996) Premotor cortex and the recognition of motor actions. *Brain Res Cogn Brain Res* 3:131–141.
- Rozand V, Lebon F, Papaxanthis C, Lepers R (2014a) Does a mental training session induce neuromuscular fatigue? *Med Sci Sports Exerc* 46:1981–1989.
- Rozand V, Pageaux B, Marcora SM, Papaxanthis C, Lepers R (2014b) Does mental exertion alter maximal muscle activation? *Front Hum Neurosci* 8:755.
- Sakamoto M, Muraoka T, Mizuguchi N, Kanosue K (2009) Combining observation and imagery of an action enhances human corticospinal excitability. *Neurosci Res* 65:23–27.
- Salama IM, Turner S, Edwards MG (2011) Automatic priming of grip force following action observation. *Q J Exp Psychol* 2006 64:833–838.
- Sale DG (1988) Neural adaptation to resistance training. *Med Sci Sports Exerc* 20:S135-145.

- Sale P, Franceschini M (2012) Action observation and mirror neuron network: a tool for motor stroke rehabilitation. *Eur J Phys Rehabil Med* 48:313–318.
- Schoenfeld BJ (2010) The mechanisms of muscle hypertrophy and their application to resistance training. *J Strength Cond Res* 24:2857–2872.
- Scott M, Taylor S, Chesterton P, Vogt S, Eaves DL (2018) Motor imagery during action observation increases eccentric hamstring force: an acute non-physical intervention. *Disabil Rehabil* 40:1443–1451.
- Serrau V, Driss T, Vandewalle H, Behm DG, Lesne-Chabran E, Le Pellec-Muller A (2012) Muscle activation of the elbow flexor and extensor muscles during self-resistance exercises: comparison of unilateral maximal cocontraction and bilateral self-resistance. *J Strength Cond Res* 26:2468–2477.
- Shepard RN (1984) Ecological constraints on internal representation: resonant kinematics of perceiving, imagining, thinking, and dreaming. *Psychol Rev* 91:417–447.
- Shields SA, MacDowell KA, Fairchild SB, Campbell ML (1987) Is mediation of sweating cholinergic, adrenergic, or both? A comment on the literature. *Psychophysiology* 24:312–319.
- Simoneau E, Martin A, Porter MM, Van Hoecke J (2006) Strength training in old age: adaptation of antagonist muscles at the ankle joint. *Muscle Nerve* 33:546–555.
- Smith D, Romano-Smith S, Wright DJ, Deller-Rust B, Wakefield CJ (2019) The Effects of Combining PETTLEP Imagery and Action Observation on Bicep Strength: A Single-Case Design. *J Appl Sport Psychol*:1–15.
- Solodkin A, Hlustik P, Chen EE, Small SL (2004) Fine modulation in network activation during motor execution and motor imagery. *Cereb Cortex N Y N 1991* 14:1246–1255.
- Stefan K, Classen J, Celnik P, Cohen LG (2008) Concurrent action observation modulates practice-induced motor memory formation. *Eur J Neurosci* 27:730–738.
- Stephan KM, Frackowiak RS (1996) Motor imagery--anatomical representation and electrophysiological characteristics. *Neurochem Res* 21:1105–1116.
- Stinear C (2010) Corticospinal facilitation during motor imagery. In: *The Neurophysiological Foundations of Mental and Motor Imagery*, pp 47–62.
- Stoykov ME, Corcos DM, Madhavan S (2017) Movement-Based Priming: Clinical Applications and Neural Mechanisms. *J Mot Behav* 49:88–97.
- Stoykov ME, Madhavan S (2015) Motor priming in neurorehabilitation. *J Neurol Phys Ther JNPT* 39:33–42.
- Sun Y, Wei W, Luo Z, Gan H, Hu X (2016) Improving motor imagery practice with synchronous action observation in stroke patients. *Top Stroke Rehabil* 23:245–253.
- Szameitat AJ, Shen S, Sterr A (2007) Motor imagery of complex everyday movements. An fMRI study. *NeuroImage* 34:702–713.

- Taube W, Lorch M, Zeiter S, Keller M (2014) Non-physical practice improves task performance in an unstable, perturbed environment: motor imagery and observational balance training. *Front Hum Neurosci* 8:972.
- Taylor AM, Christou EA, Enoka RM (2003) Multiple features of motor-unit activity influence force fluctuations during isometric contractions. *J Neurophysiol* 90:1350–1361.
- Team RC (2018) R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Austria, 2015. ISBN 3-900051-07-0: URL <http://www.R-project.org>.
- Tkach D, Reimer J, Hatsopoulos NG (2007) Congruent activity during action and action observation in motor cortex. *J Neurosci Off J Soc Neurosci* 27:13241–13250.
- Tod D, Edwards C, McGuigan M, Lovell G (2015) A Systematic Review of the Effect of Cognitive Strategies on Strength Performance. *Sports Med Auckl NZ* 45:1589–1602.
- Tod D, Iredale F, Gill N (2003) “Psyching-up” and muscular force production. *Sports Med Auckl NZ* 33:47–58.
- Vernet-Maury E, Robin O, Dittmar A (1995) The ohmic perturbation duration, an original temporal index to quantify electrodermal responses. *Behav Brain Res* 67:103–107.
- Vogt S, Di Rienzo F, Collet C, Collins A, Guillot A (2013) Multiple roles of motor imagery during action observation. *Front Hum Neurosci* 7 Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3839009/> [Accessed September 12, 2018].
- Vogt S, Thomaschke R (2007) From visuo-motor interactions to imitation learning: behavioural and brain imaging studies. *J Sports Sci* 25:497–517.
- Wagner J, Stephan T, Kalla R, Brückmann H, Strupp M, Brandt T, Jahn K (2008) Mind the bend: cerebral activations associated with mental imagery of walking along a curved path. *Exp Brain Res* 191:247–255.
- Wakefield C, Smith D (2011) From strength to strength: A single-case design study of PETTLEP imagery frequency. *Sport Psychol* 25:305–320.
- Williams SE, Cumming J, Ntoumanis N, Nordin-Bates SM, Ramsey R, Hall C (2012) Further validation and development of the movement imagery questionnaire. *J Sport Exerc Psychol* 34:621–646.
- Wright DJ, McCormick SA, Williams J, Holmes PS (2016) Viewing Instructions Accompanying Action Observation Modulate Corticospinal Excitability. *Front Hum Neurosci* 10 Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4740958/> [Accessed September 13, 2018].
- Wright DJ, Williams J, Holmes PS (2014) Combined action observation and imagery facilitates corticospinal excitability. *Front Hum Neurosci* 8:951.

- Wright DJ, Wood G, Eaves DL, Bruton AM, Frank C, Franklin ZC (2018) Corticospinal excitability is facilitated by combined action observation and motor imagery of a basketball free throw. *Psychol Sport Exerc* 39:114–121.
- Yao WX, Ranganathan VK, Alexandre D, Siemionow V, Yue GH (2013) Kinesthetic imagery training of forceful muscle contractions increases brain signal and muscle strength. *Front Hum Neurosci* 7 Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3783980/> [Accessed September 11, 2018].
- Yue G, Cole KJ (1992) Strength increases from the motor program: comparison of training with maximal voluntary and imagined muscle contractions. *J Neurophysiol* 67:1114–1123.

Figures caption


Figure 1. Experimental session settings. A: Experimental settings involved a standardized body position in front of a fixed force platform against which participants produced a maximal isometric contraction with the elbow at 90° (i.e. vertical arrow). The total force was used as the behavioural performance variable quantifying strength. B: Electromyograms from the *biceps brachii* and *anterior deltoideus* of the dominant upper limb. The smoothed/rectified signal corresponding to the 12 s of each trial was integrated and used as a dependent variable quantifying voluntary muscle activation. C: Skin conductance, with sensors on the second phalanx of the second and third digits, according to traditional recommendations. We measured the maximal amplitude of skin conductance response during each 12s-trial, as an index of the sympathetic activity.

Figure 2. Barplots with 95% confidence intervals (error bars) of self-reports data on the 10-points Likert scale used for psychometric measures. A: Perceived motivation across experimental session. B: Perceived strength during the experimental session. C: Perceived difficulty during AO and AO+MI. D: Perceived vividness during AO+MI. NS: Not statistically significant. *: $p < 0.05$.


Figure 3. Linear mixed effects analysis of EDR amplitude. A: EDR amplitude by TRIAL regression slopes during AO, CONTROL and AO+MI. Regression slopes are represented with 95% confidence interval (dotted lines). B: Barplot of the main CONDITION (upper panel) and TRIAL (lower panel) effects. Estimates are represented with 95% confidence intervals (error bars). NS: Not statistically significant. φ : $p < 0.10$. Figure 4. Linear mixed effects analysis of the total force. A: Total force by $iEMG_{ANTERIOR\ DELTOIDEUS}$ regression slopes during AO, CONTROL and AO+MI. Regression slopes are represented with 95% confidence interval (dotted lines). B: Regression slope between the total force and $iEMG_{BICEPS\ BRACHII}$, irrespective of the CONDITION, represented with 95% confidence interval (dotted lines). C: Barplot of the total force estimates during AO, CONTROL and AO+MI, represented with 95% confidence interval (error bars). D: Barplot of the total force across TRIAL, represented with 95% confidence interval (error bars). NS: Not statistically significant. ***: $p < 0.001$. **: $p < 0.01$.

Figure 4. Linear mixed effects analysis of the total force. A: Total force by iEMG_{ANTERIOR DELTOIDEUS} regression slopes ($\beta_i \pm SE$) during AO (0.09 ± 0.05), CONTROL (0.02 ± 0.02) and AO+MI (0.00 ± 0.02). Regression slopes are represented with 95% confidence interval (dotted lines). B: Regression slope ($\beta_i \pm SE$) between the total force and iEMG_{BICEPS BRACHII} (0.17 ± 0.02), irrespective of the CONDITION, represented with 95% confidence interval (dotted lines). C. Barplot of the estimates for the main CONDITION effect represented with 95% confidence interval (error bars). D. Barplot of main TRIAL effect ($\beta_i \pm SE$) on total force (-0.94 ± 0.10), represented with 95% confidence interval (error bars). NS: Not statistically significant. ***: $p < 0.001$. **: $p < 0.01$.



A




B




C




Motivation




Perceived Performance






Perceived Difficulty



Motor Imagery Vividness







Total force		iEMG _{Biceps brachii}		iEMG _{Anterior deltoideus}	
Mean	95% CI	Mean	95% CI	Mean	95% CI
2617.34 mV.s ⁻¹	2276.83 - 2957.85	7803.91 mV.s ⁻¹	6063.21 - 9544.61	2548.14 mV.s ⁻¹	1546.65 - 3549.62

Table 1. Group averages and 95 % confidence intervals for raw total force, iEMG_{BICEPS BRACHII} and iEMG_{ANTERIOR DELTOIDEUS} corresponding to the best warm-up trial. iEMG: Integrated electromyogram. 95% CI: 95 % confidence interval.