



HAL
open science

New process observed in collisions between highly charged protonated protein and $\text{Xe}8+$, $\text{Xe}5+$, $\text{He}2+$ ions

Serge Martin, L. Chen, Richard Bredy, Arnaud Vernier, Philippe Dugourd, Rodolphe Antoine, C. Ortega, Min Ji, Jérôme Bernard, O. G. Maganad, et al.

► To cite this version:

Serge Martin, L. Chen, Richard Bredy, Arnaud Vernier, Philippe Dugourd, et al.. New process observed in collisions between highly charged protonated protein and $\text{Xe}8+$, $\text{Xe}5+$, $\text{He}2+$ ions. *Journal of Physics: Conference Series*, 2014, 488 (10), pp.102004. 10.1088/1742-6596/488/10/102004 . hal-02309442

HAL Id: hal-02309442

<https://univ-lyon1.hal.science/hal-02309442v1>

Submitted on 5 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

OPEN ACCESS

New process observed in collisions between highly charged protonated protein and Xe^{8+} Xe^{5+} He^{2+} ions

To cite this article: S Martin *et al* 2014 *J. Phys.: Conf. Ser.* **488** 102004

View the [article online](#) for updates and enhancements.

Related content

- [Introduction to Pharmaceutical Biotechnology, Volume 2: Protein and enzyme engineering](#)
S Bhatia
- [Some Critical Questions in Biological Physics: How dynein works and other motor proteins](#)
T A Waigh
- [Electron Kinetic Energies in Multicapture Processes in \$\text{Xe}^{30+}\$ - \$\text{C}_{60}\$ Collisions](#)
R Brédy, L Chen, S Martin et al.



IOP | ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

New process observed in collisions between highly charged protonated protein and Xe^{8+} , Xe^{5+} , He^{2+} ions

S. Martin ^{*} ¹, L. Chen ^{*} ², R. Brédy ^{*}, A. Vernier^{*}, P. Dugourd^{*}, R. Antoine^{*}, C. Ortéga^{*}, M. Ji^{*}, J. Bernard^{*}, O. Gonzalez Maganad [†], G. Reitsma[†] and T. Schlathölter

^{*}Institut Lumière Matière, UMR5306 Université Lyon 1-CNRS, Université de Lyon, 69622 Villeurbanne cedex, France

[†]KVI Atomic and Molecular Physics, University of Groningen, Zernikelaan 25, NL-9747 AA, The Netherlands

Synopsis Electron multicapture processes have been studied in collisions between cytochrome C protein and highly charged Xe and He projectile at keV kinetic energy range. In competition with single and double electron capture, a new and unexpected channel attributed to deprotonation process of the protein has been observed.

Studies of processes in ions-ions collisions are not common in the keV energy range. We have investigated the interaction between highly charged Xe^{8+} , Xe^{5+} or He^{2+} projectiles and trapped highly protonated cytochrome C (Cyt-C) using a new set-up developed at the KVI laboratory [1]. Single electron capture SEC and double electron capture DEC processes have been studied versus the charge q of the protein Cyt-C^{q+} ($q=15-19$). Typical mass spectra are shown on figure 1 for collisions between Xe^{8+} and Cyt-C^{q+} . At first sight, collisions between selected charge state of Cyt-C and projectile involve very few processes. As an example for the trapped Cyt-C^{18+} parent ion (figure 1d) the expected Cyt-C^{19+} and Cyt-C^{20+} peaks are mainly attributed to the single (SEC) and double electron capture (DEC) processes. The unexpected Cyt-C^{17+} peak is attributed tentatively to a deprotonation (DP) process of the Cyt-C parent ions. This new and unexpected channel has never been observed for other smaller biological molecules. This DP process increases from Cyt-C^{15+} to Cyt-C^{19+} . A tentative explanation is based on the fast decreasing of proton affinities versus the charge of Cyt-C. During the collision process; the electric field of the projectile is high enough to eject the proton from the Cyt-C. Higher mass resolution spectra and kinetic energy dependence would be necessary to confirm this mechanism.




Figure 1. Mass spectra of $\text{Xe}^{8+} + \text{Cyt-C}^{q+}$ ($q=15-19$). SEC, DEC and DP processes are observed.

References

- [1] S. Bari, R. Hoekstra and T. Schlathölter
2010 *Phys. Chem. Chem. Phys.* **12** 3376

¹ E-mail: smartin@univ-lyon1.fr

² E-mail: chen@univ-lyon1.fr

