



HAL
open science

Time Evolution of the internal energy distribution of molecules studied in an electrostatic storage ring, the Mini-Ring

Li Chen, C. Ortega, Min Ji, Richard Bredy, Jérôme Bernard, Bruno Concina, Guillaume Montagne, C. Joblin, A. Cassimi, Y. Ngonon-Ravache, et al.

► **To cite this version:**

Li Chen, C. Ortega, Min Ji, Richard Bredy, Jérôme Bernard, et al.. Time Evolution of the internal energy distribution of molecules studied in an electrostatic storage ring, the Mini-Ring. *Journal of Physics: Conference Series*, 2014, 488 (10), pp.102011. 10.1088/1742-6596/488/10/102011. hal-02311250

HAL Id: hal-02311250

<https://univ-lyon1.hal.science/hal-02311250>

Submitted on 5 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

OPEN ACCESS

Time Evolution of the internal energy distribution of molecules studied in an electrostatic storage ring, the Mini-Ring

To cite this article: L Chen *et al* 2014 *J. Phys.: Conf. Ser.* **488** 102011

View the [article online](#) for updates and enhancements.

Related content

- [Optical Nanomanipulation: Optically induced mechanical forces](#)
D L Andrews and D S Bradshaw
- [Time evolution of internal energy distribution of Anthracene studied in an electrostatic storage ring, the Mini-Ring](#)
L Chen, M Ji, J Bernard *et al.*
- [PAH radiative cooling and fragmentation kinematics studied within an electrostatic ring](#)
R Brédy, C Ortéga, M Ji *et al.*



IOP | ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

Time Evolution of the internal energy distribution of molecules studied in an electrostatic storage ring, the Mini-Ring

L. Chen*¹, C. Ortéga*, M. Ji*, R. Brédy*, J. Bernard*, B. Concina*, G. Montagne*, C. Joblin[†], A. Cassimi[‡], Y. Ngono-ravache[‡] and S. Martin*²

* Institut Lumière Matière, UMR5306 Université Lyon 1-CNRS, Université de Lyon, 69622 Villeurbanne cedex, France
[†] Université de Toulouse, UPS-OMP, IRAP, Toulouse, France
[‡] CIMAP, GANIL Université de CAEN, Bd. H. Becquerel, Caen, France

Synopsis We report on studies about time evolution of internal energy of polycyclic aromatic hydrocarbon PAH and fullerene in a small electrostatic storage device. Laser heating at different time delays was used as the probe to determine the energy distributions.

We present here the latest development around our compact electrostatic ion storage ring, so called the Mini-Ring[1].

The detection system has been recently upgraded with a position sensitive detector (PSD) for recording the position of the neutral spot exiting the ring at each turn. In recent experiments, this PSD was used to measure the Kinetic Energy Release (KER) of fragmentation of anthracene cations.

A significant reduction of the Betatron oscillations has been accomplished by re-optimizing the potentials of the Mini-Ring's deflectors, lens and reflectors. Major improvements were obtained by taking pictures of the Ar⁺ beam trajectory using a CCD camera (CANON 5Dmark2) equipped with 16-35 optical system placed in front of the window at the top of the experiment (Figure 1). In the past, one night was needed to take a picture of the 5 keV Argon beam (200nA) with an argentic camera. With the CCD camera, it takes 10s only to record a picture, which is good enough to optimize visualizing online the trajectory of the beam. These improvements were also possible due to a higher kinetic energy of the beam (12 keV) and therefore higher beam intensity up to 1.5μA. This fine tuning of the stored ions trajectories improved the overlap between the ion and LASER beams and reduced considerably the oscillations observed on the decay measurements.

The actual limitation of the storage time is due to collisions with the background gas (the pressure was 2·10⁻⁹ mbars at the best conditions). We have stored ions with a rather large kinetic energy range from 2 q keV to 12 q keV.

An ECR Nanogan ion source has been used to produce beams of cations or anions. Up to now beams of He⁺, F⁺, F⁻, Ar⁺, SF_n⁺ (n=1-5), SF₆⁻ as well as

PAH and fullerene have been stored. Intense beams of molecular cations and dications have been obtained using a 10 GHz very low HF power (less than 0.5 W). Results concerning the evolution of the internal energy distribution will be presented for different PAHs. Cooling rates have been measured at different internal energies. The high values of cooling rates found for Anthracene cations are explained by a cooling process due to the “fluorescence of the thermally excited electrons” predicted 20 years ago [2].



Figure 1. Picture of Mini-Ring. The light emitted along the Ar⁺ beam trajectory is due to collisions with nitrogen gas injected into the vacuum chamber at pressure of about 10⁻⁴ mbar. The blue light is due to the excitation of the N₂ molecules. The MCP imaging detector is placed at bottom left. A small spot due to neutralized Ar is well observed on the MCP.

References

- [1] S. Martin *et al.* 2013 *Phys. Rev. Lett.* **110** 063003
- [2] A. Léger, P. Boissel, and L. d'Hendecourt 1998 *Phys. Rev. Lett.* **60** 921

E-mail: smartin@univ-lyon1.fr

E-mail: chen@univ-lyon1.fr

