



HAL
open science

The responses of NO₂- and N₂O-reducing bacteria to maize inoculation by the PGPR *Azospirillum lipoferum* CRT1 depend on carbon availability and determine soil gross and net N₂O production

Allessandro Florio, Caroline Bréfort, Jonathan Gervaix, Annette Bérard,
Xavier Le Roux

► To cite this version:

Allessandro Florio, Caroline Bréfort, Jonathan Gervaix, Annette Bérard, Xavier Le Roux. The responses of NO₂- and N₂O-reducing bacteria to maize inoculation by the PGPR *Azospirillum lipoferum* CRT1 depend on carbon availability and determine soil gross and net N₂O production. *Soil Biology and Biochemistry*, 2019, 136, pp.107524. 10.1016/j.soilbio.2019.107524 . hal-02473989

HAL Id: hal-02473989

<https://univ-lyon1.hal.science/hal-02473989>

Submitted on 25 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **The responses of NO₂⁻ and N₂O-reducing bacteria to maize inoculation by**
2 **the PGPR *Azospirillum lipoferum* CRT1 depend on carbon availability and**
3 **determine soil gross and net N₂O production**

4

5 Alessandro Florio*¹, Caroline Bréfort¹, Jonathan Gervais¹, Annette Bérard² & Xavier Le Roux¹

6

7 ¹Laboratoire d'Ecologie Microbienne LEM, INRA UMR 1418, CNRS UMR 5557, Université Lyon 1,

8 Université de Lyon, F-69622 Villeurbanne Cedex, France

9 ²INRA, UMR1114 EMMAH, Site Agroparc, 84914, Avignon, France.

10

11

12 ***Corresponding author:**

13 Tel: +33 4 72 43 13 78 ; Email: alessandro.florio@univ-lyon1.fr

14 **Abstract**

15 Seed inoculation by plant growth promoting rhizobacteria (PGPRs) is an agronomic practice that
16 stimulates root carbon (C) exudation and nitrogen (N) uptake. Inoculation thus increases and decreases C
17 and N availabilities to denitrifiers in the rhizosphere, respectively. Hence, denitrification rates in the
18 rhizosphere can be positively or negatively influenced by root activity depending on the balance
19 between these two processes. We assumed that inoculation effect on denitrifiers could strongly differ
20 according to soil conditions. Would denitrifiers be mostly limited by C, inoculation would increase
21 denitrifier abundance and activity through increased labile C availability. Would denitrifiers be limited by
22 N rather than C, inoculation would decrease denitrifier abundance and activity through increased
23 competition for N. Here we manipulated denitrification limitation by C and N (i) in a field trial through
24 the use of different fertilization levels, and (ii) in a growth chamber experiment by mimicking root
25 exudate inputs. We analyzed how the effects of maize inoculation by the PGPR *Azospirillum lipoferum*
26 *CRT1* on potential gross and net N₂O production rates and NO₂⁻ and N₂O-reducer abundances were
27 related to C and N limitation levels. An increase in potential gross (up to +113%) and to a lesser extent
28 net (+37%) N₂O production was observed for soils where denitrification was highly limited by C. This was
29 explained by strong and moderate increases in the abundances of NO₂⁻ and N₂O-reducers, respectively.
30 In contrast, when denitrification was weakly limited by C, gross and net N₂O productions were negatively
31 affected by inoculation (-15 and -40%, respectively). Our results show that the inoculation practice
32 should be evaluated in term of possible increased crop yield but also possible modified N₂O emission,
33 paying much attention to cropland soils where denitrifiers are highly limited by C.

34

35 **Key-words:** *nirK*, *nirS*, *nosZI*, *nosZII*, plant-microbes competition for nitrate, root exudates

36 1. Introduction

37 Denitrification is a microbial respiratory process during which soluble nitrogen (N) oxides (nitrate,
38 NO_3^- , and nitrite, NO_2^-) are sequentially reduced by specific reductases into gaseous forms (NO , N_2O and
39 N_2) (Tiedje et al. 1982). In particular, NO_2^- reductases, encoded by the *nirK* or *nirS* gene, catalyze the
40 reduction of NO_2^- to NO , which is the first step of denitrification, leading to the production of a gaseous
41 N oxide (Shapleigh, 2013). This is the key process for both NO and N_2O production as most NO -producing
42 denitrifying cells convert efficiently NO which is a toxic compound (this explains why most *nir*-harbouring
43 bacteria also harbor the *nor* gene catalyzing NO reduction; Graf et al., 2014). The reduction of N_2O to N_2
44 is catalyzed by N_2O reductases encoded by the *nosZI* gene or the recently discovered *nosZII* gene
45 (Sanford et al. 2012; Jones et al. 2013; Domeignoz-Horta et al., 2016). Thus, the net soil emissions of N_2O
46 (a potent greenhouse gas; Baggs, 2011) depend largely on the balance of its production and
47 consumption, and on the responses of NO_2^- - and N_2O -reducers to changes in soil environmental
48 conditions.

49 In particular, two main factors affect the ecology of NO_2^- - and N_2O -reducers in the plant rhizosphere,
50 namely the availabilities of N and carbon (C). On the one hand, root growth and activity enhance soil
51 NO_3^- uptake by plants, thus strengthening plant-denitrifiers competition for NO_3^- . This can decrease soil
52 NO_3^- availability for denitrifiers and ultimately limit their growth (Kuzyakov and Xu, 2013). On the other
53 hand, root exudation provides easily decomposable C sources to denitrifiers, which can increase their
54 activity and abundance (Berks et al., 1995). Hence, denitrification rates in the rhizosphere can be
55 positively or negatively influenced by root activity depending on the balance between these two
56 processes. However, to what extent NO_2^- - and N_2O -reducers might display different sensitivities to C
57 and/or N availability and how this might affect the responses of soil N_2O production and consumption to
58 changed C and N availabilities remains unclear. For instance, we synthesized the results of previous
59 studies analyzing the effects of soil amendments with labile C sources, mineral N, labile C plus mineral N,

60 or artificial root exudates (ARE) and we found that C or N addition increases potential gross and net N₂O
61 production (PGNP and PNNP, respectively) in only 50%-70% of the soils studied (Table 1). One
62 explanation is that the type of soil and in particular the soil C and N statuses influence the C or N addition
63 effects. This synthesis also shows that the effect of mineral N addition on the abundances of NO₂⁻ and
64 N₂O-reducers has been analyzed in many studies, the effect being generally low, without any clear
65 difference between *nirK*-, *nirS*-, *nosZI*- and *nosZII*-bacteria (Table 1). In contrast, fewer studies reported
66 the effect of labile C amendment or of the addition of labile C plus mineral N, information being missing
67 for some groups (Table 1). This shows that it is still largely unknown whether these denitrifier groups
68 respond differently to changes in mineral N or labile C availabilities.

69 Inoculation of cereal seeds by plant growth promoting rhizobacteria (PGPRs) increases root C
70 exudation (Heulin et al., 1987; Shaw et al., 2006) and enhances N uptake by inoculated plants (Sarig et
71 al., 1988; Fallik et al., 1994; Mantelin and Touraine, 2004). Recently, Florio et al. (2017) analyzed how the
72 promoting activity of PGPRs may influence the activity and abundance of denitrifiers in rhizosphere soil
73 by modifying C and N availabilities. The authors reported contrasted effects of inoculation of denitrifiers,
74 between different soil types. They suggested that inoculation could increase and decrease *nirS*
75 abundance and consequently potential gross N₂O production when denitrification was highly and lowly
76 limited by soil C, respectively. However, soil type was a confounding factor with C availability in this
77 study which compared inoculation effects between different sites and soil types. Distinguishing the
78 effects of C and N availability from the effects of other soil characteristics actually requires to manipulate
79 C and N availabilities to denitrifiers using a same soil. Further, the authors studied inoculation effect on
80 potential gross but not net N₂O production.

81 Here we manipulated C and N availabilities for a same soil background by using different mineral
82 fertilization levels and mimicking different maize root exudation rates at the field and microcosm scales,
83 and we assessed the effects of maize inoculation by the PGPR *Azospirillum lipoferum CRT1* on potential

84 gross and net N₂O production rates, and on the abundances of NO₂⁻ and N₂O-reducers, according to C
85 and N availabilities. We hypothesized that when denitrification is highly limited by C, inoculation would
86 increase the abundance and activity of NO₂⁻ and N₂O-reducers as higher maize root exudation would
87 have a key role (Fig. 1, red arrows). In contrast, when denitrification is less limited by C, the stimulating
88 effect of root exudation would be less important than the effect of the increased competition between
89 plants and denitrifiers for N, and inoculation would decrease denitrifier abundance and activity (Fig. 1,
90 blue arrows). In addition, the sensitivity of the different denitrifier groups, in particular of NO₂⁻-reducers
91 as compared to N₂O-reducers, to C and N availabilities, would determine the overall effect on gross and
92 net N₂O production, which could not be easily predicted due to the lack of sufficient information on their
93 ecology regarding C and N (Table 1). We discuss our results in term of possible implications of the
94 inoculation practice for N₂O emission according to soil type.

95

96 **2. Materials and Methods**

97 *2.1. Field experiment*

98 The experimental site is located in Sérézin-de-la-Tour, southeast of France (45°37' N, 5°16' E). The
99 soil is a Fulvic Cambisol (World Reference Base for Soil Resources, 2006), and its main physical and
100 chemical characteristics are as follows: 34.7% clay, 26.9% sand, 38.3% silt; pH (H₂O 1:2.5) 7.1; SOC 31.6 g
101 C kg⁻¹; TN 3.4 g N kg⁻¹; and Olsen P 0.153 g kg⁻¹. The experiment was set up as a randomized block design
102 with 5 blocks, and treatments randomly assigned to one plot (12 m × 9.6 m) in each block. The
103 experimental fields had been cultivated with wheat for three years previous to the experiment. Maize
104 (*Zea mays*, cv. *Seiddi*) seeds were inoculated with *A. lipoferum CRT1* isolated from the rhizosphere of
105 field-grown maize in France (Fages and Moulard, 1988). The targeted inoculum load was 10⁶ CFU added
106 per seed for inoculated plants, I, coated in a commercial peat-based Azo-Green™ formulation
107 (Agrauxine, Beaucozé, France). Coated but non-inoculated seeds, NI, were used as controls. Sowing

108 occurred on 30th April 2015 (95,000 seeds ha⁻¹). Five pairs of NI-I plots were not fertilized (nf plots) while
109 two fertilization treatments were applied using a mineral fertilizer (NH₄NO₃) at a rate of 80 kg N ha⁻¹
110 close to optimal N availability (f plots) or 40 kg N ha⁻¹ for the reduced fertilization treatment (f/2 plots).
111 Ten plots at an additional field site located in the proximity of the first site (45°57' N, 5°34' E) and
112 managed under organic farming with feather meal used as organic fertilizer at a rate of 120 kg N ha⁻¹ (F-
113 org) were also included in the experimentation. The soil is a Calcisol (siltic) (World Reference Base for
114 Soil Resources, 2006), and its main physical and chemical characteristics are as follows: 10.2% clay,
115 27.4% sand, 62.5% silt; pH (H₂O 1:2.5) 8.1; SOC 20.0 g C kg⁻¹; TN 2.1 g N kg⁻¹; and Olsen P 0.134 g kg⁻¹.
116 This led to a total of 40 plots, *i.e.* 4 treatments x 5 pairs of NI-I plots.

117 Rhizosphere soil (0-20 cm) was sampled on 5th June at the 6-leaves stage. Six individual plants were
118 randomly selected from each plot and removed using a spade to excavate the root system. Rhizosphere
119 soil was collected by gently shaking the roots. Fresh soil retrieved from the 6 plants was pooled, sieved
120 using 2-mm mesh size and stored at +4 °C a few days before activity measurements.

121

122 2.2. Growth chamber experiment

123 The soil (500 Kg) for the growth chamber experiment was collected from the surface layer (0-30 cm) of
124 non-inoculated and non-fertilized plots following the end of field experiment in Sérézin-de-la-Tour. The
125 soil was air-dried, homogenized by sieving (2-mm mesh size), pooled and stored a few days at room
126 temperature before microcosm preparation. The experiment was conducted in pots (11.3 x 11.3 x 21.5
127 cm³), each filled with 1.8 Kg soil. All pots were flooded with distilled water in order to leach excess
128 mineral N, and kept to 70% of soil water holding capacity. Pots were transferred to an environmental
129 growth chamber (photoperiod 16 h, temperature 19 °C night and 26 °C day, relative humidity 65%, PPFD:
130 350 μmol m⁻² s⁻¹). One inoculated, I, or non-inoculated, NI, maize seed (same maize cv. and same
131 inoculation load as for the field experiment) was sown in each pot at a depth of 1 cm. The experiment

132 was set up as a randomized block design with 6 blocks, and treatments randomly assigned to one pot in
133 each block. Pots were kept in the chamber until maize plants reached the 6-leaves stage (25 days).
134 During the first ten days, all maize plants were watered every other day with 10 ml H₂O_{dd}; then pots
135 were amended with artificial root exudates C (ARE-C treatment) or distilled water (H₂O_{dd} control
136 treatment). ARE-C solution was prepared with sterilized H₂O_{dd} and 9 C sources identified as main
137 constituents of maize root exudates (Krafczyk et al., 1984; Baudoin et al., 2003). The stock solution at C
138 concentration of 25 mg C ml⁻¹ contained 3 carbohydrates (66.75 mM glucose, 66.75 mM fructose and
139 79.75 mM arabinose), 3 carboxylic acids (49.5 mM succinic acid, 33.25 mM citric acid and 49.5 mM
140 fumaric acid) and 3 amino acids (33.25 mM alanine, 25 mM aspartic acid and 19.5 mM glutamic acid).
141 Working solutions consisted of 12.5 ml of stock solution mixed with H₂O_{dd} in order to ensure an
142 enrichment of 250, 100 or 20 µg C g⁻¹ soil per pot. Furthermore, additional soil pots with 100 µg C g⁻¹
143 were set up and fertilized with KNO₃ at 40 Kg N ha⁻¹. Half of the N fertilizer was added at sowing (day 0)
144 and half at 3-leaves stage (day 10). At harvest (6-leaves stage), fresh rhizosphere soil was collected by
145 gently shaking roots and stored at +4 °C or -20 °C before potential activities and molecular
146 measurements. A ~10 g subsample was weighed and dried at 105°C during 24 h to determine gravimetric
147 soil moisture. Soil nitrate concentration was measured using 5 g equivalent dry weight soil from I and NI
148 plots after extraction with 20 ml of 2 mol l⁻¹ KCl. The extraction solution was shaken at 10 °C for 1 h at
149 140 rpm, filtered at 0.2 µm and frozen at -20 °C until measurements of NO₃⁻ concentrations were made
150 using an ion chromatograph (DX120 Dionex, Salt Lake City, USA) equipped with a 4 × 250 mm column
151 (IonPac AS9 HC). The sum of substrate-induced respiration quantified by Community Level Physiological
152 Profiles (CLPP) using the MicroResp™ system (Campbell et al. 2003) was used as an index of the
153 heterotrophic microbial biomass. The MicroResp™ system consists of a 96-deep-well microplate (1.2-ml
154 volume) filled with soil from I and NI plots and with the addition of a range of aqueous C substrates (SIR),
155 as described by Bérard et al. (2012).

156

157 *2.3. Potential gross (PGNP) and net (PNNP) N₂O production measurements*

158 PGNP was measured for each defrosted fresh soil sample according to Patra et al. (2005), using 3.5 g
159 dw equivalent soil per 150 mL flask. The atmosphere of each flask was replaced by a 90:10 He-C₂H₂
160 mixture to provide anaerobic conditions and inhibit N₂O-reductase activity. PGNP was determined as the
161 linear rate of the gross production of N₂O during short-term (8 h) incubation using a gas chromatograph
162 (μ GC R3000, Santa Clara, CA, USA), the headspace being sampled every 1.5 h. PNNP was determined for
163 each defrosted fresh soil sample except for F-org soils, using 3.5 g dw equivalent soil. PNNP was
164 determined as the linear rate of the N₂O production during short-term (8 h) incubation under anaerobic
165 conditions but without C₂H₂ addition. Glucose (0.5 mg C g⁻¹ dry soil), glutamic acid (0.5 mg C g⁻¹ dry soil)
166 and KNO₃ (50 μ g NO₃⁻-N g⁻¹ dry soil) were added to the soil samples and the soil moisture was brought to
167 100% water holding capacity. To test whether soil storage at -20°C affected denitrification, PGNP had
168 also been measured on each fresh soil sample, using the same protocol as detailed above. PGNP from
169 frozen soils were strongly correlated to those from fresh soils without any significant effect of storage
170 ($y=0.99x$, $R^2=0.65$, $p<0.0001$; regression not significantly different from the 1/1 line).

171 In addition, semi-potential gross N₂O production when either C (GNP_C) or N (GNP_N) was not added
172 were measured in order to evaluate the limitation of denitrification by C and N, respectively (Florio et al.,
173 2017). In these cases, denitrification depended on the availability of soil C (including ARE-C) and N,
174 respectively. Linearity of the N₂O production rate was always observed over 8 h, whatever the substrate
175 added. Activity measurements were performed at the AME platform (Microbial Ecology UMR5557,
176 Lyon).

177

178 *2.4. Quantification of the abundances of PGPR Azospirillum CRT1, and NO₂⁻ and N₂O- reducers*

179 DNA was extracted from 0.5 g of soil using the FASTDNA SPIN Kit for Soil (BIO 101 Systems;
180 Qbiogene, Carlsbad, CA, USA). DNA concentrations were determined using a Qubit® 2.0 fluorometer with
181 Quant-iT™ dsDNA broad range (BR) Assay Kit (Invitrogen, France).

182 The abundance of *Azospirillum CRT1* was measured for 6-leaves stage soil samples by quantitative
183 PCR as described by Couillerot et al. (2010) but qPCR counts were always below detection limit (*i.e.* less
184 than 10^2 g^{-1} soil).

185 The abundances of NO_2^- -reducers were measured by quantitative PCR targeting the *nirK* and *nirS*
186 genes (encoding the copper and *cd*₁ nitrite reductases, respectively). Amplification was performed using
187 primers nirK876/nirK1040 (Henry et al., 2006) or nirSCd3aF/nirSR3cd (Throbäck et al., 2004). The 20 μL
188 final volume PCR mix contained (final concentrations) QuantiTect SybrGreen PCR Master Mix 1 x, 1 μM of
189 each *nirK* primer or 0.5 μM of each *nirS* primer, 0.4 mg of T4 protein, and 5 ng or 12.5 ng of soil DNA
190 extract for *nirK* or *nirS*, respectively. Quantitative PCR was performed as follows: 15 min at 95 °C, 40
191 (*nirK*) or 45 (*nirS*) amplification cycles (15 s at 95 °C, 30 s at 59 °C for *nirS* or 63 °C for *nirK*, 30 s at 72 °C,
192 and 10 s at 40 °C).

193 The abundances of N_2O -reducers were measured by targeting the *nosZI* and *nosZII* genes (encoding
194 the N_2O reductases corresponding to two distinct clades). Amplification was performed using primers
195 nosZ2F/nosZ2R (Henry et al., 2006) or nosZ-II-F/nosZ-II-R (Jones et al., 2013). The 25 μL final volume PCR
196 mix contained (final concentrations) QuantiTect SybrGreen PCR Master Mix 1 x (*nosZI*) or 1.2 x (*nosZII*), 1
197 μM of each primer, 0.8 mg of T4 protein (*nosZI*) or 2% BSA (*nosZII*), and 12.5 ng or 20 ng of soil DNA
198 extract for *nosZI* or *nosZII*, respectively. Quantitative PCR was performed as follows: 15 min at 95 °C, 6
199 touchdown amplification cycles for *nosZI* (15 s at 95 °C, 30 s at 65 °C, 30 s at 72 °C, and 15 s at 80 °C) and
200 45 amplification cycles (15 s at 95 °C, 30 s at 60 °C for *nosZI*; or 53 °C for *nosZII*, 30 s at 72 °C, and 10 s at
201 40 °C).

202 Standards were generated from PCR products obtained from soil DNA extracts as described by Florio
203 et al., 2017). Possible inhibitory effects of co-extracted humic compounds in soil extracts were checked
204 by dilution series, but no inhibition was observed. A melting curve analysis was performed to assess PCR
205 product specificity after amplification. The average real-time PCR efficiency for each of these genes was
206 97%, 100%, 86% and 84% for *nirK*, *nirS*, *nosZI* and *nosZII*, respectively. Gene copy number per gram of dry
207 soil was calculated from the copy number of each gene per ng of DNA multiplied by the amount of DNA
208 extracted from one gram of dry soil.

209

210 2.5. Statistical analyses

211 Significant effects of inoculation on microbial activities and abundances were identified using two-way
212 ANOVA with inoculation and C limitation as factors (JMP Pro 12, SAS Institute, Cary, North Carolina,
213 USA). Where necessary, data were log-transformed to ensure conformity with the assumptions of
214 normality and homogeneities of variances. For each pair of NI-I plots or microcosms, *i.e.* corresponding
215 to the same treatment within a given block, the effect of inoculation on a given variable V (*i.e.*, PGNP,
216 PNNP, and abundances of NO₂⁻ and N₂O -reducers) was expressed as:

$$217 \quad \% \text{ Inoculation effect} = \left(\frac{V(I)}{V(NI)} - 1 \right) * 100.$$

218 The limitations of denitrification by C or N were computed for NI treatments as follows:

$$219 \quad \text{Limitation of denitrification by C} = \left(1 - \frac{GNP(C-)}{PGNP} \right) * 100.$$

$$220 \quad \text{Limitation of denitrification by N} = \left(1 - \frac{GNP(N-)}{PGNP} \right) * 100.$$

221 Correlations were carried out to investigate the relationships (i) between the limitation of denitrification
222 by C and the amount of ARE-C amended to pots or SIR; (ii) between the limitation of denitrification by N
223 and the soil nitrate concentration; and (ii) between the inoculation effects on NO₂⁻-reducers, N₂O -
224 reducers, PGNP, PNNP, and C or N limitation.

225

226 **3. Results**

227 *3.1. Limitation of denitrification by C or N in the field and according to the amount of artificial root* 228 *exudates added to microcosms*

229 For field plots and non-amended microcosms, denitrification was strongly limited by organic C, with
230 potential gross N₂O production measured without C addition (GNP_C, *i.e.* when denitrification activity
231 depends only on soil endogenous C supply, see 2.5 section) being reduced by 65-76% as compared to
232 PGNP (Supplementary Tab. S1). For microcosms, the limitation of denitrification by C significantly and
233 gradually decreased when the amount of ARE-C amended increased (Fig. 2). For the highest ARE-C level,
234 limitation of denitrification by organic C was reduced to 31% (Fig. 2). For amended microcosms,
235 fertilization did not affect the level of denitrification limitation by soil C (Fig. 2). Furthermore, organic
236 matter amendment according to the local organic farming practices led to significant reduction in
237 limitation by C down to 51% (Fig. 2).

238 The level of denitrification limitation by NO₃⁻ was always lower than the level of limitation by C for
239 field plots (29-35% and 51-76%, respectively; Supplementary Tab. S1), including non-fertilized plots, and
240 values for microcosms were not correlated with ARE-C amendment or fertilization levels (data not
241 shown). Furthermore, values of limitation of denitrification by N or C were negatively and significantly
242 correlated with soil nitrate content (Fig. S1a) and SIR (as a measure of microbial heterotrophic biomass
243 C; Fig. S1b), respectively.

244

245 *3.2. Effects of inoculation and C limitation on potential gross and net N₂O production rates, and on* 246 *denitrifier abundances*

247 Values of PGNP for field and non-amended microcosm samples ranged from 1.5 to 3.9 μg N g⁻¹ h⁻¹,
248 whereas PGNP values for amended microcosms significantly increased (p<0.0001) with increased ARE-C

249 levels from 3.8 to 7.1 $\mu\text{g N g}^{-1} \text{h}^{-1}$ (Supplementary Tab. S1). Values of GNP_{C} ranged from 0.74 to 4.64 $\mu\text{g N}$
250 $\text{g}^{-1} \text{h}^{-1}$ for F-org NI and ARE-C₂₅₀ NI plots, respectively, whereas values of GNP_{N} ranged from 0.80 to 5.07
251 $\mu\text{g N g}^{-1} \text{h}^{-1}$ for ARE-C₁₀₀ NI and ARE-C₂₅₀ I plots, respectively (Supplementary Tab. S1). Values of PNNP for
252 field and non-amended microcosms samples ranged from 1.5 to 3.4 $\mu\text{g N g}^{-1} \text{h}^{-1}$, whereas PNNP values
253 for amended microcosms significantly increased with increased ARE-C levels from 2.1 to 4.9 $\mu\text{g N g}^{-1} \text{h}^{-1}$
254 (Supplementary Tab. S1).

255 The abundance of *nirK*-harbouring NO_2^- -reducers was in the same order of magnitude as the
256 abundance of *nirS*-bacteria for field plots, *i.e.* typically from 1.3×10^6 to 1.6×10^7 copies g^{-1} dry soil, but it
257 was slightly higher for *nirK*- than *nirS*-bacteria for microcosms (*i.e.* from 1.8×10^8 to 2.4×10^8 and from
258 3.9×10^7 to 5.7×10^7 copies g^{-1} dry soil, respectively; Supplementary Tab. S2). The abundances of *nirK*-
259 and *nirS*-bacteria were significantly and positively correlated ($p < 0.0001$). The abundance of NO_2^- -
260 reducers as the sum of *nirK* and *nirS* abundances (*i.e.* total *nir* abundance) increased with increasing ARE-
261 C levels ($p = 0.0007$).

262 The abundance of *nosZI*-harbouring N_2O -reducers was in the same order of magnitude as the
263 abundance of *nosZII* for field plots, *i.e.* typically from 4.9×10^5 to 1.5×10^7 copies g^{-1} dry soil, but it was
264 higher for *nosZI* than *nosZII* for microcosms (*i.e.* from 2.4×10^6 to 4.4×10^6 and from 3.7×10^5 to 4.6×10^5
265 copies g^{-1} dry soil, respectively; supplementary Table S2). The total abundance of N_2O -reducers (sum of
266 *nosZI* and *nosZII* abundances, *i.e.* *nosZ* abundance) increased with increasing ARE-C levels ($p < 0.0001$).

267 Two-way ANOVA results showed a significant main effect of C limitation on *nosZI* and total *nosZ*
268 abundances, and for PGNP (Table 2). Furthermore, a significant interaction effect between inoculation
269 and C limitation was observed for the abundances of the NO_2^- -reducers ($p = 0.026$, $p = 0.047$ and $p = 0.024$
270 for *nirK*, *nirS* and total *nir* abundances, respectively) and N_2O -reducers ($p = 0.019$ and $p = 0.030$ for *nosZII*
271 and total *nosZ* abundances, respectively), and for PGNP and PNNP ($p = 0.035$ and $p = 0.023$, respectively)

272 (Table 2). This indicates that denitrifier abundances and activities were significantly affected by
273 inoculation but with the effects varying depending on C limitation levels.

274

275 *3.3. Relationship between the inoculation effects on PGNP and PNNP, and C limitation*

276 A positive and exponential relationship was observed between the inoculation effect on PGNP and
277 the level of denitrification limitation by C ($R^2=0.92$, $p<0.0001$; Fig. 3a), *i.e.* the higher the C limitation, the
278 higher the inoculation-induced increase in PGNP. In particular, for field plots with the highest C limitation
279 (70-76%), the increase of PGNP in response to inoculation was highest, reaching up to +113%. Values of
280 C limitation for non C-amended microcosms (65%) were slightly lower than those observed for field
281 plots, and were associated to a lack of inoculation effect on PGNP (Fig. 3a). Conversely, for microcosm
282 soils amended with ARE-C, lower denitrification limitation by C were observed (ranging from 31 to 63%)
283 and inoculation effect on PGNP was then neutral to negative (from +1% to -17%). No relationship was
284 observed between the inoculation effect on PGNP and the level of denitrification limitation by N when
285 considering the F and G treatments together (Supplementary Fig. S2a) or separately (data not shown)".

286 A positive relationship was observed between the inoculation effect on PNNP and the level of
287 denitrification limitation by C ($R^2=0.69$, $p=0.011$; Fig. 3b). The inoculation effect on PNNP in field plots
288 exhibiting the highest increase in PGNP was positive but reached only +37% (Fig. 3b). Conversely, for
289 soils amended with ARE-C, inoculation decreased PNNP down to -46%. No relationship was observed
290 between the inoculation effect on PNNP and the level of denitrification limitation by N (Supplementary
291 Fig. S2b).

292

293 *3.4. Relationships between the inoculation effect on denitrifier abundances and C limitation*

294 The inoculation effect on the abundance of NO_2^- -reducers was significantly and positively related to
295 the limitation of denitrification by C. The best relationship was observed for the sum of *nirS*- and *nirK*-

296 harbouring bacteria ($R^2=0.86$, $p<0.0001$; Fig. 3c). In particular, inoculation increased *nir* abundance (up to
297 +91%) in field plots where limitation by C was the highest. In contrast, inoculation had a weak effect on
298 *nir* abundance (-7% to +5%) in microcosms and organic plots where C limitation was lower. No
299 relationship was observed between the inoculation effect on *nir* abundance and the level of
300 denitrification limitation by N (Supplementary Fig. S2c).

301 The inoculation effect on the total abundance of N_2O -reducers was significantly and positively
302 related to the limitation of denitrification by C ($R^2=0.66$, $p=0.012$, Fig. 3d). In particular, inoculation had
303 no or little effect on *nosZ* abundance (from -10 to +20%) in field plots that corresponded to the highest
304 values of limitation of denitrification by C, and it had a slightly negative effect in microcosms where C
305 limitation was lower (ranging from -19% for soil receiving the highest ARE-C dose to -6% for the medium
306 ARE-C dose and fertilized soil). No relationship was observed between the inoculation effect on *nosZ*
307 abundance and N limitation (Supplementary Fig. S2d).

308

309 **4. Discussion**

310 Over the last 20 years, strategies for sustainable agricultural development, including natural systems
311 agriculture and nature-based solutions (Eggermont et al. 2015), have been developed worldwide to
312 promote agroecosystem multi-functionality (Altieri, 1999). Major challenges are faced by farmers when
313 developing more sustainable agricultural systems less dependent on chemical inputs, and better use of
314 biotic interactions is part of their toolbox to promote the performance of agroecosystems under lower
315 chemical inputs (Barot et al., 2017). In this context, the practice of cereal inoculation with PGPRs is a
316 promising alternative to classical, intensive cropping systems for maintaining high yield while decreasing
317 fertilizer inputs (El Zemrany et al., 2006). Technical-economic acceptance by farmers and avoidance of
318 any negative side effects, including greenhouse gas emission from soils, are two major challenges
319 associated to cereal seed inoculation by *Azospirillum* (Bounaffaa et al., 2018). It has been reported that

320 the *Azospirillum* strains used for inoculation are unable to establish in soil, and that their abundance
321 generally drops below detection limit a few weeks after inoculation (Bashan, 1999) as also observed in
322 our study. Despite this, inoculation has been reported to indirectly affect soil microbial communities
323 (Baudoin et al. 2009) likely through lasting effects on plant and root growth and development. It is thus
324 important to assess whether inoculation can have some unintentional side effects on soil functioning, in
325 particular regarding greenhouse gas production.

326

327 *4.1. Fertilization and mimicking root exudation are effective in generating a range of levels of*
328 *denitrification limitation by C and N in the field and in microcosms*

329 In the rhizosphere, C and N availabilities are among the main factors modulating the interactions
330 between plant roots and denitrifying microorganisms (Philippot et al., 2007). Our hypothesis was that
331 maize inoculation by a PGPR would affect potential net and gross N₂O production rates and the
332 abundances of microbial groups playing a key role for N₂O production (NO₂⁻-reducers) and consumption
333 (N₂O-reducers) by altering two main biological processes which occur simultaneously in the rhizosphere
334 but act in an opposite way (see Fig. 1). The higher NO₃⁻ uptake by plants observed in response to
335 inoculation (Mantelin and Touraine, 2004) should strengthen roots-microbes competition for NO₃⁻, which
336 should limit NO₃⁻ availability for microorganisms and thus decrease the abundance and activity of the
337 denitrifier groups mostly sensitive to NO₃⁻ availability. In contrast, the higher release of C exudates that
338 maize inoculation induces (Heulin et al., 1987; Shaw et al., 2006) should favor the activity and growth of
339 microbial heterotrophs, including denitrifiers. We also assumed that the relative importance of the two
340 processes would vary according to the importance of denitrification limitation by C and N. Testing this
341 hypothesis implied to manipulate the levels of denitrification limitation by C and N for the model soil
342 studied, which was achieved by adding mineral N fertilizer (in both field and growth chamber
343 experiments) and mimicking maize root exudate inputs (through addition of ARE-C in the microcosms).

344 This allowed us to generate a wide range of values for denitrification limitation by C and N for a given
345 soil. We observed that denitrification limitation by C was always higher than limitation by N for crop field
346 conditions, even for non-fertilized plots, and that limitations by N or C were negatively and significantly
347 correlated with soil nitrate content (Fig. S1a) and SIR (Fig. S1b), respectively. This could be due to a
348 rather high N and low C status of agricultural soils, in relation to previous years' fertilization practices
349 and to the important biomass export from annual cropping systems where soil C tends to decrease with
350 time (Recous et al., 1995). The addition of ARE-C to microcosms significantly decreased C limitation as
351 expected. We used realistic rates of daily ARE-C inputs, similar to rates used in other studies (Trofymow
352 et al., 1987; Iijima et al., 2000, Baudoin et al., 2003; Henry et al., 2008). Further, we applied ARE-C
353 repeatedly through low and recurrent additions rather than a large and single pulse, to better mimic the
354 exudation process. This probably allowed denitrifiers to grow and adapt to higher C availability
355 throughout the experiment, because a few days is sufficient to observe an increase in the size of
356 denitrifying communities following C addition (Henry et al., 2008). This can explain why a certain level of
357 denitrification limitation by C was still observed even at the highest ARE-C level. The treatment G-nf-ARE
358 250 not only decreased C limitation but also decreased N limitation (see Fig. S1), and it might thus be
359 possible that mineralization of the 3 exudate compounds that include N (i.e. the 3 amino acids, among
360 the 9 compounds used) fuelled N supply to denitrifiers. However, only 3 among the 9 exudate
361 compounds added to soil included N, the overall C:N ratio of the pool of artificial exudates used being
362 quite high (23.2); furthermore, no relationship was observed between N- and C-limitation levels,
363 considering either the whole data set or the data set from the microcosm experiment (not shown). Thus,
364 we did decouple N- and C-limitations in our study (even if the use of N-free exudates only could have
365 improved this decoupling). Overall, our approach allowed us to explore inoculation effects on denitrifiers
366 over a broad range of denitrification limitation by C (limitation level from 31 to 76%) and by N (from 29

367 to 69%) using a same, manipulated soil rather than comparing different soils to avoid the problem of
368 having confounding factors.

369
370 *4.2. The contrasted effects of inoculation on potential gross and net N₂O production are explained by*
371 *denitrification limitation by soil C*

372 Carbon availability is often recognized as the main determinant of denitrification in soil (Myrold and
373 Tiedje, 1985; Weier et al., 1993; Schaeffer et al., 2003), particularly in cropland soils (Chantigny et al.,
374 2010; Attard et al., 2011). Consistently, we observed that the effect of maize inoculation on potential
375 gross N₂O production was significantly and positively related to the denitrification limitation by C but not
376 to limitation by N. This shows that organic C rather than NO₃⁻ availability to denitrifiers controls
377 inoculation effects on potential gross N₂O production. Although measuring root exudation in soil is
378 challenging (Weixin et al., 1993) and was beyond the scope of our study, we can assume that the
379 increased root C exudation from inoculated plants drove the strong inoculation-induced increase (up to
380 +113%) in potential gross N₂O production for soils where denitrifiers were highly C-limited. Such an
381 amplitude of the stimulation of denitrification is consistent with a root exudation effect, because gross
382 N₂O production has been reported to increase by 16-250% in the rhizosphere as compared to bulk soil
383 (Stefanson, 1972; Vinther et al., 1982; Højberg et al., 1996; Mahmood et al., 1997) and by 50-660% in
384 response to ARE-C inputs to soil (Mounier et al., 2004; Henry et al., 2008; Langarica-Fuentes et al., 2018).
385 When denitrification limitation by C was decreased by recurrent root exudate inputs, the negative effect
386 of the competition between roots and denitrifiers for NO₃⁻ seemed to prevail, and the resulting outcome
387 of inoculation was a slightly negative effect on potential gross N₂O production (-17%).

388 The effect of inoculation on potential net N₂O production was also mainly related to denitrifier
389 limitation by C as it was positive (up to +37%) when level of denitrification limitation by C was high,
390 whereas it was negative (down to -46%) at low C limitation levels. However, when denitrification

391 limitation by C was high, the inoculation effect was lower for PNNP than PGNP (+37% and +113%,
392 respectively). The review of the literature concerning the effects of labile C and/or mineral N addition
393 provide no clue to explain these different amplitudes of the inoculation effects, as PNNP and PGNP seem
394 to respond similarly to labile C and mineral N additions (Table 1).

395

396 *4.3. The effects of inoculation on potential gross and net N₂O production are related to inoculation*
397 *effects on the abundances of NO₂⁻ and N₂O-reducers*

398 The activity and abundance of denitrifiers are not necessarily tightly coupled, since the synthesis of
399 denitrifying enzymes is inducible (Zumft, 1997). Moreover, potential net N₂O production depends on the
400 balance between the activity and abundance of NO₂⁻ and of N₂O- reducers (Chapuis-Lardy et al., 2007;
401 Assemien et al., 2019). Here we observed that inoculation-induced changes in potential gross N₂O
402 production were strongly and positively related to inoculation-induced changes in the abundance of *nir*-
403 harbouring bacteria (relationship between inoculation effects on PGNP and *nir* abundance: $y=0.48x$;
404 $R^2=0.88$; $p<0.0001$). Several authors have already reported such a coupling between changes in potential
405 gross denitrification and *nirS* and/or *nirK* abundances in agricultural soils (Čuhel et al., 2010; Enwall et al.,
406 2010; Attard et al., 2011; Jusselme et al., 2016; Assemien et al., 2019), although this is not necessarily the
407 case (Le Roux et al., 2013). This suggests that inoculation conditioned potential gross N₂O production in
408 the rhizosphere by mediating the build-up of NO₂⁻-reducers, probably due to changed C availability.

409 The inoculation effect on NO₂⁻-reducer abundance was concomitant to an inoculation effect on N₂O
410 reducer abundance, but the magnitude of the effect varied notably between these groups. Specifically,
411 the inoculation effect on total *nosZ* abundance was only slightly positive (up to +20%) when the effect on
412 *nir* abundance was highly positive (up to +91%). As the strong increase in NO₂⁻-reducers resulted in a
413 strong increase in PGNP, the moderate increase in N₂O reducers abundance partially dampened the
414 potential N₂O production induced by inoculation. This likely explains why inoculation increased PNNP but

415 not as highly as for PGNP under these conditions. These results suggest that N₂O emissions from maize
416 croplands could be increased by the maize inoculation practice in the case soils are characterized by high
417 levels of denitrification limitation by C. This calls for specific quantification of N₂O emissions from soils of
418 inoculated and non-inoculated plots, in particular targeting croplands where limitation of denitrifiers by
419 C is high.

420 The different responses of the abundances of NO₂⁻ and N₂O reducers to inoculation might be due to
421 their different sensitivity to C and N availability. Actually, our synthesis of results from previous studies
422 regarding denitrifier abundances (Table 1) does not support the existence of any clear niche
423 differentiation between both groups. It highlights that nearly no study has compared the responses of
424 N₂O producers and reducers to labile C or ARE-C addition (Table 1), which should be better explored in
425 the future.

426

427 *4.4. Conclusions*

428 Given that denitrification is a major source of N loss and N₂O emission in agroecosystems (Syakila
429 and Kroeze, 2011), it is crucial to avoid practices that may increase this process. Our results, based on a
430 field trial and a growth chamber experiment, show that the inoculation practice can have very strong
431 effects of the activities and abundances of soil NO₂⁻ and N₂O-reducing bacteria, but that the effects vary
432 (and actually can be opposite) according to soil N and moreover C availability. More particularly, we
433 showed that the level of denitrification limitation by C predicts well the resulting effect of inoculation on
434 potential gross and net N₂O production. Inoculation by PGPRs can increase net N₂O production from
435 cropland soils characterized by high C limitation by increasing the abundance of NO₂⁻-reducers more than
436 the abundance of N₂O-reducers. In contrast, inoculation may increase the soil capacity to act as a sink for
437 N₂O for soils where denitrifiers are not heavily limited by C. Because our results are based on potential
438 (gross and net) N₂O production rates and abundances of key denitrifier groups, assessing the actual

439 effect of inoculation on N₂O emissions will require to quantify emission rates from inoculated and non-
440 inoculated plots across soils with contrasted C limitation levels. This could prove crucial for assessing and
441 mitigating the environmental consequences of such agricultural practice.

442

443 **5. Acknowledgements**

444 We acknowledge funding by the French Agency for Research, ANR, to the AZODURE project (ANR-12-
445 AGRO-0008) and to the associated post-doctoral fellowship of AF. We thank all the members from the
446 AZODURE project who participated to the soil sampling campaign, and A. Dumas, A. Frutos and L.
447 Capowiez for realizing CLPP measurements.

448

449 **6. References**

- 450 Altieri, M.A. 1999. The ecological role of biodiversity in agroecosystems. *Agriculture, Ecosystem and*
451 *Environment* 74, 19–31.
- 452 Assemien, F., Cantarel, A., Florio, A., Lerondelle, C., Pommier, T., Gonnety, T.J., Le Roux, X. 2019.
453 Different groups of nitrite-reducers and N₂O-reducers have distinct ecological niches and functional
454 roles in West African cultivated soils. *Soil Biology and Biochemistry* 129, 39–47.
- 455 Attard, E., Recous, S., Chabbi, A., De Berranger, C., Guillaumaud, N., Labreuche, J., Philippot, L., Schmid,
456 B., Le Roux, X. 2011. Soil environmental conditions rather than denitrifier abundance and diversity
457 drive potential denitrification after changes in land uses. *Global Change Biology* 17, 1975–1989.
- 458 Baggs, E.M., 2011. Soil microbial sources of nitrous oxide: recent advances in knowledge, emerging
459 challenges and future direction. *Current Opinion in Environmental Sustainability* 3, 321–327.
- 460 Barot, S., Allard, V., Cantarel, A., Enjalbert, J., Gauffreteau, A., Goldringer, I., Lata, J.C., Le Roux, X.,
461 Niboyet, A., Porcher, E., 2017. Designing mixtures of varieties for multifunctional agriculture with
462 the help of ecology: a review. *Agronomy for Sustainable Development* 37, 13.

463 Barrett, M., Khalil, M.I., Jahangir, M.M.R., Lee, C., Cárdenas, L.M., Collins, G., Richards, K.G., O'Flaherty,
464 V. 2016. Carbon amendment and soil depth affect the distribution and abundance of denitrifiers in
465 agricultural soils. *Environmental Science and Pollution Research* 23, 7899–7910.

466 Bashan, Y. 1999. Interactions of *Azospirillum spp.* in soils: a review. *Biology and Fertility of Soils* 29, 246–
467 256.

468 Baudoin, E., Benizri, E., Guckert, A. 2003. Impact of artificial root exudates on the bacterial community
469 structure in bulk soil and maize rhizosphere. *Soil Biology and Biochemistry* 35, 183–1192.

470 Baudoin, E., Philippot, L., Cheneby, D., Chapuis-Lardy, L., Fromin, N., Bru, D., Rabary, B., Brauman, A.
471 2009. Direct seeding mulch-based cropping increases both the activity and the abundance of
472 denitrifier communities in a tropical soil. *Soil Biology and Biochemistry*, 41, 1703–1709.

473 Bérard, A., Sassi, M.B., Renault, P., Gros, R. 2012. Severe drought-induced community tolerance to heat
474 wave. An experimental study on soil microbial processes. *Journal of soils and sediments*, 12, 513–
475 518.

476 Berks, B.C., Ferguson, S.J., Moir, J.W.B., Richardson, D.J.D. 1995. Enzymes and associated electron
477 transports systems that catalyse the respiratory reduction of nitrogen oxides and oxyanions.
478 *Biochimica and Biophysica Acta* 1232, 97–173.

479 Bounaffaa, M., Florio, A., Le Roux, X., Jayet, P.A. 2018. Economic and environmental analysis of maize
480 inoculation by plant growth promoting rhizobacteria in the French Rhône-Alpes region. *Ecological*
481 *Economics* 146, 334–346.

482 Campbell, C.D., Chapman, S.J., Cameron, C.M., Davidson, M.S., Potts, J.M. 2003. A rapid microtiter plate
483 method to measure carbon dioxide evolved from carbon substrate amendments so as to determine
484 the physiological profiles of soil microbial communities by using whole soil. *Applied and*
485 *Environmental Microbiology* 69, 3593–3599.

486 Chantigny, M.H., Rochette, P., Angers, D.A., Bittman, S., Buckley, K., Massé, D., Bélanger, G., Eriksen-
487 Hamel, N., Gasser, M.O. 2010. Soil nitrous oxide emissions following band-incorporation of fertilizer
488 nitrogen and swine manure. *Journal of environmental quality*, 39, 1545–1553.

489 Chapuis-Lardy, L., Wrage, N., Metay, A., Chotte, J.L., Bernoux, M. 2007. Soils, a sink for N₂O? A review.
490 *Global Change Biology*, 13, 1–17.

491 Clark, I.M., Buchkina, N., Jhurrea, D., Goulding, K.W., Hirsch, P.R. 2012. Impacts of nitrogen application
492 rates on the activity and diversity of denitrifying bacteria in the Broadbalk Wheat Experiment.
493 *Philosophical Transaction of the Royal Society B* 367, 1235–1244.

494 Couillerot, O., Bouffaud, M. L., Baudoin, E., Muller, D., Caballero-Mellado, J., Moëgne-Loccoz, Y. (2010).
495 Development of a real-time PCR method to quantify the PGPR strain *Azospirillum lipoferum CRT1* on
496 maize seedlings. *Soil Biology and Biochemistry* 42, 2298–2305.

497 Čuhel, J., Šimek, M., Laughlin, R.J., Bru, D., Chèneby, D., Watson, C.J., Philippot, L. 2010. Insights into the
498 effect of soil pH on N₂O and N₂ emissions and denitrifier community size and activity. *Applied and*
499 *environmental microbiology* 76, 1870–1878.

500 Dandie, C.E., Burton, D.L., Zebarth, B.J., Henderson, S.L., Trevors, J.T., Goyer, C. 2008. Changes in
501 bacterial denitrifier community abundance over time in an agricultural field and their relationship
502 with denitrification activity. *Applied and environmental microbiology* 74, 5997–6005.

503 Dendooven, L., Duchateau, L., Anderson, J.M. 1996. Gaseous products of the denitrification process as
504 affected by the antecedent water regime of the soil. *Soil Biology and Biochemistry* 28, 239–245.

505 Domeignoz-Horta, L.A., Putz, M., Spor, A., Bru, D., Breuil, M.C., Hallin, S., Philippot, L. 2016. Non-
506 denitrifying nitrous oxide-reducing bacteria - An effective N₂O sink in soil. *Soil Biology and*
507 *Biochemistry* 31, 376–379.

508 Eggermont, H., Balian, E., Azevedo, J. M. N., Beumer, V., Brodin, T., Claudet, J., Fady, B., Grube, M.,
509 Keune, H., Lamarque, P., Reuter, K., Smith, M., van Ham, C., Weisser, W.W. Le Roux, X. 2015.

510 Nature-based solutions: new influence for environmental management and research in Europe.
511 GAIA-Ecological Perspectives for Science and Society, 24, 243–248.

512 El Zemrany, H., Cortet, J., Lutz, M.P., Chabert, A., Baudoin, E., Haurat, J., Maughan, N., Félix, D., Défago,
513 G., Bally, R., Moenne-Loccoz, Y. 2006. Field survival of the phytostimulator *Azospirillum lipoferum*
514 *CRT1* and functional impact on maize crop, biodegradation of crop residues, and soil faunal
515 indicators in a context of decreasing nitrogen fertilisation. Soil Biology and Biochemistry 38, 1712–
516 1726.

517 Enwall, K., Philippot, L., Hallin, S. 2005. Activity and composition of the denitrifying bacterial community
518 respond differently to long-term fertilization. Applied and environmental microbiology 71, 8335–
519 8343.

520 Enwall, K., Throbäck, I.N., Stenberg, M., Söderström, M., Hallin, S. 2010. Soil resources influence spatial
521 patterns of denitrifying communities at scales compatible with land management. Applied and
522 Environmental Microbiology 76, 2243–2250.

523 Fages, J., Mulard, D. 1988. Isolement de bactéries rhizosphériques et effet de leur inoculation en pots
524 chez *Zea mays*. Agronomie 8, 309–312.

525 Fallik, E., Sarig, S., Okon, Y. 1994. Morphology and physiology of plant roots associated with
526 *Azospirillum*. *Azospirillum/plant associations*, 77–85.

527 Florio, A., Pommier, T., Gervais, J., Bérard, A., Le Roux, X. 2017. Soil C and N statuses determine the
528 effect of maize inoculation by plant growth-promoting rhizobacteria on nitrifying and denitrifying
529 communities. Scientific Reports 7, 8411.

530 Giles, M.E., Daniell, T.J., Baggs, E.M. 2017. Compound driven differences in N₂ and N₂O emission from
531 soil; the role of substrate use efficiency and the microbial community. Soil Biology and Biochemistry
532 106, 90–98.

533 Gillam, K.M., Zebarth, B.J., Burton, D.L. 2008. Nitrous oxide emissions from denitrification and the
534 partitioning of gaseous losses as affected by nitrate and carbon addition and soil aeration. *Canadian*
535 *Journal of Soil Science* 88, 133–143.

536 Hallin, S., Jones, C. M., Schloter, M., Philippot, L. 2009. Relationship between N-cycling communities and
537 ecosystem functioning in a 50-year-old fertilization experiment. *The ISME journal* 3, 597.

538 Henderson, S.L., Dandie, C.E., Patten, C.L., Zebarth, B.J., Burton, D.L., Trevors, J.T., Goyer, C. 2010.
539 Changes in denitrifier abundance, denitrification gene mRNA levels, nitrous oxide emissions, and
540 denitrification in anoxic soil microcosms amended with glucose and plant residues. *Applied and*
541 *Environmental Microbiology* 76, 2155–64.

542 Henry, S., Bru, D., Stres, B., Hallet, S., Philippot, L. 2006. Quantitative detection of the *nosZ* Gene,
543 encoding nitrous oxide reductase, and comparison of the abundances of 16S rRNA, *narG*, *nirK*, and
544 *nosZ* genes in soils. *Applied and Environmental Microbiology* 72, 5181–5189.

545 Henry, S., Texier, S., Hallet, S., Bru, D., Dambreville, C., Chèneby, D., Bizouard, F., Germon, J.C., Philippot,
546 L. 2008. Disentangling the rhizosphere effect on nitrate reducers and denitrifiers: insight into the
547 role of root exudates. *Environmental Microbiology* 10, 3082–3092.

548 Heulin, T., Guckert, A., Balandreau, J. 1987. Stimulation of root exudation of rice seedlings by
549 *Azospirillum* strains-carbon budget under gnotobiotic conditions. *Biology and Fertility of Soils* 4, 9-14.

550 Højberg, O., Binnerup, S.J., Sørensen, J. 1996. Potential rates of ammonium oxidation, nitrate reduction,
551 and denitrification in the young barley rhizosphere. *Soil Biology and Biochemistry* 28, 47–54.

552 Iijima, M., Griffiths, B., Bengough, A.G. 2000. Sloughing of cap cells and carbon exudation from maize
553 seedling roots in compacted sand. *New Phytologist* 145, 477–482.

554 Jones, C.M., Graf, D.R., Bru, D., Philippot, L., Hallin, S. 2013. The unaccounted yet abundant nitrous
555 oxide-reducing microbial community: a potential nitrous oxide sink. *The ISME Journal* 7, 417–426.

556 Jusselme, M.D., Saccone, P., Zinger, L., Faure, M., Le Roux, X., Guillaumaud, N., Bernard, L., Clement, J.C.,
557 Poly, F. 2016. Variations in snow depth modify N-related soil microbial abundances and functioning
558 during winter in subalpine grassland. *Soil Biology and Biochemistry* 92, 27–37.

559 Kastl, E.M., Schloter-Hai, B., Buegger, F., Schloter, M. 2015. Impact of fertilization on the abundance of
560 nitrifiers and denitrifiers at the root–soil interface of plants with different uptake strategies for
561 nitrogen. *Biology and fertility of soils* 51, 57–64.

562 Krafczyk, I., Trolldenier, G., Beringer, H. 1984. Soluble root exudates of maize: influence of potassium
563 supply and rhizosphere microorganisms. *Soil Biology and Biochemistry* 16, 315–322.

564 Krause, H. M., Thonar, C., Eschenbach, W., Well, R., Mäder, P., Behrens, S., Kappler, A., Gatterer, A.
565 2017. Long term farming systems affect soils potential for N₂O production and reduction processes
566 under denitrifying conditions. *Soil Biology and Biochemistry* 114, 31–41.

567 Kuzyakov, Y., Xu, X. 2013. Competition between roots and microorganisms for N: mechanisms and
568 ecological relevance. *New Phytologist* 198, 656–669.

569 Langarica-Fuentes, A., Manrubia, M., Giles, M. E., Mitchell, S., Daniell, T. J. 2018. Effect of model root
570 exudate on denitrifier community dynamics and activity at different water-filled pore space levels in a
571 fertilised soil. *Soil Biology and Biochemistry* 120, 70–79.

572 Le Roux, X., Schmid, B., Poly, F., Barnard, R. L., Niklaus, P. A., Guillaumaud, N., Habekost, M., Oelmann, Y.,
573 Philippot, L., Salles, J.F., Schloter, M., Steinbeiss, S., Weigelt, A. 2013. Soil environmental conditions
574 and microbial build-up mediate the effect of plant diversity on soil nitrifying and denitrifying enzyme
575 activities in temperate grasslands. *PLoS One*, 84, e61069.

576 Loick, N., Dixon, E. R., Abalos, D., Vallejo, A., Matthews, G. P., McGeough, K. L., Well, R., Watson, C.J.,
577 Laughlin, R.J., Cardenas, L.M. 2016. Denitrification as a source of nitric oxide emissions from
578 incubated soil cores from a UK grassland soil. *Soil Biology and Biochemistry* 95, 1–7.

579 Ma, W., Jiang, S., Assemien, F., Qin, M., Ma, B., Xie, Z. Liu, Y., Feng, H., Du, G., Ma, X., Le Roux, X. 2016.
580 Response of microbial functional groups involved in soil N cycle to N, P and NP fertilization in Tibetan
581 alpine meadows. *Soil Biology and Biochemistry* 101, 195–206.

582 Mahmood, T., Ali, R., Malik, K.A., Shamsi, S.R.A. 1997. Denitrification with and without maize plants (*Zea*
583 *mays L.*) under irrigated field conditions. *Biology and Fertility of Soils* 24, 323–328.

584 Mantelin, S., Touraine, B. 2004. Plant growth-promoting bacteria and nitrate availability: impacts on root
585 development and nitrate uptake. *Journal of Experimental Botany* 55, 27–34.

586 Miller, M.N., Zebarth, B., Dandie, C.E., Burton, D.L., Goyer, C., Trevors, J.T. 2008. Crop residue influence
587 on denitrification, N₂O emissions and denitrifier community abundance in soil. *Soil Biology and*
588 *Biochemistry* 40, 2553–2562.

589 Miller, M.N., Dandie, C.E., Zebarth, B.J., Burton, D.L., Goyer, C., Trevors, J.T. 2012. Influence of carbon
590 amendments on soil denitrifier abundance in soil microcosms. *Geoderma* 170, 48–55.

591 Morley, N.J., Richardson, D.J., Baggs, E.M. 2014. Substrate induced denitrification over or under
592 estimates shifts in soil N₂/N₂O ratios. *PLoS one* 9, e108144.

593 Mounier, E., Hallet, S., Cheneby, D., Benizri, E., Gruet, Y., Nguyen, C., Piutti, S., Robin, C., Slezack-
594 Deschaumes, S., Martin-Laurent, F., Germon, J. C., Philippot, L. 2004. Influence of maize mucilage on
595 the diversity and activity of the denitrifying community. *Environmental Microbiology* 6, 301–312.

596 Murray, P.J., Hatch, D.J., Dixon, E.R., Stevens, R.J., Laughlin, R.J., Jarvis, S.C. 2004. Denitrification
597 potential in a grassland subsoil: effect of carbon substrates. *Soil Biology and Biochemistry* 36, 545–
598 547.

599 Myers, R.J.K., McGarity, J.W. 1971. Factors influencing high denitrifying activity in the subsoil of
600 solodized solonetz. *Plant and Soil* 35, 145–160.

601 Myrold D.D., Tiedje, J.M. 1985. Establishment of denitrification capacity in soil: Effects of carbon, nitrate
602 and moisture. *Soil Biology and Biochemistry* 17, 819–822.

603 Niboyet, A., Barthes, L., Hungate, B.A., Le Roux, X., Bloor, J.M., Ambroise, A., Fontaine, S., Price, P.M.,
604 Leadley, P.W. 2010. Responses of soil nitrogen cycling to the interactive effects of elevated CO₂ and
605 inorganic N supply. *Plant and soil* 327, 35–47.

606 Patra, A.K., Abbadie, L., Clays-Josserand, A., Degrange, V., Grayston, S.J., Loiseau, P., Louault, F.,
607 Mahmood, S., Nazaret, S., Philippot, L., Poly, F., Prosser, J., Le Roux, X. 2005. Effect of grazing on
608 microbial functional groups involved in soil N dynamics. *Ecological Monographs* 75, 65–80.

609 Philippot, L., Hallin, S., Schloter, M. 2007. Ecology of denitrifying prokaryotes in agricultural
610 soil. *Advances in Agronomy* 96, 249–305.

611 Recous, S., Robin, D., Darwis, D., Mary, B. 1995. Soil inorganic N availability: effect on maize residue
612 decomposition. *Soil Biology and Biochemistry* 31, 1529–38.

613 Sanford, R.A., Wagner, D.D., Wu, Q., Chee-Sanford, J.C., Thomas, S.H., Cruz-García, C., Rodriguez, G.,
614 Massol-Deyá, A., Krishnanif, K.K., Ritalahtig, K.M., Nissen, S., Konstantinidis, K.T., Löffler, F.E. 2012.
615 Unexpected non-denitrifier nitrous oxide reductase gene diversity and abundance in soils.
616 *Proceedings of the National Academy of Sciences* 109, 19709–19714.

617 Sarig, S., Blum, A., Okon, Y. 1988. Improvement of water status and yield of field-grown grain sorghum
618 (*Sorghum bicolor*) by inoculation with *Azospirillum brasilense*. *The Journal of Agricultural Science*
619 110, 271–277.

620 Schaeffer, S.M., Billings, S.A., Evans, R.D. 2003. Responses of soil nitrogen dynamics in a Mojave Desert
621 ecosystem to manipulation in soil carbon and nitrogen availability. *Oecologia* 134, 547–553.

622 Shapleigh, J.P. 2013. Denitrifying prokaryotes. In *The prokaryotes* pp. 405-425. Springer, Berlin,
623 Heidelberg.

624 Shaw, L.J., Morris, P., Hooker, J.E. 2006. Perception and modification of plant flavonoid signals by
625 rhizosphere microorganisms. *Environmental Microbiology* 8, 1867–1880.

626 Stefanson, R.C. 1972. Soil denitrification in sealed soil–plant systems. I. Effect of plant, soil water content
627 and soil organic matter content. *Plant and Soil* 33, 113–127.

628 Syakila, A., Kroeze, C. 2011. The global nitrous oxide budget revisited. *Greenhouse Gas Measurement*
629 *Management* 1, 17–26.

630 Tatti, E., Goyer, C., Zebarth, B.J., Burton, D.L., Giovannetti, L., Viti, C. 2013. Short-term effects of mineral
631 and organic fertilizer on denitrifiers, nitrous oxide emissions and denitrification in long-term amended
632 vineyard soils. *Soil Science Society of America Journal* 77, 113–122.

633 Throbäck, I.N., Enwall, K., Jarvis, Å., Hallin, S. 2004. Reassessing PCR primers targeting *nirS*, *nirK* and *nosZ*
634 genes for community surveys of denitrifying bacteria with DGGE. *FEMS Microbiology Ecology* 49, 401–
635 417.

636 Tiedje, J.M. 1982. Denitrification. *Methods of Soil Analysis. Part 2. Chemical and Microbiological*
637 *Properties*, 1011–1026.

638 Torralbo, F., Menéndez, S., Barrena, I., Estavillo, J.M., Marino, D., González-Murua, C. 2017. Dimethyl
639 pyrazol-based nitrification inhibitors effect on nitrifying and denitrifying bacteria to mitigate N₂O
640 emission. *Scientific reports* 7, 13810.

641 Trofymow, J.A., Coleman, D.C., Cambardella, C. 1987. Rates of rhizodeposition and ammonium depletion
642 in the rhizosphere of axenic oat roots. *Plant and Soil* 97, 333–344.

643 Vinther, F.P., Memon, H.G., Jensen, V. 1982. Populations of denitrifying bacteria in agricultural soil under
644 continuous barley cultivation. *Pedobiologia* 24, 319–328.

645 Weier, K.L., Doran, J.W., Power, J.F., Walters, D.T. 1993. Denitrification and the dinitrogen nitrous oxide
646 ratio as affected by soil water, available carbon, and nitrate. *Soil Science Society of America Journal*
647 57, 66–72.

648 Weixin, C., Coleman, D.C., Carroll, C.R., Hoffman, C.A. 1993. In situ measurement of root respiration and
649 soluble C concentrations in the rhizosphere. *Soil Biology and Biochemistry*, 25, 1189–1196.

- 650 World Reference Base for Soil Resources 2006. A framework for international classification, correlation
651 and communications. World Soil Resources Reports 103
- 652 Zumft, W.G. 1997. Cell biology and molecular basis of denitrification. Microbiology and Molecular
653 Biology Reviews 61, 533–616.

654 **Table 1.** A review of the effects of C and N amendments to soil on the abundances of *nirK*- and *nirS*-nitrite reducers, the abundances of *nosZI*-
655 and *nosZII*-N₂O reducers, and potential gross (PGNP) and net (PNNP) N₂O production rates. Arrows indicate an increase (↑), a decrease (↓) or no
656 change (→) in the variable considered as compared with non-amended control. Two arrows indicate two effects for half of cases each. G and F
657 indicates growth chamber and field studies, respectively.

	Type of amendment	Reference	Land type	Type of experiment	Effects reported on denitrifier abundance and activity					
					<i>nirK</i>	<i>nirS</i>	<i>nosZI</i>	<i>nosZII</i>	PGNP	PNNP
Labile C amendment	Glucose	Myers and McGarity (1971)	Unplanted	G					↑	
	Glucose	Weier et al. (1993)	Unplanted	G					↑	↑
	Glucose	Dandie et al. (2008)	Unplanted	G					↑	
	Glucose	Miller et al. (2008)	Cropland	G					→	↓
	Glucose	Henderson et al. (2010)	Mixed crops	G		→	→		→	→
	Glucose	Miller et al. 2012)	Unplanted	G					→	↑
	Glucose	Barrett et al. (2016)	Unplanted	G		↑				
						-	↑50%	↑50%	-	↑50%
					-	→50%	→50%	-	→50%	→25%
					-	-	-	-	-	↓25%
Mineral N amendment	NO ₃	Weier et al. (1993)	Unplanted	G					→	→
	NO ₃	Enwall et al. (2005); Hallin et al. (2009)	Cropland	F	↑	→	↑		↑	
	NO ₃	Gillam et al. (2008)	Unplanted	G					→	↑
	NO ₃	Miller et al. (2008)	Cropland	G					→	↑
	NO ₃	Miller et al. (2008)	Cropland	G					↑	↑
	NO ₃	Miller et al. (2008)	Cropland	G					↑	↑
	NO ₃ + NH ₄	Niboyet et al. (2010)	Grassland	G					↑	
	NO ₃ + NH ₄	Clark et al. (2012)	Cropland	F	↑	→	→			
	NO ₃ + NH ₄	Clark et al. (2012)	Cropland	F	↑	↓	→			
	NO ₃ + NH ₄	Tatti et al. (2013)	Orchard	G	→	→	→		↑	↑→
	NO ₃ + NH ₄	Tatti et al. (2013)	Orchard	G	→	↑	→		↑	↑→
	NO ₃ + NH ₄	Kastl et al. (2015)	Grasslands	G	→	→	→			
	NO ₃ + NH ₄	Kastl et al. (2015)	Grasslands	G	→	→	→			
	NO ₃ + NH ₄	Kastl et al. (2015)	Grasslands	G	→	→	→			
	NO ₃ + NH ₄	Ma et al. (2016)	Grasslands	F	→	→	→			
	NO ₃ + NH ₄	Florio et al. (2017)	Cropland	F	→	→	→	→	→	
	NO ₃ + NH ₄	Krause et al. (2017)	Cropland	G	→	→	→	→	↑	→

	Type of amendment	Reference	Land type	Type of experiment	Effects reported on denitrifier abundance and activity					
					<i>nirK</i>	<i>nirS</i>	<i>nosZI</i>	<i>nosZII</i>	PGNP	PNNP
Mineral N amendment	NH ₄	Enwall et al. (2005); Hallin et al. (2009)	Cropland	F	↓	↓	→		↑	
	NH ₄	Torralbo et al. (2017)	Cropland	G	→	→	→	↓		
	NH ₄	Torralbo et al. (2017)	Cropland	G	→	→	→	→		
					↑22%	↑7%	↑7%	-	↑67%	↑62%
				→71%	→79%	→93%	→75%	→33%	→38%	
				↓7%	↓14%	-	↓25%	-	-	
Labile C + mineral N amendment	Glucose + NO ₃	Weier et al. (1993)	Unplanted	G					↑	↑
	Glucose + NO ₃	Murray et al. (2004)	Unplanted	G					↑	↑
	Glucose + NO ₃	Gillam et al. (2008)	Unplanted	G					→	→
	Glucose + NO ₃	Miller et al. (2008)	Cropland	G					↑	↑
	Glucose + NO ₃	Miller et al. (2008)	Cropland	G					→	↑
	Glucose + NO ₃	Miller et al. (2008)	Cropland	G					↑	↑
	Glucose + NO ₃	Miller et al. (2008)	Cropland	G					→	↓
	Glucose + NO ₃	Miller et al. (2008)	Cropland	G					↑	→
	Glucose + NO ₃	Miller et al. (2008)	Cropland	G			→		↑	↑
	Glucose + NO ₃	Loick et al. (2016)	Unplanted	G						↑
	Glucose, cellulose + NO ₃	Dendooven et al. (1996)	Unplanted						↑	↑
	Starch + NO ₃	Murray et al. (2004)	Unplanted	G					↑	→
	Cellulose + NO ₃	Murray et al. (2004)	Unplanted	G					→	→
	Organic acids, aminoacids + NO ₃	Morley et al. (2014)	Unplanted	G					↑	↑
Glucose, citric acid, glutamine + NO ₃	Giles et al. (2017)	Unplanted	G					↑	↑	
					-	-	-	↑71%	↑66%	
					-	-	→100%	→29%	→27%	
					-	-	-	-	↓7%	
ARE-C amendment	Mucilage	Mounier et al. (2004)	Unplanted	G					↑	
	Artificial exudates	Henry et al. (2008)	Cropland	G	↑	↑	→		↑	↑
	Artificial exudates	Langarica-Fuentes et al. (2018)	Unplanted	G	↑	↑	↑	→	↑	→
					↑100%	↑100%	↑50%	-	↑100%	↑50%
				-	-	→50%	→100%	-	→50%	
				-	-	-	-	-	-	

661

662 **Table 2.** Overall effects of maize inoculation by *Azospirillum lipoferum* CRT1 on the abundances of *nirK*-, *nirS*- and *nir*- (*nirK+nirS*) NO₂⁻ reducers,
663 the abundances of *nosZI*-, *nosZII*- and *nosZ*- (*nosZI+nosZII*) N₂O reducers, and potential gross (PGNP) and net (PNNP) N₂O production rates.

	<i>nirK</i>	<i>nirS</i>	<i>nir</i>	<i>nosZI</i>	<i>nosZII</i>	<i>nosZ</i>	PGNP	PNNP
Inoculation	NS	NS	NS	NS	NS	NS	NS	NS
C limitation	NS	NS	NS	0.005	NS	0.003	<0.0001	0.056
Inoculation x C limitation	0.026	0.047	0.024	0.060	0.019	0.030	0.035	0.023

664 Results were obtained using two-way ANOVA with inoculation and C limitation as fixed effects.

665

666 **Figure legends**

667

668 **Fig. 1.** We assumed that the effect of cereal inoculation with plant growth promoting rhizobacteria on
669 potential gross and net N₂O production results from the balance between the inoculation-induced increase
670 in root exudation and increased plant-microbes competition for NO₃⁻ which may differently affect the main
671 denitrifier groups. In soils with low limitation of denitrification by organic C, exudation would play a minor
672 role and enhanced competition for nitrate would lead to lower the abundance and activity of NO₂⁻-
673 reducers (*nir*-harbouring bacteria) and N₂O-reducers (*nosZ*-harbouring bacteria). In soils with high limitation
674 of denitrification by C, positive effect of increased exudation would prevail, and increased C availability
675 would lead to higher denitrifier abundance and activity. The sensitivity of the different denitrifier groups, in
676 particular of NO₂⁻- reducers as compared to N₂O-reducers, to C and N availabilities would determine the
677 overall inoculation effect on gross and net N₂O production.

678

679 **Fig. 2.** Relationship between the level of denitrification (*i.e.* potential gross N₂O production) limitation by
680 soil organic carbon, C, and the amount of artificial root exudates-C (ARE-C) added to the microcosms under
681 growth chamber (G) conditions (light and dark grey dots). White symbols corresponding to field (F) plots
682 are presented for comparison. ARE-C₀, ARE-C₂₀, ARE-C₁₀₀, ARE-C₂₅₀ refer to 0, 20, 100 and 250 μg C g⁻¹ soil
683 treatments. f/2, f, org and nf refer to reduced, optimal, organic and no fertilization, respectively.


684

685 **Fig. 3.** Relationship between the inoculation effects on (a) potential gross N₂O production, PGNP, (b)
686 potential net N₂O production, PNNP, (c) the total abundance of NO₂⁻- reducers, and (d) the total abundance
687 of N₂O-reducers, and the level of limitation of denitrification by C. Each point corresponds to the mean of
688 the inoculation effect calculated for each pair of NI-I plots or microcosms as described in section 2.5.
689 Symbols for treatments are as in Fig. 2.

690

691 Fig. 1


692





693

694

695 **Fig. 2.**



696



699 **Supplementary Material**

700

701 **The responses of NO₂⁻- and N₂O-reducing bacteria to maize inoculation by the**
702 **PGPR *Azospirillum lipoferum* CRT1 depend on carbon availability and**
703 **determine gross and net N₂O production**

704

705 by Alessandro Florio, Caroline Bréfort, Jonathan Gervais, Annette Bérard & Xavier Le Roux

706

707 2 Supplementary Tables S1 & S2

708 2 Supplementary Figure S1 & S2.

709 **Table S1.** Potential gross N₂O production (PGNP), gross N₂O production when either C (GNP_C) or N (GNP_N) was not added to flasks, limitation of
710 denitrification by C or N, and potential net (PNNP) N₂O production rates in inoculated (I) and non-inoculated (NI) soils from growth chamber (G) and
711 field (F) experiments. Acronyms for fertilization treatments are as in Fig. 2. Values are means ± s.e. (n=6 or 5 for G and F, respectively).
712


Type of experiment	Inoculation	Treatment	PGNP ($\mu\text{g N}_2\text{O-N g}^{-1} \text{ soil h}^{-1}$)	GNP _C ($\mu\text{g N}_2\text{O-N g}^{-1} \text{ soil h}^{-1}$)	GNP _N ($\mu\text{g N}_2\text{O-N g}^{-1} \text{ soil h}^{-1}$)	Limitation of denitrification by C ((1-GNP _C /PGNP)*100) (%)	Limitation of denitrification by N ((1-GNP _N /PGNP)*100) (%)	PNNP ($\mu\text{g N}_2\text{O-N g}^{-1} \text{ soil h}^{-1}$)
G	NI	ARE-C ₀	3.70±0.18	1.07±0.14	2.00±0.67	70.65±4.04	46.73±16.54	3.40±0.24
		ARE-C ₂₀	4.00±0.16	1.48±0.20	1.55±0.50	62.67±5.17	61.37±12.27	2.13±0.12
		ARE-C ₁₀₀	4.99±0.23	1.94±0.20	0.80±0.20	60.38±5.26	84.35±3.47	4.20±0.18
		ARE-C ₂₅₀	7.07±0.57	4.64±0.62	4.93±0.65	30.56±13.07	27.22±11.67	4.87±0.31
		ARE-C _{100f}	4.76±0.28	1.94±0.18	3.40±0.54	58.23±5.13	27.34±11.44	2.25±0.12
	I	ARE-C ₀	3.66±0.17	1.24±0.08	1.71±0.54			3.45±0.15
		ARE-C ₂₀	3.84±0.18	1.42±0.16	1.38±0.43			2.29±0.04
		ARE-C ₁₀₀	4.58±0.35	1.98±0.18	2.51±1.25			3.46±0.26
		ARE-C ₂₅₀	6.46±0.32	4.36±0.70	5.07±0.92			2.62±0.24
		ARE-C _{100f}	4.74±0.37	2.24±0.27	3.19±0.51			2.47±0.16
F	NI	nf	3.15±0.92	0.98±0.12	2.53±0.65	69.60±7.95	31.24±6.48	1.92±0.52
		f/2	2.63±0.74	0.85±0.07	2.57±0.60	72.45±6.08	35.03±5.71	2.24±0.65
		f	1.98±0.58	1.00±0.16	2.24±0.61	76.48±15.01	29.49±4.66	1.61±0.46
		org	1.54±0.05	0.74±0.03	1.31±0.13	51.43±2.19	18.15±5.19	n.d.
	I	nf	3.49±0.98	1.06±0.14	2.40±0.56			1.82±0.53
		f/2	3.92±0.89	1.08±0.10	2.55±0.46			1.69±0.31
		f	3.63±0.78	0.85±0.19	2.56±0.45			1.49±0.24
		org	1.46±0.06	0.79±0.05	1.17±0.07			n.d.

713 **Table S2.** Abundances of *nirK*-, *nirS*- and *nir* (*nirK+nirS*) NO₂⁻-reducers, and of *nosZI*-, *nosZII*- and *nosZ*- (*nosZI+nosZII*) N₂O-reducers in inoculated (I) and
714 non-inoculated (NI) soils from growth chamber (G) and field (F) experiments. Acronyms for fertilization treatments are as in Fig. 2. Values are means ±
715 s.e. (n=6 or 5 for G and F, respectively).

Type of experiment	Inoculation	Treatment	<i>nirK</i> (<i>nirK</i> copies g ⁻¹ soil)	<i>nirS</i> (<i>nirS</i> copies g ⁻¹ soil)	<i>nir</i> (<i>nirK+nirS</i> copies g ⁻¹ soil)	<i>nosZI</i> (<i>nosZI</i> copies g ⁻¹ soil)	<i>nosZII</i> (<i>nosZII</i> copies g ⁻¹ soil)	<i>nosZ</i> (<i>nosZI+nosZII</i> copies g ⁻¹ soil)
G	NI	ARE-C ₀	1.87E+08±1.31E+07	3.93E+07±7.51E+06	2.26E+08±1.72E+07	2.91E+06±2.17E+05	4.46E+05±9.24E+04	3.36E+06±3.30E+05
		ARE-C ₂₀	1.83E+08±9.11E+06	4.22E+07±5.24E+06	2.25E+08±1.86E+07	2.91E+06±1.73E+05	4.28E+05±6.74E+04	3.34E+06±2.05E+05
		ARE-C ₁₀₀	2.04E+08±1.14E+07	4.82E+07±7.78E+06	2.52E+08±1.08E+07	3.62E+06±2.07E+05	4.62E+05±7.74E+04	4.08E+06±2.12E+05
		ARE-C ₂₅₀	2.25E+08±2.43E+07	5.66E+07±1.10E+07	2.82E+08±5.14E+07	4.44E+06±4.38E+05	4.49E+05±9.75E+04	4.89E+06±3.51E+05
		ARE-C _{100f}	2.01E+08±8.01E+06	4.52E+07±8.31E+06	2.46E+08±1.43E+07	3.24E+06±1.42E+05	3.99E+05±8.24E+04	3.64E+06±2.05E+05
	I	ARE-C ₀	1.82E+08±1.04E+07	5.24E+07±1.29E+07	2.34E+08±1.83E+07	2.37E+06±2.20E+05	3.68E+05±7.49E+04	2.74E+06±1.87E+05
		ARE-C ₂₀	1.94E+08±8.90E+06	4.20E+07±3.48E+06	2.36E+08±1.40E+07	2.48E+06±2.90E+05	4.31E+05±7.75E+04	2.91E+06±2.48E+05
		ARE-C ₁₀₀	2.04E+08±1.23E+07	4.94E+07±9.07E+06	2.53E+08±3.43E+07	2.76E+06±3.20E+05	4.43E+05±1.20E+05	3.20E+06±3.68E+05
		ARE-C ₂₅₀	2.39E+08±1.98E+07	5.57E+07±8.48E+06	2.95E+08±2.59E+07	3.73E+06±3.55E+05	4.53E+05±9.60E+04	4.18E+06±2.88E+05
		ARE-C _{100f}	1.88E+08±5.27E+07	7.14E+07±1.15E+07	2.59E+08±6.90E+07	3.54E+06±7.63E+05	4.01E+05±8.28E+04	3.94E+06±1.87E+05
F	NI	nf	8.45E+06±1.51E+06	1.15E+07±3.03E+06	1.99E+07±3.87E+06	8.54E+05±1.69E+05	1.22E+06±1.55E+05	2.08E+06±1.08E+05
		f/2	1.19E+07±3.44E+06	1.28E+07±3.19E+06	2.47E+07±5.56E+06	9.17E+05±1.88E+05	1.08E+06±1.92E+05	2.00E+06±1.20E+05
		f	9.68E+06±3.95E+06	1.07E+07±2.38E+06	2.04E+07±5.75E+06	8.43E+05±1.73E+05	1.35E+06±3.27E+05	2.20E+06±1.83E+05
		org	7.38E+06±1.86E+06	2.15E+06±2.16E+05	9.53E+06±1.88E+06	5.14E+05±3.77E+04	1.55E+07±1.91E+06	1.62E+07±1.92E+06
		nf	9.42E+06±2.75E+06	1.32E+06±3.42E+06	1.07E+07±5.84E+06	7.56E+05±1.14E+05	1.08E+06±2.16E+05	1.83E+06±1.57E+05
	I	f/2	1.47E+07±2.78E+06	1.56E+07±3.10E+06	3.03E+07±3.99E+06	1.03E+06±1.85E+05	1.06E+06±2.77E+05	2.35E+06±3.10E+05
		f	1.42E+07±3.44E+06	1.45E+07±2.65E+06	2.87E+07±5.70E+06	9.32E+05±1.39E+05	1.20E+06±2.33E+05	2.13E+06±1.22E+05
		org	6.03E+06±8.78E+05	1.95E+06±2.14E+05	7.98E+06±1.17E+06	4.91E+05±6.77E+04	1.47E+07±1.74E+06	1.53E+07±1.79E+06
		nf	9.42E+06±2.75E+06	1.32E+06±3.42E+06	1.07E+07±5.84E+06	7.56E+05±1.14E+05	1.08E+06±2.16E+05	1.83E+06±1.57E+05
		f/2	1.47E+07±2.78E+06	1.56E+07±3.10E+06	3.03E+07±3.99E+06	1.03E+06±1.85E+05	1.06E+06±2.77E+05	2.35E+06±3.10E+05

716


717



718

719 **Fig. S1.** Relationship between (a) the limitation of denitrification by N and soil nitrate content, and

720 between (b) the limitation of denitrification by C and SIR. Symbols for treatments are as in Fig. 2.



721

722 **Fig. S2.** Relationship between the inoculation effects on (a) potential gross N₂O production, PGNP, (b)
 723 potential net N₂O production, PNNP, (c) the total abundance of NO₂⁻-reducers (*nir*), and (d) the total
 724 abundance of N₂O-reducers (*nosZ*), and the level of limitation of denitrification by N. Symbols for
 725 treatments are as in Fig. 2.