



**HAL**  
open science

## The Gauls experienced the Roman Warm Period: Oxygen isotope study of the Gallic site of Thézy-Glimont, Picardie, France

Thibault Clauzel, Pascale Richardin, Jannick Ricard, Yves Le Béchenec,  
Romain Amiot, François Fourel, Brian Phouybanhdyt, Arnauld  
Vinçon-Laugier, Jean-Pierre Flandrois, Christophe Lécuyer

### ► To cite this version:

Thibault Clauzel, Pascale Richardin, Jannick Ricard, Yves Le Béchenec, Romain Amiot, et al..  
The Gauls experienced the Roman Warm Period: Oxygen isotope study of the Gallic site of  
Thézy-Glimont, Picardie, France. *Journal of Archaeological Science: Reports*, 2020, 34, pp.102595.  
10.1016/j.jasrep.2020.102595 . hal-02998511

HAL Id: hal-02998511

<https://univ-lyon1.hal.science/hal-02998511v1>

Submitted on 24 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 The Gauls experienced the Roman Warm Period: oxygen isotope study of  
2 the Gallic site of Thézy-Glimont, Picardie, France

3 Thibault Clauzel<sup>a</sup>, Pascale Richardin<sup>b,b'</sup>, Jannick Ricard<sup>c</sup>, Yves Le Béchenec<sup>c</sup>, Romain  
4 Amiot<sup>a</sup>, François Fourel<sup>d</sup>, Brian Phouybanhdyt<sup>b,e</sup>, Arnauld Vinçon-Laugier<sup>a</sup>, Jean-Pierre  
5 Flandrois<sup>f</sup> and Christophe Lécuyer<sup>a,\*, §</sup>

6  
7 a: Univ Lyon, Univ Lyon 1, ENSL, CNRS, LGL-TPE, F-69622, Villeurbanne, France

8 b: Centre de Recherche et de Restauration des Musées de France C2RMF, Palais du Louvre, Porte des Lions, 14  
9 quai François Mitterrand, 75001 Paris, France

10 b': PRETECH – Préhistoire et Technologie, CNRS UMR7055, Université Paris Nanterre, Nanterre, France

11 c: TRAME (Université de Picardie Jules Verne) & Unité Mixte de Recherche n° 8546, AOrOc, CNRS, ENS  
12 Ulm, Paris, France

13 d: Laboratoire d'Ecologie des Hydrosystèmes Naturels et Anthropisés, CNRS UMR 5023, Université Claude  
14 Bernard, Lyon 1, France

15 e: GEOTRAC - Géochronologie Traceurs Archéométrie LSCE- Laboratoire des Sciences du Climat et de  
16 l'Environnement [Gif-sur-Yvette] : DRF/LSCE

17 f: Université de Lyon, CNRS, UMR 5558, Laboratoire de Biométrie et Biologie Evolutive, Villeurbanne, France

18  
19 \*: also at Institut Universitaire de France

20 §: corresponding author, email address: [christophe.lecuyer@univ-lyon1.fr](mailto:christophe.lecuyer@univ-lyon1.fr)

21  
22 **Abstract (190 words)**

23 The Roman Warm Period (from ≈300 BCE to ≈300 AD) is a climatic optimum, which had a  
24 key role in the development of the Roman civilization. This study provides new Mean Air  
25 Temperatures (MATs) inferred from the oxygen isotope composition of 80 bones and teeth  
26 apatite from 8 humans and 8 animals of the Gallic site of Thézy-Glimont, Picardie, France,  
27 dated between the 3<sup>rd</sup> and 2<sup>nd</sup> century BCE. Various bones from the cephalic, axial and

28 appendicular skeleton of three human individuals were sampled. All bones have similar  
29 phosphate  $\delta^{18}\text{O}$  values for each individual, showing that the oxygen isotope signal of  
30 phosphate groups may be homogeneously recorded in bones. The sampled individuals were  
31 assumed to drink water from the nearby Avre river collecting local meteoric waters. The  
32 MATs were reconstructed combining (1) oxygen isotope fractionation equations between  
33 phosphatic tissues ( $\delta^{18}\text{O}_p$ ) and drinking water ( $\delta^{18}\text{O}_w$ ), and (2) relationships between MATs  
34 and  $\delta^{18}\text{O}_w$ . Depending on the used MAT- $\delta^{18}\text{O}_w$  relationship, the reconstructed MATs are  
35  $10.2^\circ\text{C}\pm 1.4^\circ\text{C}$  and  $10.5^\circ\text{C}\pm 1.7^\circ\text{C}$ , which are comparable to present days ( $10.9^\circ\text{C}$ ). This study  
36 shows that the Gallic civilization also experienced relatively high temperatures since the  
37 initiation of the Roman Warm Period.

38

39 Keywords: oxygen isotopes; human skeleton; climate; Gauls, Roman Warm Period

40

## 41 **1. Introduction**

42

43 Although the Holocene is considered to be a very stable interglacial period, climatic  
44 variations from the centennial- to millennial-scale have been identified (Holzhauser et al.,  
45 2005; Caroli and Caldara, 2007; Lirer et al., 2014; Luterbacher et al., 2016; Cisneros et al.,  
46 2016). These variations are considered to be key elements in the rise and fall of civilizations.  
47 For instance, the Medieval Warm Period witnessed the development of rural populations in  
48 Europe linked to very productive harvests. Cities developed and major cultural innovations  
49 took place during warm periods due to land shortages and increasing food requirements  
50 (Fagan, 2008). On the other hand, at the end of the Late Bronze Age, most eastern  
51 Mediterranean urban centers were either destroyed or abandoned throughout the Middle East  
52 and the Aegean area. Some authors have recently suggested that a centuries-long

53 megadrought caused the widespread collapse of Bronze Age Palatial civilization (Kaniewski  
54 et al., 2010; Drake, 2012). The Roman Warm Period (RWP), also called the Roman Climatic  
55 Optimum or the Roman Humid Period, is a climatic variation of interest clearly identified in  
56 the European record. The RWP is identified as a key period for the development of the  
57 Roman culture all around the Mediterranean basin.

58         The concept of the RWP in Europe was first evoked by Lamb (1995) who provided a  
59 synthesis of archaeological, glaciological and palynological studies. The temporal definition  
60 of the RWP varies from one study to another, which may be due to the diversity of climate  
61 proxies and records in Europe. On average, its period extends between  $\approx$ 300 BCE and  $\approx$ 300  
62 AD. The most direct proxies of the RWP are archaeological testimonies. Pliny the Elder (23-  
63 79 AD) testifies that, contrary to what had been the custom in earlier centuries, vines and  
64 olive trees were cultivated farther north in Italy. Ptolemy of Alexandria (100-168 AD) noticed  
65 that, around 140 AD, it rained every month of the year except August. McCormick et al.  
66 (2012) developed an online database referencing archaeological evidence of climate  
67 conditions during the Roman Period.

68         Although the RWP has been intensively studied in the European record, its climate  
69 characteristics remain unclear. Most of the studies of continental western Europe (essentially  
70 Spain, France and Italy) suggest a warm (e.g. McDermott et al., 2001; Roberts et al., 2001;  
71 Desprat et al., 2003; Holzhauser et al., 2005; Frisia et al., 2005; Berger and Bravard, 2012;  
72 Esper et al., 2012) and humid (e.g. Nesje et al., 2000; Wick et al., 2003; Ferrio et al., 2006;  
73 Magny et al., 2007; Migowski et al., 2006; Zanchetta et al., 2007) climatic episode.

74         All the studies reconstructing the level of the Dead Sea also suggest a humid episode  
75 (Frumkin et al., 1991; Heim et al., 1997; Bookman et al., 2004; Migowski et al., 2006). The  
76 data coming from coastal and marine sediments from all over the Mediterranean basin  
77 highlight a dry climate prevailing during the RWP (Schilman et al., 2001; Marquer et al.,

78 2008; Piva et al., 2008; Combourieu Nebout et al., 2009). This conflicts with the data coming  
79 from continental Europe presented above, and suggests that the Mediterranean basin was  
80 under the influence of a distinct climate regime during this episode.

81 High latitude records show a rather cold episode as documented in Iceland (Dahl-Jensen  
82 et al., 1998; Seidenkrantz et al., 2007). There are some records of a cold climate period in  
83 mid-latitudes, either in continental (Issar and Yakir, 1997; Baier et al., 2004; Mangini et al.,  
84 2005; Büntgen et al., 2011) or oceanic (Bond et al., 2001; Taricco et al., 2009; Wang et al.,  
85 2012; Lirer et al., 2014) environments. These temperature discrepancies may be explained by  
86 the existence of three cold and dry intervals during the RWP, which are named Roman I to III  
87 (Lirer et al., 2014; Margaritelli et al., 2016).

88 Variations of solar irradiance are currently invoked to explain the RWP (Bond et al.,  
89 2001; Jiang et al., 2005; Mangini et al., 2005; Wang et al., 2012). Some studies highlight  
90 correlations between solar activity variations and oceanic (Taricco et al., 2009; Lirer et al.,  
91 2014) or continental (Mangini et al., 2005; Margaritelli et al., 2016) records, suggesting that  
92 variations in solar activity may be the main mechanism responsible for the RWP. Dermody et  
93 al. (2012) have presented a model of a climatic “see-saw” between western/coastal Europe  
94 and Mediterranean/Eastern Europe, where solar activity variations trigger a change in the  
95 position and intensity of jet streams through thermal variations of the North Atlantic Ocean.  
96 However, these interpretations must be taken with caution, as the role of solar activity as a  
97 driver for climate variations is questionable and has been challenged in the last decades (e.g.  
98 Csiro, 1983; Haigh, 2007; Lockwood, 2012; Oldenborgh et al., 2013).

99 All the presented records suffer from significant dating uncertainties and, for most of  
100 them, are not focused on the RWP. No climatic reconstruction coming from human or animal  
101 bones is available so far for this period. This study aims to present stable isotope proxy  
102 records of Mean Air Temperature (MAT) inferred from vertebrate remains. Samples have

103 been collected from an archaeological site, well constrained in space and time, which is  
104 located in the North of France where an especially important lack of data has been noticed.  
105 The studied population belongs to the Gallic civilization. The Gauls, who came from Celtic  
106 populations, occupied most of Western Europe during the Iron Age (800 BCE – 50 BCE).  
107 Between the 2<sup>nd</sup> and 1<sup>st</sup> century BCE, the Roman Empire was involved in territorial conquests  
108 from minor Asia through Northern Africa and Greece to Western Europe, which includes  
109 Gallic territories. Therefore, the study of Gallic archaeological sites may bring information  
110 dealing with the transition from the Iron Age Cold Epoch and the RWP or the first part of the  
111 RWP, until  $\approx 0$  AD when all populations are expected to have been under the control of the  
112 Roman Empire.

113         The climate characteristics of the RWP can be studied at mid-to-low latitudes using the  
114 oxygen isotope composition of meteoric waters ( $\delta^{18}\text{O}_{\text{mw}}$ ), which strongly depends on air  
115 temperature, humidity, the trajectories of humid air masses and the amount of precipitation  
116 (Dansgaard, 1964). Reconstructing the past  $\delta^{18}\text{O}_{\text{mw}}$  variations of meteoric waters permits an  
117 estimation of the past variations of those climate parameters.

118         It is known that the oxygen isotope composition of phosphate group ( $\delta^{18}\text{O}_{\text{p}}$ ) from  
119 mineralized tissues of mammalian individuals (bones, teeth) is related to the oxygen isotope  
120 composition of drinking water ( $\delta^{18}\text{O}_{\text{w}}$ ) (Longinelli, 1984; Luz et al., 1984; Luz and Kolodny,  
121 1985; Kohn, 1996). During pre-industrial times, most of the consumed water came from local  
122 sources (wells, springs, nearby watercourses), which are derived from local meteoric waters  
123 (Daux et al., 2005; Darling, 2006). The contribution of water from food varies from one  
124 species to another, according to the water-turnover of the animal (Kohn, 1996). Water  
125 turnover scales to body-mass for most terrestrial vertebrates (Eberhardt, 1969). Humans are  
126 medium omnivores with moderate water turnover, around 5 to 10% of total body water per  
127 day (Shimamoto and Komiya, 2000), which means that the  $\delta^{18}\text{O}$  of water ingested is strongly

128 and mainly imprinted by the  $\delta^{18}\text{O}$  of drinking water. The cooking operation of brewing or  
129 stewing may lead to an isotopic enrichment of the water contained in food compared to  
130 meteoric water (Brettell et al., 2012), but its effect is not unanimous (e.g. Daux et al., 2008).

131 Therefore, measuring the oxygen isotope composition of phosphatic tissues from  
132 terrestrial mammals allows the  $\delta^{18}\text{O}_{\text{mw}}$  of contemporary meteoric waters to be calculated,  
133 hence climate parameters such as MAT to be quantified. We demonstrate in this study that the  
134 reconstructed MATs at Thézy-Glimont were similar to those of today.

135

## 136 **2. Material and methods**

137

### 138 *2.1. The Gallic site of Thézy-Glimont*

139

140 The Gallic site of Thézy-Glimont is located in the Valley of the Somme, in the region of  
141 Picardie, France, 15 km southeast from the city of Amiens. (Figure 1). The site was excavated  
142 in the context of rescue archaeology by the Service d'Archéologie Préventive Amiens  
143 Métropole (SAAM) between November 2012 and May 2013 (Le Béchenec et al., 2016).

144 Picardie is located at mid-latitude (49°N-50°N) in the northwest of Europe, climatically  
145 under the prevailing fresh and humid westerlies from the North Atlantic Ocean. The climate is  
146 rated as Cfb (warm temperate, fully humid with warm summers) according to the Köppen-  
147 Geiger classification (Kottek et al., 2006) with a yearly average temperature of 10.9°C and a  
148 small seasonality amplitude of  $\pm 5.4^\circ\text{C}$  throughout the years 1980 to 2010 (data from Météo  
149 France<sup>TM</sup>). The yearly precipitation rate is 631 mm, with a small seasonality amplitude of  
150  $\pm 7$  mm, July and December being the two most humid months.

151 Thézy-Glimont (49°48'N – 2°25'E) is located at the bottom of the Avre Valley, which  
152 is a tributary of the Somme river. The Avre river is assumed to have been the drinking water

153 source of the inhabitants of Thézy-Glimont as it is located on the edge of the archaeological  
154 site and no water wells have been observed.

155 The SAAM highlighted 17 rectangular graves during the excavation. The graves contain  
156 horses (*Equus caballus*), bovids (*Bos taurus*) and a human (*Homo sapiens*) for most of them.  
157 Sometimes a pig (*Sus scrofa*) has been found. All deposits seem to be intentional and  
158 organized. They were filled with the same sediment (white chalk) coming from the digging,  
159 whereas the tradition of Gallic funeral practices is usually cremation. The excavation report  
160 chooses to present the graves as “offering graves”, even if their true nature is not fully  
161 understood (Le Béchenec et al., 2016). A total of 8 human, 12 horse, 39 bovid and 3 pig  
162 skeletons were excavated from the 17 graves.

163 All 8 humans, 2 horses, 5 bovids and 1 pig were sampled from this corpus (see Table 1).  
164 A great number of sampled individuals come from TGV 1135 as it was the first grave to be  
165 excavated, the biggest of the archaeological site and the one with the highest number of  
166 individuals. The humans are all young males without any noticeable disease. The human  
167 identified as sample TG-02 is the youngest, around 15 years old, while TG-05 is the oldest  
168 (between 30 and 59 years old). The forensic expertise suggested a quick burying after death  
169 before the rigor mortis happened. Although the

170 Eighty samples (bones and teeth) were collected (Table 1). Bone samples were collected  
171 using a toothless diamond saw blade fitted to a Dremel<sup>TM</sup> drill. Various elements coming  
172 from the axial, cephalic and appendicular skeletons of TG-01, TG-02 and TG-03 were  
173 sampled along with 19 bones for TG-01, 14 for TG-02 and 19 for TG-03, respectively (Figure  
174 2). The bone samples were first abraded using a surgical knife in order to remove the altered  
175 layer at the surface and grossly crushed using a hammer. All samples were washed with  
176 acetone to degrease them, rinsed three times with Ultrapure<sup>TM</sup> water and dried in an oven  
177 under medium vacuum, at 60°C, for 24 hours. Clean samples were then crushed in fine



178 powder using a Pulverisette 7™ planetary ball mill composed of zirconium oxide. Teeth were  
179 extracted from the skulls by gently pulling them from their sockets. They were rinsed with  
180 Ultrapure™ water and cleaned using a toothbrush. The powder was obtained by drilling the  
181 enamel of the teeth with a Dremel™ diamond-head drill bit.

182

## 183 2.2. *Dating of the archaeological site*

184

185 During the excavation of the archaeological site, Beta Analytics™ realized <sup>14</sup>C datings  
186 on 13 samples, giving dates with 68.3% probability (1σ) between 200 BCE and 5 AD. New  
187 <sup>14</sup>C dates were realized on collagen extracted from 7 human, 2 horse and 1 bovid bones. The  
188 protocol of the extraction of soluble collagen was based on the method described by Longin  
189 (1971) and has been previously described in detail by Richardin et al. (2017).

190 All measurements were performed with Accelerator Mass Spectrometry techniques  
191 from the Artemis AMS facility at Saclay (France) (Moreau et al., 2013). Calendar dates were  
192 determined using the OxCal v.4.2 procedure and the most recent calibration curve data for the  
193 Northern Hemisphere, IntCal13 (Reimer et al., 2013). Calibration of radiocarbon ages was  
194 performed considering that the contribution of marine products to the diet of the Gauls  
195 remained marginal. Calibrated date ranges (Table 1) correspond to 95.4% probability (2σ).  
196 The archaeological site is dated with 95.4% certainty between 350 BCE and 60 BCE with the  
197 highest probability being between 350 BCE and 200 BCE, one century earlier than previously  
198 proposed. There are no obvious differences among and between the humans and the animals,  
199 therefore all the graves seem contemporary, their digging possibly spread over a century.

200

## 201 2.3. *Bone turnover and isotope homogeneity*

202

203           The bone tissue is vascularized, composed at 60 to 70% of hydroxyapatite crystals  
204 tightly inter-grown with an organic matrix predominantly composed of type I collagen for 20  
205 to 30%. Several ion substitutions can be observed in the crystalline matrix (LeGeros, 1981;  
206 Wopenka and Pasteris, 2005). Enamel from teeth has much larger bioapatite crystals and a  
207 very low organic content, which makes it less sensitive to diagenesis (Tütken and  
208 Vennemann, 2011). Bone and tooth formation starts in the embryo, but human teeth complete  
209 accretion long before bone growth has finished. The human permanent dentition is expected  
210 to be fully formed at about 12 years, while a complete bone structure can be achieved at the  
211 age of 20. Tooth enamel and dentine grow by accretion and do not renew afterwards. On the  
212 contrary, there is a set-up of re-absorption of old bone (both organic and inorganic matrix)  
213 and deposition of new bone tissue. At maturity, this mechanism happens to be continuous and  
214 the bone is said to “turnover”. For a young adult, the total bone tissue is expected to turnover  
215 every five to six years, although it may depend on the bone region or the individual health  
216 status (Glimcher, 2006). Therefore, the stable isotope composition of the bone tissue only  
217 represents the last 5 to 10 years of the individual life, while that of a tooth will represent the  
218 first 12 years of life. Some studies use the discrepancy of bone turnover to highlight a change  
219 of the diet (e.g. Lamb et al., 2014).

220           Because of poor conservation of the material in the field, the lack of articulations  
221 between bones or the preciousness of samples, most isotope studies on bone material are realized  
222 on one bone (generally the femur or the rib) and assume the value is representative of the  
223 whole individual characteristics (e.g. Prowse et al., 2004; Jay and Richards, 2006). However,  
224 if bone turnover is highly variable at the intra- and inter-individual scale, we can wonder to  
225 what extent the isotope study of one bone is representative of the whole skeleton? To  
226 investigate this issue, several parts (bones and teeth) of the same skeleton have been sampled  
227 from the three best conserved and most complete skeletons of human individuals (TG-01, TG-

228 02 and TG-03) excavated in Thézy-Glimont (see Figure 2). Intra-comparisons of the oxygen  
229 isotope composition of mineralized tissues were realized for each individual.

230

#### 231 2.4. *Oxygen isotope ratios of apatite phosphate*

232

233 Oxygen isotopic analyses were performed with stable isotope mass spectrometry  
234 techniques. Phosphate ions were extracted from the apatite using a wet chemistry procedure.  
235 After the dissolution of about 30 mg of each sample in 2 mL of 2M HF at ambient  
236 temperature for 24 hours, the CaF<sub>2</sub> precipitate was separated from the sample solution by  
237 centrifugation. The phosphate ions were neutralized with KOH and isolated using a strong  
238 anionic resin (Amberlite IRN 78) before being precipitated as silver phosphate following the  
239 method first described by Crowson et al. (1991) and slightly modified by Lécuyer et al.  
240 (1993). The oxygen isotope data were obtained using a high-temperature pyrolysis continuous  
241 flow technique developed by Lécuyer et al. (2007) and Fourel et al. (2011). For each sample,  
242 five aliquots of 400–500 µg of silver phosphate were mixed with 500 µg of carbon loaded into  
243 3.5x5 mm silver foil isotope grade capsules (Eurovector<sup>TM</sup>). Pyrolysis was performed at  
244 1450°C using a PYROcube<sup>TM</sup> elemental analyzer (Elementar GmbH-Germany) interfaced in  
245 continuous flow mode with an isotopic ratio mass spectrometer Isoprime<sup>TM</sup> (Elementar UK  
246 Ltd UK). Measurements were performed with CO gas separated by purge and trap technology  
247 within the PYROcube<sup>TM</sup> system at the Laboratoire de Géologie de Lyon and at the  
248 Laboratoire d'Ecologie des Hydrosystèmes Naturels et Anthropisés, both hosted by the  
249 Université Claude Bernard Lyon 1. The official unit for isotopic ratios is the Urey (Ur),  
250 although the unit commonly used is the permil (‰, 1‰ = 1mUr). For practical reasons, data  
251 are reported as δ<sup>18</sup>O values with respect to V-SMOW in ‰ units.

252 Oxygen isotope compositions were drift corrected to account for instrumental  
253 variations. Results were calibrated against silver phosphate samples of known  $\delta^{18}\text{O}$  values  
254 precipitated from the NBS120c standard for which the  $\delta^{18}\text{O}$  value was fixed at 21.7‰ (V-  
255 SMOW) (Lécuyer et al., 1993; Chenery et al., 2010; Halas et al., 2011), as well as with  
256 NBS127 (barium sulfate) having a certified value of 9.3‰ (V-SMOW) (Hut, 1987). NBS120c  
257 standards were converted into silver phosphate along with the samples from each chemistry  
258 batch and considered as “internal standards”. They have  $\delta^{18}\text{O}$  values that range from 21.4‰  
259 to 22.2‰ with a mean value of  $21.76 \pm 0.25$ ‰ ( $n=12$ ). The analytical uncertainty for samples  
260 is always lower than 0.3‰ ( $2\sigma$ ).

261

### 262 **3. Results**

263

#### 264 *3.1. Intra-skeletal variability of the $\delta^{18}\text{O}_p$ from human bones*

265

266 The intra-skeletal variability of the  $\delta^{18}\text{O}_p$  values from the three human individuals  
267 TG-01, TG-02, TG-03 range from 16.1‰ to 16.9‰ with a mean value of 16.6‰ for TG-01,  
268 from 17.4‰ to 17.9‰ with a mean value of 17.6‰ for TG-02 and from 16.4‰ to 17.3‰  
269 with a mean value of 16.8‰ for TG-03 (Figure 2). The standard deviation, representing the  
270 intra-skeletal variability of oxygen isotope composition, is respectively 0.2‰, 0.2‰ and  
271 0.3‰ for TG-01, TG-02 and TG-03. Local deviations of bones  $\delta^{18}\text{O}_p$  from the average  
272 skeleton  $\delta^{18}\text{O}_p$  value has been calculated for each individual as follows:

273

$$\delta^{18}O_{skeleton}^{bone}(\text{‰}) = \delta^{18}O_{V-SMOW}^{bone} + \delta^{18}O_{skeleton}^{V-SMOW} + \delta^{18}O_{V-SMOW}^{bone} \times \delta^{18}O_{skeleton}^{V-SMOW} \times 10^{-3}$$

274

275 No localized and repeatable enrichment or depletion in  $^{18}\text{O}$  relative to the average  $\delta^{18}\text{O}_p$   
276 value is observed among the three individuals.

277

### 278 3.2. *Oxygen isotope values of phosphatic tissues*

279

280 The oxygen isotope compositions of apatite phosphate from the 80 samples were  
281 analyzed (Figure 3, Supplementary Table 1). The  $\delta^{18}\text{O}_p$  values of human bones and teeth  
282 range from 15.1‰ to 18.0‰. Teeth are systematically enriched in  $^{18}\text{O}$  relative to bones by  
283 respectively 0.9‰, 0.3‰ and 0.9‰ for TG-01, TG-02 and TG-03. Human bones have a mean  
284  $\delta^{18}\text{O}_p$  value of 16.2‰ with a standard deviation of 0.8‰. TG-02, who is the youngest  
285 analyzed individual, is significantly enriched by 1.3‰ compared to the other humans. The  
286 analyzed pig has a mean  $\delta^{18}\text{O}_p$  value of 15.6‰ with a standard deviation of 0.5‰. The bones  
287 from 5 bovids have a mean  $\delta^{18}\text{O}_p$  value of 17.3‰ with a standard deviation of 0.2‰, while  
288 the two teeth from individuals TG-B4 and TG-B5 have a mean value of 16.1‰ and a standard  
289 deviation of 0.1‰. Teeth are significantly depleted in  $^{18}\text{O}$  by 1.3‰ relative to bones. No  
290 significant isotopic difference is observed relative to sex (TG-B1 versus the others) or to the  
291 grave the individuals were buried in (TG-B5 versus the others). Molars from the two horses  
292 have a mean value of 15.8‰ with a standard deviation of 0.6‰. In the case of the horse TG-  
293 CH1, the bone  $\delta^{18}\text{O}_p$  value is 1.4‰ higher than that of the tooth. The bones of herbivores  
294 (horse, bovid) have a mean  $\delta^{18}\text{O}_p$  of 17.5‰, while the average  $\delta^{18}\text{O}_p$  for omnivores (human,  
295 pig) is 15.9‰.

296

## 297 4. Discussion

298

### 299 4.1. *Sample preservation and diagenesis*

300

301 The bones from Thézy-Glimont showed a visible brownish layer of alteration, which  
302 was removed using a surgical knife. Some of the human teeth showed attrition, which is a  
303 common feature for the gallic civilization. It may be due to the habit of eating gritter bread  
304 containing sediment particles. Attrition may have altered mechanically the teeth without  
305 scrambling the isotopic signal recorded in enamel.

306 In apatite, P-O bond is known to be very resistant to inorganic isotope exchange with  
307 aqueous fluids at surface temperatures (e.g. Blake et al., 1997; Lécuyer et al., 1999). No trace  
308 of microbially induced diagenesis has been detected. An expected offset of the  $\delta^{18}\text{O}_p$  signal  
309 between teeth and bones from the same individual has been noticed (Touzeau et al., 2013;  
310 Martin et al., 2017). All three individuals TG-01, TG-02 and TG-03 have very low  $\delta^{18}\text{O}_p$   
311 intra-skeletal variability (Figure 2). When considering all the measurements from each  
312 skeleton the three standard deviations of 0.2‰, 0.2‰ and 0.3‰, respectively, are in the same  
313 order of magnitude as the analytical uncertainty ( $\pm 0.3\%$ ). No noticeable and localized  
314 isotopic enrichment or depletion relative to the average  $\delta^{18}\text{O}_p$  value has been noticed. All  
315 these observations support the hypothesis of the preservation of the pristine oxygen isotope  
316 compositions of phosphatic tissues.

317

#### 318 4.2. *Homogeneity of oxygen isotope values in human skeleton apatite phosphate*

319

320 The mapping of the oxygen isotope composition of human bone apatite at the skeletal  
321 level of observation (Figure 2) revealed that, in the case of  $\delta^{18}\text{O}_p$  values, the study of one  
322 selected bone is representative of the composition of the whole individual. We observe an  
323 isotopic homogeneity at the skeleton scale, although the bones have various turnover rates.  
324 Two complementary explanations are proposed: TG-01, TG-02 and TG-03 being young, they

325 might have only realized one cycle of bone turnover at most and their diet did not change  
326 during this interval of time. As migrations can be recorded by apatite  $\delta^{18}\text{O}_p$  values (e.g.  
327 Lightfoot and O'Connell, 2016), we also propose that they were sedentary and lived all their  
328 life in Picardie, France, which could be confirmed with supplementary analysis such as the  
329 use of strontium isotopes. The oxygen isotope variability inside a bone has not been  
330 investigated, but the epiphysis of some bones (e.g. the humerus) and the diaphysis of others  
331 (e.g. the tibia) were analyzed and no noticeable offset was detected. With caution, following  
332 studies of the oxygen isotope composition of bones in archaeological sedentary populations  
333 may assume that one bone is representative of the whole individual. The  $\delta^{18}\text{O}_p$  value of  
334 vertebrates is also dependent on body temperature (Kohn, 1996) and has been used to study  
335 the thermophysiology of terrestrial (Barrick and Showers, 1994, 1995; Amiot et al., 2006) and  
336 marine vertebrates (Bernard et al., 2010; Séon et al., 2020). Thermal variations (or regional  
337 heterothermy), which exist at a human body scale, are not documented at the light of our  
338 study of the oxygen isotope ratios of phosphatic tissues.

339

#### 340 4.3. *The $\delta^{18}\text{O}_p$ record in teeth and bones from individuals of Thézy-Glimont*

341

342 There is a very low  $\delta^{18}\text{O}_p$  variability observed between humans and animals from  
343 Thézy-Glimont. The  $\delta^{18}\text{O}_p$  values from herbivore bones are on average 1.7‰ higher than  
344 those from omnivorous individuals. This can be explained by the diet of the herbivores  
345 mainly eating grass and leaves containing water which is  $^{18}\text{O}$ -enriched in relation to the plant  
346 evaporative system (Barbour, 2007). Comparing the oxygen isotope value from herbivores  
347 and omnivores (mainly humans) enables to address the question of the influence of food  
348 processing on  $\delta^{18}\text{O}_p$ . Although no information is available about the dietary practices on this  
349 archaeological site, stewing or brewing may be detected through an isotopic enrichment of

350 body water later translated in a higher  $\delta^{18}\text{O}_p$  of human bones compared to herbivores (Brettell  
351 et al., 2012). This effect is not detected in this study, which leads to two possible conclusions:  
352 either the inhabitants of Thézy-Glimont had dietary practices which did not include brewing  
353 or stewing, or these cooking processes are not detected in the oxygen isotope composition of  
354 bone apatite. This latter conclusion agrees the works of Daux et al., (2008) on modern human  
355 teeth samples and Chesson et al. (2010) on the similarity of oxygen isotope values of different  
356 modern beverages and meteoric waters in a same location.

357 The  $\delta^{18}\text{O}_p$  value of human teeth enamel is higher than the  $\delta^{18}\text{O}_p$  value of bones,  
358 especially for TG-01 (+0.9‰) and TG-03 (+0.9‰). These values are comparable to the offset  
359 between infant and adult bones of  $\approx +1.2\text{‰}$  found by Britton et al. (2015) which can be  
360 attributed to a specific diet during childhood. A large part of the child's diet consists of milk,  
361 which is  $^{18}\text{O}$ -enriched from the mother's body water. The second and third molars,  
362 respectively M2 and M3, are considered post-weaning teeth and do not record any breast-  
363 feeding. Although the M3 is not discernable from the oxygen isotope value of bones from  
364 TG-02 and is  $^{18}\text{O}$ -depleted compared to the two other teeth for TG-03, the M2 has similar  
365 values than pre-weaning teeth (M1 for TG-01 and PM1 for TG-03). On the contrary,  
366 herbivore teeth are depleted in  $^{18}\text{O}$  compared to bones:  $-1.3\text{‰}$  for bovids and  $-1.4\text{‰}$  for the  
367 horse TG-CH1. Herbivore hypsodont teeth, which grow continuously and are often used to  
368 study climate seasonality (e.g. Bernard et al., 2009), record a signal which is more limited in  
369 time than bones. For the last decade, the average interannual change in yearly MAT of  
370 Picardie, France, is  $\pm 1^\circ\text{C}$  (data from historique-meteo.net) and can go as high as  $2 - 2.5^\circ\text{C}$   
371 (for instance between 2013 and 2014). Translated into  $\delta^{18}\text{O}$  of local meteoric waters, this  
372 interannual change is  $\pm 0.6\text{‰}$  on average and can go as far as a  $\pm 1.6\text{‰}$ , which is in the range  
373 of the teeth-bones offset noticed for the herbivores. We propose that herbivore teeth recorded  
374 more time-limited interval and colder climatic conditions, hence presenting an offset from the



375 bones  $\delta^{18}\text{O}$  record, which represents a pluriannual time scale. Bones are a better material to  
376 reconstruct MAT on the pluriannual time scale than teeth which record either a drinking water  
377 environmental signal mixed with breastfeeding in the case of humans or a time-limited signal  
378 in the case of herbivores. Therefore, only the oxygen isotope values coming from bones are  
379 considered for the calculation of  $\delta^{18}\text{O}_w$  of drinking water and MATs reconstruction.

380 TG-02 is  $^{18}\text{O}$ -enriched compared to all the other humans and there is no significant  
381 difference between the  $\delta^{18}\text{O}_p$  of his bones and his teeth (see Figure 3). This isotopic offset can  
382 be attributed to his young age (estimated at 15 years) keeping in mind that the set-up of bone  
383 turnover is only achieved around 20 years-old. The bones from TG-02 might still record a  
384 residual signal from the breast-feeding period, mixed with a signal coming from drinking  
385 water. Therefore, this individual was removed from the reconstruction of the  $\delta^{18}\text{O}$  of  
386 environmental drinking water.

387 The oxygen isotope composition of phosphatic tissues from individuals with a medium  
388 to low water turnover is mostly representative of the  $\delta^{18}\text{O}$  of the drinking water. Thézy-  
389 Glimont is located in a valley bottom, a few meters away from the Avre river, which is  
390 assumed to be the main drinking water source of local inhabitants. The  $\delta^{18}\text{O}_p$  record is,  
391 therefore, representative of the climate parameters linked to the Avre river. The relationship  
392 between the  $\delta^{18}\text{O}_p$  of phosphatic tissues and the  $\delta^{18}\text{O}_w$  of drinking water has been determined  
393 empirically in numerous groups of vertebrates. These oxygen isotope fractionation equations  
394 mainly depend on the basal metabolism, body mass and diet (herbivorous vs. carnivorous vs.  
395 omnivorous) of species. Several equations have been used in this study and are represented in  
396 Figure 4 as follows:

397

398 Daux et al. (2008) for humans:

399

$$\delta^{18}O_w(\pm 1.51\text{‰}) = 1.54 \times \delta^{18}O_p - 33.72$$

400

401 Longinelli (1984) for pigs:

402

$$\delta^{18}O_w = \frac{\delta^{18}O_p - 22.71}{0.86}$$

403

404 D'Angela and Longinelli (1990) for bovids:

405

$$\delta^{18}O_w = \frac{\delta^{18}O_p - 24.90}{1.01}$$

406

407 Bryant et al. (1994) for equids:

408

$$\delta^{18}O_w(\pm 2.46\text{‰}) = \frac{\delta^{18}O_p - 22.90}{0.6863}$$

409

410 The errors associated with the linear regression of empirical data ( $y = mx + b$ ) have  
411 been computed as the mean standard error for the “y” value when not available in the original  
412 publication and are  $\pm 0.2\text{‰}$  for the equation of Longinelli (1984) and  $\pm 0.1\text{‰}$  for the equation  
413 of D'Angela and Longinelli (1990). We highlight that these uncertainties may be  
414 underestimated as both of the isotope fractionation equations have been established with a  
415 very low number of data, respectively 7 average  $\delta^{18}O$  values for Longinelli (1984) and 5  
416 average  $\delta^{18}O$  values for D'Angela and Longinelli (1990).

417 Using these equations,  $\delta^{18}O_p$  were converted into  $\delta^{18}O_w$  values (Figure 4,  
418 Supplementary Table 2), which were interpreted as reflecting the oxygen isotope composition  
419 of the Avre river during the 2<sup>nd</sup> to 3<sup>rd</sup> century BCE. The mean  $\delta^{18}O_w$  inferred from humans

420 and animals is  $-8.4\text{‰}$ , with a standard deviation of  $\pm 1\text{‰}$ . The propagation of uncertainty,  
421 represented as error bars in Figure 4, have been calculated as the quadratic sum of the  
422 analytical uncertainty, errors associated with the equations and variability in the case of  
423 several measurements obtained for a single individual (TG-01, TG-02, TG-03 and TG-C1).  
424 Taking into account these uncertainties, all the  $\delta^{18}\text{O}_w$  values are homogenous and we can  
425 affirm that they represent one main water source.

426 The present-day isotopic composition of the Avre river water was measured by Négrel  
427 and Petelet-Giraud (2005) twice in a year (Avril and October 2001), with respective  $\delta^{18}\text{O}_w$   
428 values of  $-6.86\text{‰}$  and  $-7.00\text{‰}$ . The  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  values of the Avre river match the Global  
429 Meteoric Water Line (Craig, 1961) as well as the Brest-Orléans isotopic rainwater domain,  
430 confirming that the signal from the Avre river reflects the composition of local meteoric  
431 waters. The oxygen isotope composition of meteoric waters is dependent on temperature, rate  
432 of precipitation and the trajectory of humid air masses (Craig and Gordon, 1965; Rozanski et  
433 al., 1993; Gat, 1996). During the last two millennia, the climatic moisture delivered through  
434 Europe has been driven by the North Atlantic Oscillation (NAO) (Smith et al., 2016) and it is  
435 reasonable to consider that there has not been any major change in the balance between the  
436 water cycle reservoirs for this interval of time (Darling, 2006). For mid- to high-latitudes,  
437 rough linear correlations are observed between the  $\delta^{18}\text{O}$  of meteoric water and air temperature  
438 (e.g. Dansgaard, 1964; Rozanski et al., 1992; Fricke and O'Neil, 1999). Such correlations  
439 result from isotopic compositions that obey a Rayleigh distillation law in both evolving  
440 reservoirs of vapor water and condensed water. Condensation in clouds globally proceeds  
441 from warm low latitudes toward cold high latitudes. Air temperature in the residual clouds  
442 tends to decrease resulting in an increasing isotopic difference between precipitation and  
443 water vapor.

444 For Europe, the slope between  $\delta^{18}\text{O}_{\text{mw}}$  and Mean Air Temperature (MAT) is considered  
445 to be 0.58‰ per °C (Rozanski et al., 1992). We combine the  $\delta^{18}\text{O}_w$  (now assumed to be local  
446  $\delta^{18}\text{O}_{\text{mw}}$ ) obtained from measurements performed on individuals from Thézy-Glimont to  
447 relationships between  $\delta^{18}\text{O}_{\text{mw}}$  and MATs. Several relationships (Dansgaard, 1964; Rozanski  
448 et al., 1992; Fricke and O’Neil, 1999; von Grafenstein et al., 1996; Lécuyer, 2013) are  
449 available in the literature. These different equations may provide sizeable differences in the  
450 calculated air temperature (Daux et al., 2005). We chose two equations we considered the  
451 most appropriate in the framework of this study. First, the equation provided by von  
452 Grafenstein et al. (1996), which has been established at Lake Ammersee, Bavaria, Germany  
453 (48°00’N - 11°07’E), only located  $\approx 2^\circ$  of latitude south of Thézy-Glimont. The equation is  
454 expressed as follows:

$$MAT (^{\circ}C) = \frac{\delta^{18}O_{mw} + 14.48}{0.58 (\pm 0.11)} \quad (1)$$

456  
457 Secondly, we also considered the isotope fractionation equation established by Lécuyer  
458 (2013) as it has been constructed from European meteorological stations excluding those from  
459 the Mediterranean basin, which is under the influence of a specific climatic mode (Lécuyer et  
460 al., 2018). The equation is expressed as follows:

$$MAT (^{\circ}C) = 1.41 (\pm 0.09) \times \delta^{18}O_{mw} + 22.02 (\pm 0.83) \quad (2)$$

461  
462  
463  
464 A reconstruction of mean annual  $\delta^{18}\text{O}_{\text{mw}}$  value at the location of Thézy-Glimont (49°N,  
465 2°E) using The Online Isotopes in Precipitation Calculator (Bowen and Revenaugh, 2003;  
466 IAEA/WMO, 2015; Bowen, 2020) gives a value of  $-6.7 \pm 0.2\text{‰}$ , which is very close to the

467 April (-6.86‰) and October (-7.00‰)  $\delta^{18}\text{O}_w$  values measured on the Avre river by Négrel  
468 and Petelet-Giraud, (2005). We emphasize that at mid latitudes, sinusoidal-like seasonality in  
469 meteoric water  $\delta^{18}\text{O}_{\text{mw}}$  values mathematically leads to spring and autumn  $\delta^{18}\text{O}_{\text{mw}}$  values  
470 matching the mean annual one. Therefore, we use these inter-season oxygen isotope values to  
471 test both  $\delta^{18}\text{O}_{\text{mw}}$ -MAT equations for the Picardie region. The calculated temperatures are  
472  $13.1\pm 1.6^\circ\text{C}$  and  $12.3\pm 1.6^\circ\text{C}$  in April and  $12.9\pm 1.6^\circ\text{C}$  and  $12.1\pm 1.6^\circ\text{C}$  in October for equations  
473 (1) and (2), respectively. The isotope data from April 2001 may be biased as floods happened  
474 in the Somme department during this month, which led to an  $^{18}\text{O}$  enrichment of the Avre river  
475 water sampled for the study. As far as October data are concerned, the calculated  
476 temperatures are near the average temperature of  $11.6^\circ\text{C}$  for this month (data from Météo  
477 France<sup>TM</sup>). Those results indicate that both equations are reliable to infer MAT from  $\delta^{18}\text{O}_{\text{mw}}$   
478 and could safely be applied to the late Iron Age.

479 When applying equations (1) and (2) to the calculated  $\delta^{18}\text{O}_{\text{mw}}$ , the reconstituted air  
480 temperatures for the late Iron Age are on average  $10.5^\circ\text{C}\pm 1.7^\circ\text{C}$  for equation (1) and  
481  $10.2^\circ\text{C}\pm 1.4^\circ\text{C}$  for equation (2) (Figure 5). Therefore, the results for both of these equations  
482 being very close, those calculated MATs are ascribed to Thézy-Glimont during the 3<sup>rd</sup> to 2<sup>nd</sup>  
483 century BCE. These calculated MAT well compare the present-day average temperature of  
484  $10.9^\circ\text{C}$  recorded in Amiens, Picardie. Our result implies that mid-latitudes also experienced a  
485 climate optimum that can be attributed to the RWP, with continental temperatures close to  
486 present-days. This MAT is consistent with results obtained from other studies realized at  
487 mid-latitudes, such as palynological data from Tinner et al. (2003), sedimentological data  
488 from Holzhauser et al. (2005) and geochemical data obtained from speleothems by  
489 McDermott et al. (2001).

490 We assign such mild air temperatures to what is considered to be the initiation of the  
491 RWP. The timing of the RWP is still unsure as well as its climatic characteristics. There may

492 always remain some discrepancies in the European temperature record for the RWP because  
493 no global coherent warm period has occurred on Earth (Neukom et al., 2019) excluding the  
494 current climate change. Nonetheless, this study provides a new climatic anchor point which is  
495 well constrained in space and time, in a region lacking climatic data for the RWP.

496

## 497 **5. Conclusion**

498

499 The Gallic site of Thézy-Glimont provides an insight into the climatic conditions  
500 occurring during the 3<sup>rd</sup> to 2<sup>nd</sup> century BCE in the North Gaul. An oxygen isotope mapping of  
501 three human skeletons revealed a great homogeneity of  $\delta^{18}\text{O}_p$  values within the range of  
502 analytical uncertainty. This result is of paramount importance in archaeological studies as it  
503 means that, in the case of  $\delta^{18}\text{O}_p$  values of sedentary populations, the use of one bone as  
504 representative of the whole skeleton is justified.

505 The  $\delta^{18}\text{O}_p$  of bones from Gauls and associated animals record the  $\delta^{18}\text{O}$  value of drinking  
506 water, which illustrates the climatic conditions prevailing in Picardie, France. The  
507 reconstructed air temperatures are  $10.2^\circ\text{C}\pm 1.4^\circ\text{C}$  and  $10.5^\circ\text{C}\pm 1.7^\circ\text{C}$  depending on the selected  
508  $\delta^{18}\text{O}_{\text{mw}}$ -MAT equation. Here, we emphasize that those estimated mean air temperatures are  
509 similar to present-day ones. Mild air temperatures recorded in Picardie, France, during the  
510 beginning of the RWP is a new result that needs to be taken into consideration for  
511 understanding the climatic regime of the RWP in Europe. This study also shows that  
512 archaeological artefacts may provide quantified climatic anchor points with a high spatial and  
513 temporal resolution, playing a key role in our knowledge of the Late Holocene short-term  
514 climatic changes.

515

516 **Declaration of competing interest**

517

518 The authors declare that they have no known competing financial interests or personal  
519 relationships that could have appeared to influence the work reported in this paper.

520

## 521 **Acknowledgments**

522

523 This study was funded by the Centre National de la Recherche Scientifique and the Institut  
524 Universitaire de France (CL). We thank the SAAM and its director Josabeth Millereux- Le  
525 Béchennec for providing the archaeological samples presented in this study.

526

## 527 **References**

528

- 529 Amiot, R., Lécuyer, C., Buffetaut, E., Escarguel, G., Fluteau, F., Martineau, F., 2006. Oxygen  
530 isotopes from biogenic apatites suggest widespread endothermy in Cretaceous  
531 dinosaurs. *Earth and Planetary Science Letters* 246, 41–54.  
532 <https://doi.org/10.1016/j.epsl.2006.04.018>
- 533 Baier, J., Lücke, A., Negendank, J.F., Schleser, G.-H., Zolitschka, B., 2004. Diatom and  
534 geochemical evidence of mid-to late Holocene climatic changes at Lake Holzmaar, West-  
535 Eifel (Germany). *Quaternary international* 113, 81–96.
- 536 Barbour, M.M., 2007. Stable oxygen isotope composition of plant tissue: a review. *Functional*  
537 *Plant Biol.* 34, 83–94. <https://doi.org/10.1071/FP06228>
- 538 Barrick, R.E., Showers, W.J., 1995. Oxygen isotope variability in juvenile dinosaurs  
539 (*Hypacrosaurus*): evidence for thermoregulation. *Paleobiology* 21, 552–560.  
540 <https://doi.org/10.1017/S0094837300013531>

541 Barrick, R.E., Showers, W.J., 1994. Thermophysiology of *Tyrannosaurus rex*: Evidence from  
542 Oxygen Isotopes. *Science* 265, 222–224. <https://doi.org/10.1126/science.265.5169.222>

543 Berger, J.-F., Bravard, J.-P., 2012. Le développement économique romain face à la crise  
544 environnementale : le cas de la Gaule narbonnaise. *Des climats et des hommes. La*  
545 *Découverte*, Paris 269–289.

546 Bernard, A., Daux, V., Lécuyer, C., Brugal, J.-P., Genty, D., Wainer, K., Gardien, V., Fourel,  
547 F., Jaubert, J., 2009. Pleistocene seasonal temperature variations recorded in the  $\delta^{18}\text{O}$  of  
548 *Bison priscus* teeth. *Earth and Planetary Science Letters* 283, 133–143.  
549 <https://doi.org/10.1016/j.epsl.2009.04.005>

550 Bernard, A., Lécuyer, C., Vincent, P., Amiot, R., Bardet, N., Buffetaut, E., Cuny, G., Fourel,  
551 F., Martineau, F., Mazin, J.-M., Prieur, A., 2010. Regulation of body temperature by some  
552 mesozoic marine reptiles. *Science* 328, 1379–1382.  
553 <https://doi.org/10.1126/science.1187443>

554 Blake, R.E., O’Neil, J.R., Garcia, G.A., 1997. Oxygen isotope systematics of biologically  
555 mediated reactions of phosphate: I. Microbial degradation of organophosphorus  
556 compounds. *Geochimica et Cosmochimica Acta* 61, 4411–4422.  
557 [https://doi.org/10.1016/S0016-7037\(97\)00272-X](https://doi.org/10.1016/S0016-7037(97)00272-X)

558 Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M.N., Showers, W., Hoffmann, S.,  
559 Lotti-Bond, R., Hajdas, I., Bonani, G., 2001. Persistent Solar Influence on North Atlantic  
560 Climate During the Holocene. *Science* 294, 2130–2136.  
561 <https://doi.org/10.1126/science.1065680>

562 Bookman, R., Enzel, Y., Agnon, A., Stein, M., 2004. Late Holocene lake levels of the Dead  
563 Sea. *GSA Bulletin* 116, 555–571. <https://doi.org/10.1130/B25286.1>

564 Bowen, G.J., 2020. The Online Isotopes in Precipitation Calculator, version 3.1 [WWW  
565 Document]. URL <http://www.waterisotopes.org>



566 Bowen, G.J., Revenaugh, J., 2003. Interpolating the isotopic composition of modern meteoric  
567 precipitation. *Water Resources Research* 39. <https://doi.org/10.1029/2003WR002086>

568 Brettell, R., Montgomery, J., Evans, J., 2012. Brewing and stewing: the effect of culturally  
569 mediated behaviour on the oxygen isotope composition of ingested fluids and the  
570 implications for human provenance studies. *J. Anal. At. Spectrom.* 27, 778–785.  
571 <https://doi.org/10.1039/C2JA10335D>

572 Britton, K., Fuller, B.T., Tütken, T., Mays, S., Richards, M.P., 2015. Oxygen isotope analysis  
573 of human bone phosphate evidences weaning age in archaeological populations. *American*  
574 *Journal of Physical Anthropology* 157, 226–241. <https://doi.org/10.1002/ajpa.22704>

575 Büntgen, U., Tegel, W., Nicolussi, K., McCormick, M., Frank, D., Trouet, V., Kaplan, J.O.,  
576 Herzig, F., Heussner, K.-U., Wanner, H., Luterbacher, J., Esper, J., 2011. 2500 Years of  
577 European Climate Variability and Human Susceptibility. *Science* 331, 578–582.  
578 <https://doi.org/10.1126/science.1197175>

579 Caroli, I., Caldara, M., 2007. Vegetation history of Lago Battaglia (eastern Gargano coast,  
580 Apulia, Italy) during the middle-late Holocene. *Vegetation History and Archaeobotany* 16,  
581 317–327.

582 Chenery, C., Müldner, G., Evans, J., Eckardt, H., Lewis, M., 2010. Strontium and stable  
583 isotope evidence for diet and mobility in Roman Gloucester, UK. *Journal of*  
584 *Archaeological Science* 37, 150–163. <https://doi.org/10.1016/j.jas.2009.09.025>

585 Chesson, L.A., Valenzuela, L.O., O’Grady, S.P., Cerling, T.E., Ehleringer, J.R., 2010. Links  
586 between Purchase Location and Stable Isotope Ratios of Bottled Water, Soda, and Beer in  
587 the United States. *J. Agric. Food Chem.* 58, 7311–7316. <https://doi.org/10.1021/jf1003539>

588 Cisneros, M., Cacho, I., Frigola, J., Canals, M., Martrat, B., Casado, M., Grimalt, J., Pena, L.,  
589 Margaritelli, G., Lirer, F., 2016. Sea surface temperature variability in the central-western  
590 Mediterranean Sea during the last 2700 years: a multi-proxy and multi-record approach.

591 Combourieu Nebout, N., Peyron, O., Dormoy, I., Desprat, S., Beaudouin, C., Kotthoff, U.,  
592 Marret, F., 2009. Rapid climatic variability in the west Mediterranean during the last 25  
593 000 years from high resolution pollen data. *Climate of the Past* 5, 503–521.

594 Craig, H., 1961. Isotopic Variations in Meteoric Waters. *Science* 133, 1702–1703.  
595 <https://doi.org/10.1126/science.133.3465.1702>

596 Craig, H., Gordon, L.I., 1965. Deuterium and oxygen 18 variations in the ocean and the  
597 marine atmosphere.

598 Crowson, R.A., Showers, W.J., Wright, E.K., Hoering, T.C., 1991. Preparation of phosphate  
599 samples for oxygen isotope analysis. *Anal. Chem.* 63, 2397–2400.  
600 <https://doi.org/10.1021/ac00020a038>

601 Csiro, A.P.B., 1983. Solar variability, weather and climate: An update. *Quarterly Journal of*  
602 *the Royal Meteorological Society* 109, 23–55. <https://doi.org/10.1002/qj.49710945903>

603 Dahl-Jensen, D., Mosegaard, K., Gundestrup, N., Clow, G.D., Johnsen, S.J., Hansen, A.W.,  
604 Balling, N., 1998. Past Temperatures Directly from the Greenland Ice Sheet. *Science* 282,  
605 268–271. <https://doi.org/10.1126/science.282.5387.268>

606 D'Angela, D., Longinelli, A., 1990. Oxygen isotopes in living mammal's bone phosphate:  
607 Further results. *Chemical Geology: Isotope Geoscience section* 86, 75–82.  
608 [https://doi.org/10.1016/0168-9622\(90\)90007-Y](https://doi.org/10.1016/0168-9622(90)90007-Y)

609 Daniel Bryant, J., Luz, B., Froelich, P.N., 1994. Oxygen isotopic composition of fossil horse  
610 tooth phosphate as a record of continental paleoclimate. *Palaeogeography,*  
611 *Palaeoclimatology, Palaeoecology, Stable Isotope and Trace-Element Geochemistry of*  
612 *Vertebrate Fossils: Interpreting Ancient Diets and Climates* 107, 303–316.  
613 [https://doi.org/10.1016/0031-0182\(94\)90102-3](https://doi.org/10.1016/0031-0182(94)90102-3)

614 Dansgaard, W., 1964. Stable isotopes in precipitation. *Tellus* 16, 436–468.  
615 <https://doi.org/10.3402/tellusa.v16i4.8993>

616 Darling, G., 2006. Isotopes in water, in: Leng, M.J. (Ed.), *Isotopes in Palaeoenvironmental*  
617 *Research, Developments in Palaeoenvironmental Research*. Springer Netherlands,  
618 Dordrecht, pp. 1–66. [https://doi.org/10.1007/1-4020-2504-1\\_01](https://doi.org/10.1007/1-4020-2504-1_01)

619 Daux, V., Lécuyer, C., Adam, F., Martineau, F., Vimeux, F., 2005. Oxygen isotope  
620 composition of human teeth and the record of climate changes in France (Lorraine) during  
621 the last 1700 years. *Climatic change* 70, 445–464.

622 Daux, V., Lécuyer, C., Héran, M.-A., Amiot, R., Simon, L., Fourel, F., Martineau, F.,  
623 Lynnerup, N., Reychler, H., Escarguel, G., 2008. Oxygen isotope fractionation between  
624 human phosphate and water revisited. *Journal of human evolution* 55, 1138–1147.

625 Dermody, B., De Boer, H.J., Bierkens, M.F.P., Weber, S.L., Wassen, M.J., Dekker, S.C.,  
626 2012. A seesaw in Mediterranean precipitation during the Roman Period linked to  
627 millennial-scale changes in the North Atlantic. *Climate of the Past* 8, 637–651.

628 Desprat, S., Goñi, M.F.S., Loutre, M.-F., 2003. Revealing climatic variability of the last three  
629 millennia in northwestern Iberia using pollen influx data. *Earth and Planetary Science*  
630 *Letters* 213, 63–78.

631 Drake, B.L., 2012. The influence of climatic change on the Late Bronze Age Collapse and the  
632 Greek Dark Ages. *Journal of Archaeological Science* 39, 1862–1870.  
633 <https://doi.org/10.1016/j.jas.2012.01.029>

634 Eberhardt, L.L., 1969. Similarity, allometry and food chains. *Journal of Theoretical Biology*  
635 24, 43–55. [https://doi.org/10.1016/S0022-5193\(69\)80005-6](https://doi.org/10.1016/S0022-5193(69)80005-6)

636 Esper, J., Büntgen, U., Timonen, M., Frank, D.C., 2012. Variability and extremes of northern  
637 Scandinavian summer temperatures over the past two millennia. *Global and Planetary*  
638 *Change* 88, 1–9.

639 Fagan, B., 2008. *The Great Warming: Climate Change and the Rise and Fall of Civilizations*.  
640 Bloomsbury Publishing USA.

641 Ferrio, J.P., Alonso, N., López, J.B., Araus, J.L., Voltas, J., 2006. Carbon isotope composition  
642 of fossil charcoal reveals aridity changes in the NW Mediterranean Basin. *Global Change*  
643 *Biology* 12, 1253–1266.

644 Fourel, F., Martineau, F., Lécuyer, C., Kupka, H.-J., Lange, L., Ojeimi, C., Seed, M., 2011.  
645  $^{18}\text{O}/^{16}\text{O}$  ratio measurements of inorganic and organic materials by elemental analysis–  
646 pyrolysis–isotope ratio mass spectrometry continuous-flow techniques. *Rapid*  
647 *Communications in Mass Spectrometry* 25, 2691–2696. <https://doi.org/10.1002/rcm.5056>

648 Fricke, H.C., O’Neil, J.R., 1999. The correlation between  $^{18}\text{O}/^{16}\text{O}$  ratios of meteoric water and  
649 surface temperature: its use in investigating terrestrial climate change over geologic time.  
650 *Earth and Planetary Science Letters* 170, 181–196. [https://doi.org/10.1016/S0012-](https://doi.org/10.1016/S0012-821X(99)00105-3)  
651 [821X\(99\)00105-3](https://doi.org/10.1016/S0012-821X(99)00105-3)

652 Frisia, S., Borsato, A., Spötl, C., Villa, I.M., Cucchi, F., 2005. Climate variability in the SE  
653 Alps of Italy over the past 17 000 years reconstructed from a stalagmite record. *Boreas* 34,  
654 445–455.

655 Frumkin, A., Magaritz, M., Carmi, I., Zak, I., 1991. The Holocene climatic record of the salt  
656 caves of Mount Sedom Israel. *The Holocene* 1, 191–200.  
657 <https://doi.org/10.1177/095968369100100301>

658 Gat, J.R., 1996. Oxygen and Hydrogen Isotopes in the Hydrologic Cycle. *Annual Review of*  
659 *Earth and Planetary Sciences* 24, 225–262. <https://doi.org/10.1146/annurev.earth.24.1.225>

660 Glimcher, M.J., 2006. Bone: Nature of the Calcium Phosphate Crystals and Cellular,  
661 Structural, and Physical Chemical Mechanisms in Their Formation. *Reviews in*  
662 *Mineralogy and Geochemistry* 64, 223–282. <https://doi.org/10.2138/rmg.2006.64.8>

663 Haigh, J.D., 2007. The Sun and the Earth’s Climate. *Living Rev. Sol. Phys.* 4, 2.  
664 <https://doi.org/10.12942/lrsp-2007-2>

665 Halas, S., Skrzypek, G., Meier- Augenstein, W., Pelc, A., Kemp, H.F., 2011. Inter-laboratory  
666 calibration of new silver orthophosphate comparison materials for the stable oxygen  
667 isotope analysis of phosphates. *Rapid Communications in Mass Spectrometry* 25, 579–  
668 584. <https://doi.org/10.1002/rcm.4892>

669 Heim, C., Nowaczyk, N.R., Negendank, J.F., Leroy, S.A., Ben-Avraham, Z., 1997. Near East  
670 desertification: evidence from the Dead Sea. *Naturwissenschaften* 84, 398–401.

671 Holzhauser, H., Magny, M., Zumbühl, H.J., 2005. Glacier and lake-level variations in west-  
672 central Europe over the last 3500 years. *The Holocene* 15, 789–801.

673 Hut, G., 1987. Consultants' group meeting on stable isotope reference samples for  
674 geochemical and hydrological investigations.

675 IAEA/WMO, 2015. Global Network of Isotopes in Precipitation. The GNIP Database.  
676 [WWW Document]. URL <https://nucleus.iaea.org/wiser>

677 Issar, A.S., Yakir, D., 1997. Isotopes from Wood Buried in the Roman Siege Ramp of  
678 Masada: The Roman Period's Colder Climate. *The Biblical Archaeologist* 60, 101–106.  
679 <https://doi.org/10.2307/3210599>

680 Jay, M., Richards, M.P., 2006. Diet in the Iron Age cemetery population at Wetwang Slack,  
681 East Yorkshire, UK: carbon and nitrogen stable isotope evidence. *Journal of*  
682 *Archaeological Science* 33, 653–662. <https://doi.org/10.1016/j.jas.2005.09.020>

683 Jiang, H., Eiríksson, J., Schulz, M., Knudsen, K.-L., Seidenkrantz, M.-S., 2005. Evidence for  
684 solar forcing of sea-surface temperature on the North Icelandic Shelf during the late  
685 Holocene. *Geology* 33, 73–76. <https://doi.org/10.1130/G21130.1>

686 Kaniewski, D., Paulissen, E., Campo, E.V., Weiss, H., Otto, T., Bretschneider, J., Lerberghe,  
687 K.V., 2010. Late second–early first millennium BC abrupt climate changes in coastal Syria  
688 and their possible significance for the history of the Eastern Mediterranean. *Quaternary*  
689 *Research* 74, 207–215. <https://doi.org/10.1016/j.yqres.2010.07.010>

690 Kohn, M.J., 1996. Predicting animal  $\delta^{18}\text{O}$ : Accounting for diet and physiological adaptation.  
691 *Geochimica et Cosmochimica Acta* 60, 4811–4829. <https://doi.org/10.1016/S0016->  
692 7037(96)00240-2

693 Kottek, M., Grieser, J., Beck, C., Rudolf, B., Rubel, F., 2006. World map of the Köppen-  
694 Geiger climate classification updated. *Meteorologische Zeitschrift* 15, 259–263.

695 Lamb, A.L., Evans, J.E., Buckley, R., Appleby, J., 2014. Multi-isotope analysis demonstrates  
696 significant lifestyle changes in King Richard III. *Journal of Archaeological Science* 50,  
697 559–565. <https://doi.org/10.1016/j.jas.2014.06.021>

698 Lamb, H.H., 1995. *Climate, History and the Modern World*. Psychology Press.

699 Le Béchenec, Y., Frère, S., Corsiez, A., Rolland, J., Sauvageot, P., 2016. Rapport final  
700 d’opération : Thézy-Glimont, Les Vergnes, (Somme). Novembre 2012 - Mai 2013. Service  
701 Archéologie Préventive d’Amiens Métropole.

702 Lécuyer, C., 2013. *Water on Earth*, 1st ed. John Wiley & Sons, Ltd.  
703 <https://doi.org/10.1002/9781118574928>

704 Lécuyer, C., Atrops, F., Amiot, R., Angst, D., Daux, V., Flandrois, J.-P., Fourel, F., Rey, K.,  
705 Royer, A., Seris, M., Touzeau, A., Rousseau, D., 2018. Tsunami sedimentary deposits of  
706 Crete records climate during the ‘Minoan Warming Period’ ( $\approx 3350$  yr BP). *The Holocene*  
707 28, 914–929. <https://doi.org/10.1177/0959683617752840>

708 Lécuyer, C., Fourel, F., Martineau, F., Amiot, R., Bernard, A., Daux, V., Escarguel, G.,  
709 Morrison, J., 2007. High-precision determination of  $^{18}\text{O}/^{16}\text{O}$  ratios of silver phosphate by  
710 EA-pyrolysis-IRMS continuous flow technique. *Journal of Mass Spectrometry* 42, 36–41.  
711 <https://doi.org/10.1002/jms.1130>

712 Lécuyer, C., Grandjean, P., O’Neil, J.R., Cappetta, H., Martineau, F., 1993. Thermal  
713 excursions in the ocean at the Cretaceous—Tertiary boundary (northern Morocco):  $\delta^{18}\text{O}$

714 record of phosphatic fish debris. *Palaeogeography, Palaeoclimatology, Palaeoecology* 105,  
715 235–243. [https://doi.org/10.1016/0031-0182\(93\)90085-W](https://doi.org/10.1016/0031-0182(93)90085-W)

716 Lécuyer, C., Grandjean, P., Sheppard, S.M.F., 1999. Oxygen isotope exchange between  
717 dissolved phosphate and water at temperatures  $\leq 135^{\circ}\text{C}$ : inorganic versus biological  
718 fractionations. *Geochimica et Cosmochimica Acta* 63, 855–862.  
719 [https://doi.org/10.1016/S0016-7037\(99\)00096-4](https://doi.org/10.1016/S0016-7037(99)00096-4)

720 LeGeros, R., 1981. Apatites in biological systems. *Progress in Crystal Growth and*  
721 *Characterization* 4, 1–45. [https://doi.org/10.1016/0146-3535\(81\)90046-0](https://doi.org/10.1016/0146-3535(81)90046-0)

722 Lightfoot, E., O’Connell, T.C., 2016. On the Use of Biomineral Oxygen Isotope Data to  
723 Identify Human Migrants in the Archaeological Record: Intra-Sample Variation, Statistical  
724 Methods and Geographical Considerations. *PLoS One* 11.  
725 <https://doi.org/10.1371/journal.pone.0153850>

726 Lirer, F., Sprovieri, M., Vallefucio, M., Ferraro, L., Pelosi, N., Giordano, L., Capotondi, L.,  
727 2014. Planktonic foraminifera as bio-indicators for monitoring the climatic changes that  
728 have occurred over the past 2000 years in the southeastern Tyrrhenian Sea. *Integrative*  
729 *Zoology* 9, 542–554. <https://doi.org/10.1111/1749-4877.12083>

730 Lockwood, M., 2012. Solar Influence on Global and Regional Climates. *Surv Geophys* 33,  
731 503–534. <https://doi.org/10.1007/s10712-012-9181-3>

732 Longin, R., 1971. New Method of Collagen Extraction for Radiocarbon Dating. *Nature* 230,  
733 241–242. <https://doi.org/10.1038/230241a0>

734 Longinelli, A., 1984. Oxygen isotopes in mammal bone phosphate: A new tool for  
735 paleohydrological and paleoclimatological research? *Geochimica et Cosmochimica Acta*  
736 48, 385–390. [https://doi.org/10.1016/0016-7037\(84\)90259-X](https://doi.org/10.1016/0016-7037(84)90259-X)

737 Luterbacher, J., Werner, J.P., Smerdon, J.E., Fernández-Donado, L., González-Rouco, F.J.,  
738 Barriopedro, D., Ljungqvist, F.C., Büntgen, U., Zorita, E., Wagner, S., 2016. European  
739 summer temperatures since Roman times. *Environmental research letters* 11, 024001.

740 Luz, B., Kolodny, Y., 1985. Oxygen isotope variations in phosphate of biogenic apatites, IV.  
741 Mammal teeth and bones. *Earth and Planetary Science Letters* 75, 29–36.  
742 [https://doi.org/10.1016/0012-821X\(85\)90047-0](https://doi.org/10.1016/0012-821X(85)90047-0)

743 Luz, B., Kolodny, Y., Horowitz, M., 1984. Fractionation of oxygen isotopes between  
744 mammalian bone-phosphate and environmental drinking water. *Geochimica et*  
745 *Cosmochimica Acta* 48, 1689–1693. [https://doi.org/10.1016/0016-7037\(84\)90338-7](https://doi.org/10.1016/0016-7037(84)90338-7)

746 Magny, M., de Beaulieu, J.-L., Drescher-Schneider, R., Vanni re, B., Walter-Simonnet, A.-  
747 V., Miras, Y., Millet, L., Bossuet, G., Peyron, O., Brugiapaglia, E., Leroux, A., 2007.  
748 Holocene climate changes in the central Mediterranean as recorded by lake-level  
749 fluctuations at Lake Accesa (Tuscany, Italy). *Quaternary Science Reviews* 26, 1736–1758.  
750 <https://doi.org/10.1016/j.quascirev.2007.04.014>

751 Mangini, A., Sp t l, C., Verdes, P., 2005. Reconstruction of temperature in the Central Alps  
752 during the past 2000 yr from a  $\delta^{18}\text{O}$  stalagmite record. *Earth and Planetary Science Letters*  
753 235, 741–751. <https://doi.org/10.1016/j.epsl.2005.05.010>

754 Margaritelli, G., Vallefuc o, M., Di Rita, F., Capotondi, L., Bellucci, L.G., Insinga, D.D.,  
755 Petrosino, P., Bonomo, S., Cacho, I., Cascella, A., Ferraro, L., Florindo, F., Lubritto, C.,  
756 Lurcock, P.C., Magri, D., Pelosi, N., Rettori, R., Lirer, F., 2016. Marine response to  
757 climate changes during the last five millennia in the central Mediterranean Sea. *Global and*  
758 *Planetary Change* 142, 53–72. <https://doi.org/10.1016/j.gloplacha.2016.04.007>

759 Marquer, L., Pomel, S., Abichou, A., Schulz, E., Kaniewski, D., Campo, E.V., 2008. Late  
760 Holocene high resolution palaeoclimatic reconstruction inferred from Sebkh  Mhabeul,



761 southeast Tunisia. Quaternary Research 70, 240–250.  
762 <https://doi.org/10.1016/j.yqres.2008.06.002>

763 Martin, C., Maureille, B., Amiot, R., Touzeau, A., Royer, A., Fourel, F., Panczer, G.,  
764 Flandrois, J.-P., Lécuyer, C., 2017. Record of Nile seasonality in Nubian neonates.  
765 Isotopes in environmental and health studies 53, 223–242.

766 McCormick, M., Büntgen, U., Cane, M.A., Cook, E.R., Harper, K., Huybers, P., Litt, T.,  
767 Manning, S.W., Mayewski, P.A., More, A.F.M., Nicolussi, K., Tegel, W., 2012. Climate  
768 Change during and after the Roman Empire: Reconstructing the Past from Scientific and  
769 Historical Evidence. The Journal of Interdisciplinary History 43, 169–220.  
770 [https://doi.org/10.1162/JINH\\_a\\_00379](https://doi.org/10.1162/JINH_a_00379)

771 McDermott, F., Matthey, D.P., Hawkesworth, C., 2001. Centennial-Scale Holocene Climate  
772 Variability Revealed by a High-Resolution Speleothem  $\delta^{18}\text{O}$  Record from SW Ireland.  
773 Science 294, 1328–1331. <https://doi.org/10.1126/science.1063678>

774 Migowski, C., Stein, M., Prasad, S., Negendank, J.F.W., Agnon, A., 2006. Holocene Climate  
775 Variability and Cultural Evolution in the Near East from the Dead Sea Sedimentary  
776 Record. Quaternary Research 66, 421–431. <https://doi.org/10.1016/j.yqres.2006.06.010>

777 Moreau, C., Caffy, I., Comby, C., Delqué-Količ, E., Dumoulin, J.-P., Hain, S., Quiles, A.,  
778 Setti, V., Souprayen, C., Thellier, B., Vincent, J., 2013. Research and Development of the  
779 Artemis  $^{14}\text{C}$  AMS Facility: Status Report. Radiocarbon 55, 331–337.  
780 <https://doi.org/10.1017/S0033822200057441>

781 Négrel, Ph., Petelet-Giraud, E., 2005. Strontium isotopes as tracers of groundwater-induced  
782 floods: the Somme case study (France). Journal of Hydrology 305, 99–119.  
783 <https://doi.org/10.1016/j.jhydrol.2004.08.031>

784 Nesje, A., Lie, Ø., Dahl, S.O., 2000. Is the North Atlantic Oscillation reflected in  
785 Scandinavian glacier mass balance records? *Journal of Quaternary Science* 15, 587–601.  
786 [https://doi.org/10.1002/1099-1417\(200009\)15:6<587::AID-JQS533>3.0.CO;2-2](https://doi.org/10.1002/1099-1417(200009)15:6<587::AID-JQS533>3.0.CO;2-2)

787 Neukom, R., Steiger, N., Gómez-Navarro, J.J., Wang, J., Werner, J.P., 2019. No evidence for  
788 globally coherent warm and cold periods over the preindustrial Common Era. *Nature* 571,  
789 550–554. <https://doi.org/10.1038/s41586-019-1401-2>

790 Oldenborgh, G.J. van, Laat, A.T.J. de, Luterbacher, J., Ingram, W.J., Osborn, T.J., 2013.  
791 Claim of solar influence is on thin ice: are 11-year cycle solar minima associated with  
792 severe winters in Europe? *Environ. Res. Lett.* 8, 024014. [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/8/2/024014)  
793 [9326/8/2/024014](https://doi.org/10.1088/1748-9326/8/2/024014)

794 Piva, A., Asioli, A., Trincardi, F., Schneider, R.R., Vigliotti, L., 2008. Late-Holocene climate  
795 variability in the Adriatic Sea (Central Mediterranean). *The Holocene* 18, 153–167.  
796 <https://doi.org/10.1177/0959683607085606>

797 Prowse, T., Schwarcz, H.P., Saunders, S., Macchiarelli, R., Bondioli, L., 2004. Isotopic  
798 paleodiet studies of skeletons from the Imperial Roman-age cemetery of Isola Sacra,  
799 Rome, Italy. *Journal of Archaeological Science* 31, 259–272.  
800 <https://doi.org/10.1016/j.jas.2003.08.008>

801 Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck, C.E.,  
802 Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Haflidason, H.,  
803 Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser,  
804 K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M.,  
805 Southon, J.R., Staff, R.A., Turney, C.S.M., Plicht, J. van der, 2013. IntCal13 and Marine13  
806 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP. *Radiocarbon* 55, 1869–1887.  
807 [https://doi.org/10.2458/azu\\_js\\_rc.55.16947](https://doi.org/10.2458/azu_js_rc.55.16947)

808 Richardin, P., Porcier, S., Ikram, S., Louarn, G., Berthet, D., 2017. Cats, Crocodiles, Cattle,  
809 and More: Initial Steps Toward Establishing a Chronology of Ancient Egyptian Animal  
810 Mummies. *Radiocarbon* 59, 595–607. <https://doi.org/10.1017/RDC.2016.102>

811 Roberts, N., Reed, J.M., Leng, M.J., Kuzucuoğlu, C., Fontugne, M., Bertaux, J., Woldring,  
812 H., Bottema, S., Black, S., Hunt, E., Karabiyoğlu, M., 2001. The tempo of Holocene  
813 climatic change in the eastern Mediterranean region: new high-resolution crater-lake  
814 sediment data from central Turkey. *The Holocene* 11, 721–736.  
815 <https://doi.org/10.1191/09596830195744>

816 Rozanski, K., Araguás-Araguás, L., Gonfiantini, R., 1993. Isotopic patterns in modern global  
817 precipitation. *Climate change in continental isotopic records* 78, 1–36.

818 Rozanski, K., Araguás-Araguás, L., Gonfiantini, R., 1992. Relation Between Long-Term  
819 Trends of Oxygen-18 Isotope Composition of Precipitation and Climate. *Science* 258,  
820 981–985. <https://doi.org/10.1126/science.258.5084.981>

821 Schilman, B., Bar-Matthews, M., Almogi-Labin, A., Luz, B., 2001. Global climate instability  
822 reflected by Eastern Mediterranean marine records during the late Holocene.  
823 *Palaeogeography, Palaeoclimatology, Palaeoecology* 176, 157–176.  
824 [https://doi.org/10.1016/S0031-0182\(01\)00336-4](https://doi.org/10.1016/S0031-0182(01)00336-4)

825 Seidenkrantz, M.-S., Aagaard-Sørensen, S., Sulsbrück, H., Kuijpers, A., Jensen, K.G.,  
826 Kunzendorf, H., 2007. Hydrography and climate of the last 4400 years in a SW Greenland  
827 fjord: implications for Labrador Sea palaeoceanography. *The Holocene* 17, 387–401.  
828 <https://doi.org/10.1177/0959683607075840>

829 Séon, N., Amiot, R., Martin, J.E., Young, M.T., Middleton, H., Fourel, F., Picot, L., Valentin,  
830 X., Lécuyer, C., 2020. Thermophysiology of Jurassic marine crocodylomorphs inferred  
831 from the oxygen isotope composition of their tooth apatite. *Philosophical Transactions of*

832 the Royal Society B: Biological Sciences 375, 20190139.  
833 <https://doi.org/10.1098/rstb.2019.0139>

834 Shimamoto, H., Komiya, S., 2000. The turnover of body water as an indicator of health. *J*  
835 *Physiol Anthropol Appl Human Sci* 19, 207–212. <https://doi.org/10.2114/jpa.19.207>

836 Smith, A.C., Wynn, P.M., Barker, P.A., Leng, M.J., Noble, S.R., Tych, W., 2016. North  
837 Atlantic forcing of moisture delivery to Europe throughout the Holocene. *Scientific*  
838 *Reports* 6, 1–7. <https://doi.org/10.1038/srep24745>

839 Taricco, C., Ghil, M., Alessio, S.M., Vivaldo, G., 2009. Two millennia of climate variability  
840 in the Central Mediterranean 5, 171–181. <https://doi.org/10.5194/cp-5-171-2009>

841 Tinner, W., Lotter, A.F., Ammann, B., Conedera, M., Hubschmid, P., van Leeuwen, J.F.N.,  
842 Wehrl, M., 2003. Climatic change and contemporaneous land-use phases north and south  
843 of the Alps 2300 BC to 800 AD. *Quaternary Science Reviews* 22, 1447–1460.  
844 [https://doi.org/10.1016/S0277-3791\(03\)00083-0](https://doi.org/10.1016/S0277-3791(03)00083-0)

845 Touzeau, A., Blichert-Toft, J., Amiot, R., Fourel, F., Martineau, F., Cockitt, J., Hall, K.,  
846 Flandrois, J.-P., Lécuyer, C., 2013. Egyptian mummies record increasing aridity in the Nile  
847 valley from 5500 to 1500 yr before present. *Earth and Planetary Science Letters* 375, 92–  
848 100.

849 Tütken, T., Vennemann, T.W., 2011. Fossil bones and teeth: Preservation or alteration of  
850 biogenic compositions? *Palaeogeography, Palaeoclimatology, Palaeoecology* 1–2, 1–8.  
851 <https://doi.org/10.1016/j.palaeo.2011.06.020>

852 von Grafenstein, U., Erlenkeuser, H., Müller, J., Trumborn, P., Alefs, J., 1996. A 200-year  
853 mid-European air temperature record preserved in lake sediments: An extension of the  
854  $\delta^{18}\text{O}_{\text{p}}$ -air temperature relation into the past. *Geochimica et Cosmochimica Acta* 60, 4025–  
855 4036. [https://doi.org/10.1016/S0016-7037\(96\)00230-X](https://doi.org/10.1016/S0016-7037(96)00230-X)

- 856 Wang, T., Surge, D., Mithen, S., 2012. Seasonal temperature variability of the Neoglacial  
857 (3300–2500BP) and Roman Warm Period (2500–1600BP) reconstructed from oxygen  
858 isotope ratios of limpet shells (*Patella vulgata*), Northwest Scotland. *Palaeogeography,*  
859 *Palaeoclimatology,* *Palaeoecology* 317–318, 104–113.  
860 <https://doi.org/10.1016/j.palaeo.2011.12.016>
- 861 Wick, L., Lemcke, G., Sturm, M., 2003. Evidence of Lateglacial and Holocene climatic  
862 change and human impact in eastern Anatolia: high-resolution pollen, charcoal, isotopic  
863 and geochemical records from the laminated sediments of Lake Van, Turkey. *The*  
864 *Holocene* 13, 665–675. <https://doi.org/10.1191/0959683603hl653rp>
- 865 Wopenka, B., Pasteris, J.D., 2005. A mineralogical perspective on the apatite in bone.  
866 *Materials Science and Engineering: C, NATO Advanced Study Institute (ASI on Learning*  
867 *from Nature How to design New Implantable Biomaterials: From Biomineralization*  
868 *Fundamentals to Biomimetic Materials and Processing Routes)* 25, 131–143.  
869 <https://doi.org/10.1016/j.msec.2005.01.008>
- 870 Zanchetta, G., Drysdale, R.N., Hellstrom, J.C., Fallick, A.E., Isola, I., Gagan, M.K., Pareschi,  
871 M.T., 2007. Enhanced rainfall in the Western Mediterranean during deposition of sapropel  
872 S1: stalagmite evidence from Corchia cave (Central Italy). *Quaternary Science Reviews*  
873 26, 279–286. <https://doi.org/10.1016/j.quascirev.2006.12.003>

874

## 875 **Tables**

876

877 Table 1: Grave, nature, identification and corresponding dates (at 95.4% certainty) inferred  
878 from <sup>14</sup>C dating of the samples from Thézy-Glimont. The archaeological data come from Le  
879 Béchenec et al. (2016).

880 \* corresponds to the three individuals for which various bones were sampled, see Figure 2.

881 §: If there are two or three dates from the same sample, the radiocarbon dates have been  
882 combined before calibration. Such a combination is checked for internal consistency by a  
883 chi-squared test which is performed automatically by the OxCal™ program.

884

## 885 **Figures**

886

887 Figure 1: (a) Geographic map of France with the location of the Gallic site of Thézy-Glimont,  
888 Picardie, France. PM = Prime Meridian. The datum used for latitude and longitude is WGS84.  
889 Picture was taken from Google Earth™. (b) General plan of the archaeological site. 1: Area  
890 where the excavated graves are located. 2: Intermediate area. 3: Enclosure with iron furniture  
891 and weapons. In gray is represented a trench from World War 1 (1918). The background  
892 image was taken from Google Maps™.

893

894 Figure 2: Sampling points and intra-skeletal  $\delta^{18}\text{O}_p$  (‰, V-SMOW) variations represented as  
895 local deviation from the average  $\delta^{18}\text{O}_p$  value of bones coming from TG-01, TG-02 and TG-03  
896 individuals.

897

898 Figure 3:  $\delta^{18}\text{O}_p$  (‰, V-SMOW) values of bones (blue squares) and teeth (red triangles) from  
899 humans and animals of Thézy-Glimont. Bones and teeth which are aligned vertically  
900 correspond to a same individual. Humans, bovids and horses are represented by increasing  
901 sample number from left to right.

902

903 Figure 4: Comparison of oxygen isotope fractionation equations between apatite phosphate  
904 ( $\delta^{18}\text{O}_p$ , ‰, V-SMOW) and drinking water ( $\delta^{18}\text{O}_w$ , ‰, V-SMOW) proposed by Daux et al.


905 (2008) for humans, Longinelli (1984) for pigs, D'Angela and Longinelli (1990) for bovids  
906 and Bryant et al. (1994) for horses.

907


908 Figure 5:  $\delta^{18}\text{O}_w$  values (‰, V-SMOW) of drinking water inferred from the bones of humans  
909 and animals from Thézy-Glimont. Humans and bovids are represented by increasing sample  
910 number from left to right. Error bars represent the total uncertainties. The red dashed lines  
911 represent the modern  $\delta^{18}\text{O}_w$  values measured in the Avre river by Négrel and Petelet-Giraud  
912 (2005) in April (A) and October (O) 2001.

913

914 Figure 6: Reconstructed Mean Annual Temperatures (MAT) inferred from the oxygen isotope  
915 composition of drinking water, using the oxygen isotope fractionation equation of von  
916 Grafenstein et al. (1996) (blue squares) and Lécuyer (2013) (red triangles). Humans and  
917 bovids are represented by increasing sample number from left to right. The horizontal dashed  
918 line corresponds to the modern yearly MAT in Picardie, France, averaged throughout the  
919 years 1980 to 2010 (data from Météo France<sup>TM</sup>).




(a)




(b)







TG-01  
 Mean  $\delta^{18}\text{O}_p$  (V-SMOW): 16.6‰  
 Standard Deviation: 0.2‰





TG-02  
 Mean  $\delta^{18}\text{O}_p$  (V-SMOW): 17.6‰  
 Standard Deviation: 0.2‰




TG-03  
 Mean  $\delta^{18}\text{O}_p$  (V-SMOW): 16.8‰  
 Standard Deviation: 0.3‰









Grave	Individual	Type	Sex	Age (years)	Sample type	<sup>14</sup> C dates (BP)	Calibrated dates - 2σ (95.4%) <sup>§</sup>
TGV 1135	TG-01	<i>Homo sapiens</i>	Male	< 30	*	2160 ± 30 2190 ± 30	358 cal BCE (56.7%) 280 cal BCE 258 cal BCE (2.6%) 243 cal BCE 236 cal BCE (36.1%) 170 cal BCE
					1st premolar (PM1)	-	-
					1st molar (M1)	-	-
					2 <sup>nd</sup> molar (M2)	-	-
TG-CH1	<i>Equus caballus</i>	Male	3	Left phalanx	2130 ± 30	350 cal BCE (10.5%) 308 cal BCE 210 cal BCE (84.9%) 52 cal BCE	
				3rd molar (M3)	2150 ± 30 2180 ± 30 2190 ± 30	356 cal BCE (59.0%) 286 cal BCE 234 cal BCE (36.4%) 170 cal BCE	
TG-B1	<i>Bos taurus</i>	Female	4	Right phalanx	-	-	
TG-B2	<i>Bos taurus</i>	Male	young	Left talus	-	-	
TG-B3	<i>Bos taurus</i>	Male	2	Left phalanx	-	-	
TG-B4	<i>Bos taurus</i>	Male	4	Left phalanx	-	-	
TG-CO1	<i>Sus scrofa</i>	Male	-	3rd molar (M3)	2170 ± 30 2205 ± 30	359 cal BCE (59.2%) 274 cal BCE 261 cal BCE (36.2%) 184 cal BCE	
				Right Radius	-	-	
				Rib	-	-	
				Vertebra	-	-	

<b>TGV 1249</b>	TG-02	<i>Homo sapiens</i>	Male	15	*	2200 ± 30 2170 ± 30	358 cal BCE (59.2%) 276 cal BCE 260 cal BCE (36.2%) 180 cal BCE
					1st premolar (PM1)	-	-
					1st molar (M1)	-	-
					3rd molar (M3)	-	-
<b>TGV 1450</b>	TG-03	<i>Homo sapiens</i>	Male	Adult	*	2185 ± 30	361 cal BCE (95.4%) 172 cal BCE
					1st premolar (PM1)	-	-
					2nd molar (M2)	-	-
					3rd molar (M3)	-	-
<b>TGV 1219</b>	TG-04	<i>Homo sapiens</i>	Male	Adult	Right femur	2195 ± 30 2175 ± 30	358 cal BCE (59.0%) 276 cal BCE 260 cal BCE (36.4%) 180 cal BCE
<b>TGV 1138</b>	TG-05	<i>Homo sapiens</i>	Male	30 to 59	Right femur	2180 ± 30 2145 ± 30	356 cal BCE (46.7%) 284 cal BCE 256 cal BCE (0.5%) 250 cal BCE 235 cal BCE (46.0%) 156 cal BCE 134 cal BCE (2.1%) 116 cal BCE
	TG-CH2	<i>Equus caballus</i>	Male	5	Molar	2120 ± 30 2135 ± 30	344 cal BCE (4.3%) 322 cal BCE 206 cal BCE (87.9%) 88 cal BCE 76 cal BCE (3.2%) 58 cal BCE
<b>TGV 1140</b>	TG-06	<i>Homo sapiens</i>	Male	Adult	Right femur	2170 ± 30 2145 ± 30	355 cal BCE (40.5%) 289 cal BCE 232 cal BCE (51.0%) 151 cal BCE 136 cal BCE (3.9%) 114 cal BCE
<b>TGV1206</b>	TG-07	<i>Homo sapiens</i>	Male	Adult	Right femur	2200 ± 30 2225 ± 30	364 cal BCE (95.4%) 203 cal BCE
<b>TGV 1246</b>	TG-08	<i>Homo sapiens</i>	Male	Adult	Tibia	-	-

<b>TGV 1222</b>	TG-B5	<i>Bos taurus</i>	Male	2 to 3	3rd molar	2135 ± 30 2160 ± 30 2115 ± 30	346 cal BCE (8.2%) 320 cal BCE 206 cal BCE (87.2%) 102 cal BCE
-----------------	-------	-------------------	------	--------	-----------	-------------------------------------	---