



HAL
open science

Evidence for centennial-scale Mid-Holocene episodes of hypolimnetic anoxia in a high-altitude lake system from central Tian Shan (Kyrgyzstan)

Philippe Sorrel, Kevin Jacq, Antonin van Exem, Gilles Escarguel, Benjamin Dietre, Maxime Debret, Suzanne McGowan, Ducept Jules, Gauthier Emilie, Hedi Oberhänsli

► To cite this version:

Philippe Sorrel, Kevin Jacq, Antonin van Exem, Gilles Escarguel, Benjamin Dietre, et al.. Evidence for centennial-scale Mid-Holocene episodes of hypolimnetic anoxia in a high-altitude lake system from central Tian Shan (Kyrgyzstan). *Quaternary Science Reviews*, 2021, 252, pp.106748. 10.1016/j.quascirev.2020.106748 . hal-03079515

HAL Id: hal-03079515

<https://univ-lyon1.hal.science/hal-03079515>

Submitted on 2 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Evidence for centennial-scale Mid-Holocene episodes of hypolimnetic anoxia in a high-**
2 **altitude lake system from Central Tian Shan (Kyrgyzstan)**

3
4 Philippe Sorrel^a, Kévin Jacq^b, Antonin Van Exem^b, Gilles Escarguel^c, Benjamin Dietre^d,
5 Maxime Debret^b, Suzanne McGowan^e, Jules Ducept^a, Emilie Gauthier^f and Hedi Oberhänsli^g

6
7 ^a Univ Lyon, Univ Lyon 1, ENSL, CNRS, LGL-TPE, F-69622, Villeurbanne, France

8 ^b Normandie Univ, UNIROUEN, UNICAEN, CNRS, M2C, 76000 Rouen, France

9 ^c Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-
10 69622, Villeurbanne, France

11 ^d Institut für Botanik, Universität Innsbruck, Sternwartestrasse 15, 6020 Innsbruck, Austria

12 ^e School of Geography, University of Nottingham, University Park, Nottingham, UK

13 ^f UMR 6249, Laboratoire Chrono-environnement, 16 route de Gray, 25030 Besançon cedex,
14 France

15 ^g Museum für Naturkunde, Leibnitz-Institute Berlin (Mineralogy), Invalidenstrasse 43, 10115
16 Berlin, Germany

17

18 *Corresponding author: P. Sorrel; philippe.sorrel@univ-lyon1.fr

19

20 **Keywords**

21 Hyperspectral imaging, Paleolimnology, Paleoclimate, Lake stratification, Hypolimnetic
22 anoxia, Chemocline, Central Asia, Lake Son Kol, Holocene

23

24

25

26 **Abstract**

27 Few sedimentary archives of lake meromixis are available in palaeolimnological records,
28 because long-term observations are limited in time and indisputable sediment proxies of
29 hypolimnetic anoxia are still scarce. Here we use visible and near infrared (VNIR), and short-
30 wave infra-red (SWIR) hyperspectral imaging combined with geochemical analyses to
31 reconstruct lake stratification history, redox status and mixing conditions in the water column
32 of Lake Son Kol (Kyrgyzstan) in Central Tian Shan during the last 8500 years. In particular,
33 the detection of Bacteriopheophytin *a* (Bphe *a*), a pigment produced by anoxygenic
34 phototrophic bacteria at the chemocline of meromictic lakes, emphasizes episodes of multi-
35 decadal to centennial hypolimnetic anoxia in Lake Son Kol. Phases of hypolimnetic anoxia
36 are inferred from the deposition of dark organic sediments within a stratified lake system,
37 which occurred during periods of increased snowmelt (and solid winter precipitation) and
38 warmer spring/summer temperatures that promoted floods and the **export** of terrestrial
39 material from the catchment. Prolonged euxinic conditions in bottom waters (involving the
40 development of a stable chemocline) are reported around 8500, 8400, 8200–7800, 7700–
41 7500, 7300–7000, 6500–6100, 6000–5700 and 5500–5250 cal. yr BP. At ca. 5250 cal. yr BP,
42 the chemocline abruptly vanished as Lake Son Kol tipped into a regime with predominantly
43 cooler and well-mixed conditions (predominance of oxygenic phototrophs), coeval with
44 higher lake levels. The disappearance of hypolimnetic anoxia in Lake Son Kol coincides with
45 strengthened wind conditions that imply enhanced lake overturning and upward mixing of
46 nutrients in the water column. This study reveals the strong potential of hyperspectral
47 imaging, in combination with more classical palaeolimnological approaches, to reconstruct
48 the lake trophic and mixing history and explore the controlling mechanisms at work on
49 decadal to centennial timescales. Our results outline how abrupt ecosystem changes may
50 occur even in the absence of anthropogenic climate change.

51

52 **1. Introduction**

53 Anthropogenic global warming is changing high-altitude ecosystems and environmental
54 dynamics (Solomon et al., 2007; IPCC 2013). Regional model outputs for Central Asia
55 predict that seasonal surface temperatures in the region will increase from 3 °C up to 11.4 °C
56 on average, coeval with a decline in the amount of precipitation (Ozturk et al., 2012),
57 although uncertainties in hydrological/rainfall projections are existing. High elevation regions
58 of Central Asia (e.g., Tian Shan and Pamirs mountains, Tibetan Plateau) are particularly
59 vulnerable to warmer climate conditions (e.g., Christensen et al., 2007). In high-altitude areas,
60 a consequence of global warming may be shorter and wetter (and milder) winters along with
61 longer and drier summers (Christensen et al., 2007; IPCC, 2013). Increases in solid winter
62 precipitation, spring/summer temperatures, and (summer) rainstorms are likely to increase
63 flooding of snowmelt-fed lake basins. Warmer temperatures are also expected to lead to
64 eutrophication of lakes, depletion of oxygen in deep waters (Jenny et al., 2013), and longer
65 and stronger temperature stratification in summer (e.g., Adrian et al., 1995; Livingstone,
66 2003; Foley et al., 2012). Incomplete lake overturning may result in the development of a
67 chemocline (Davison, 1993) with an anoxic hypolimnion underneath (Butz et al., 2016),
68 which may last for prolonged periods (meromixis). Meromixis implies persistent lake
69 stratification over multiple years caused by chemical differentiation of the bottom waters, (cf.
70 seasonal thermal stratification which is common in deep temperate lakes and may occur
71 above the chemocline in a meromictic lake). Investigations of long-term trends in oxygen
72 depletion in lakes, and their relationship with climatic changes on timescales beyond the
73 instrumental period, are still scarce in high Central Asia.

74 Located at the intercept between the influences of the Westerlies and the Siberian Anticyclone
75 (Chen et al., 2008; 2010), the Tian Shan holds key information about past climate variability

76 and particularly temperature and hydrological changes in western Central Asia (e.g.,
77 Rousseau et al., 2020). A number of valuable and high-quality lake sediment studies have
78 examined Holocene climate development in the Tian Shan and Pamir mountains (e.g.,
79 Aichner et al., 2015; Beer et al., 2007; Li et al., 2011; Mischke et al., 2010; Chen et al., 2016;
80 Heinecke et al., 2016, 2017, 2018; Rousseau et al., 2020), including several at Lake Son Kol
81 (or Son Kul) in Kyrgyzstan (Huang et al., 2014; Mathis et al., 2014; Pacton et al., 2015;
82 Lauterbach et al., 2014; Schwarz et al., 2017, Wolff et al., 2017; Laug et al., 2020). The focus
83 of these works has been on the Mid- to Late Holocene, with however limited
84 palaeoenvironmental reconstructions between 8500 and 6000 cal. yr BP. Thus there are
85 virtually no high-resolution palaeolimnological records of long-term lake mixing conditions
86 from this region yet, which could provide information on the complex links between
87 environmental, climatic and anthropogenic changes in high-altitude settings.

88 Here we combine hyperspectral imaging (HSI) and X-Ray fluorescence (XRF) analyses of a
89 new core from Lake Son Kol (SK06) to obtain an unprecedented record of sedimentary
90 pigment changes at ultra-high sampling resolution (μm -scale; sub-annual resolution).

91 Building upon the work of Huang et al. (2014), who inferred that most of the annual
92 precipitation switched from the summer to the cold (winter) season between the Mid- and
93 Late Holocene, we used sedimentary pigments and geochemical data to (i) decipher lake
94 trophic status during the last 8500 years, (ii) investigate when anoxic conditions developed in
95 the hypolimnion, and (iii) assess whether the occurrence of hypolimnetic anoxia can be linked
96 to climatic changes in Central Tian Shan. We further evaluate whether phase boundaries
97 identified in the multi-proxy dataset can be attributed to event-based phase transitions or more
98 gradual shifts using multivariate statistical analyses. Comparisons with other
99 palaeolimnological data from Lake Son Kol and regional records will be used to understand
100 regional climate development in the Tian Shan, as well as implications for lake stratification

101 dynamics during the Holocene.

102

103 **2. Study site**

104 *2.1. Geological, environmental and lacustrine setting*

105 The geology, geomorphology, climatic and lacustrine characteristics of Lake Son Kol have
106 been extensively studied (Mathis et al., 2014; Huang et al., 2014; Lauterbach et al., 2014).
107 Son Kol (41°50'52"N, 75°07'45"E, 3016 m a.s.l.) is the second largest natural lake in
108 Kyrgyzstan, within the central Tian Shan mountain ranges (Fig. 1). It fills the central
109 depression of an intramontane basin located between two generally east-west stretching
110 mountain ranges, which are the Son Kol Too and Baidula Ranges (3200–3500 m a.s.l.) in the
111 north and the Boor-Albas Range (3800–4000 m a.s.l.) in the south (Shitnikov, 1980). These
112 mountain ranges are composed of Carboniferous sedimentary rocks and Permian granitoids
113 (east and south) and Cambro-Ordovician metamorphic granodioritic and granitic gneisses, as
114 well as strongly folded sedimentary rocks in the north
115 (<http://portal.onegeology.org/OnegeologyGlobal/>). The modern equilibrium line altitude
116 (ELA) of glaciers at north-facing slopes in the Central Tian Shan is ~3850–4100 m a.s.l.,
117 (Narama et al., 2007; Koppes et al., 2008), hence there are no glaciers in the lake watershed
118 today and, based on ELA reconstructions, seem unlikely to have fed Lake Son Kol during the
119 Holocene (Savoskul and Solomina, 1996). The widespread pasture-covered plains
120 surrounding the lake are made up of Quaternary eroded material arranged in old Pleistocene
121 moraines (Abramowski et al., 2006; Narama and Okuno, 2006) and inactive fan deltas. A
122 large fluvial fan-delta nourished by cirque valleys of the Boor Albas range progrades from the
123 southern shore into the lake. Along the southeastern and northeastern shore, large coastal
124 sedimentary features such as sand spits and associated open lagoons have been developed by
125 the prevailing westerly winds, indicating a stable lake level during at least the Late Holocene.

126 Today, Lake Son Kol is located above the treeline, on a high plateau surrounded by mountain
127 meadows. Because of pronounced altitudinal changes over short distances, the vegetation
128 rapidly changes within a few kilometers with the development of a lush alpine meadow in the
129 lake catchment at ~3000 m a.s.l., while conversely steppe to semi-desert environments prevail
130 in open lowlands (below 2000 m a.s.l) with sagebrush steppe (*Artemisia*) and
131 Chenopodiaceae (Mathis et al., 2014).

132 Lake Son Kol is ~29-km long and ~18-km wide, with a surface area of ~275 km² and a
133 catchment of ~1130 km². The lake water volume is ~2.82 km³ and the lake is presently ~13-m
134 deep (Lauterbach et al., 2014). According to the lake stratification classifications of Kalff
135 (2002), the depth and climatic location of Lake Son Kol should make it susceptible to shifts in
136 mixing regime. The lake has a single outlet at its southern end, which discharges into the
137 Naryn River and then to the Syr Darya down to the Aral Sea basin. Apart from direct
138 precipitation on the lake surface, most surface inflow is from several small mostly episodic
139 rivers fed by rainfall, groundwater and snowmelt from neighbouring mountains (Huang et al.,
140 2014). A few perennial rivers supply the lake from the west where they braid in a small
141 alluvial plain at the shore. Today, however, most of the inflow is contributed by groundwater
142 (Beloglasov and Smirnova, 1987). Average lake water temperature fluctuates between 16 °C
143 (14.5 °C in summer 2007; Huang et al., 2014) and 0–2 °C in winter, with almost no stable
144 thermal stratification throughout the year (Huang et al., 2014; Lauterbach et al., 2014). Earlier
145 monitoring data reported average summer water temperatures of 12–13 °C (Beloglasov and
146 Smirnova, 1987), with maxima of 20–23 °C during warmer years. Freezing starts in October
147 and ice break-up may begin as early as March (up to late April) depending on short-term
148 climatic variability (Huang et al., 2014).

149 The water column of Son Kol is well mixed and oxygenated to the lake bottom in the
150 summer, though a small oxygen minimum has been noted at a water depth of 1.5 m revealing

151 respiration in the epilimnion (July 2007; Huang et al., 2014). Monitoring data from the 1960s
152 and 1970s similarly reported a well-oxygenated water body throughout the whole year, but
153 Beloglasova and Smirnova (1987) observed oxygen concentrations of 0.8–1.8 mg.l⁻¹ or less in
154 the shallow water zone during winter. The strong fetch (maximum effective fetch of ~17 km;
155 Håkanson et al., 1983) and the high wind frequency (wind speeds up to 20 km.h⁻¹ in summer
156 and fall) impart efficient and complete mixing of the shallow water body. According to
157 previous surveys the weather is calm only during 11% of the ice-free season (Beloglasov and
158 Smirnova, 1987). In line with this, surface/intermediate flow currents reaching 2–8 cm.s⁻¹
159 across the water body (and up to 10 cm.s⁻¹ in the deepest part of the basin) reflect a highly
160 dynamic lacustrine system (Beloglasov and Smirnova, 1987). Such conditions guarantee for a
161 steady mixing of the whole water column and thus makes Son Kol mostly a polymictic lake
162 nowadays (apart from under ice in winter). Son Kol is a moderate hard-water lake (4.56–4.96
163 mg⁻¹) with a mineralisation of 465 mg.l⁻¹ (summer) and 306 mg.l⁻¹ (winter) (Beloglasov and
164 Smirnova, 1987). Accordingly, today it is a bicarbonate lake with weak magnesium
165 dominance. Conductivity throughout the water column ranged between 515 and 530 mS.cm⁻¹
166 in the summer of 2012 (Schwarz et al., 2017).

167

168 2.2. Climatic setting

169 Local climatic conditions lead to short temperate summers and severe cold winters, with snow
170 cover between November and April. Mean annual air temperature is -3.5 °C (mean of 11 °C
171 in summer and -20 °C in winter), based on the nearest meteorological station (Dolon, 41.8° N,
172 75.8° E, 3040 m a.s.l.) (Shnitnikov, 1980). Mean annual precipitation is 500–600 mm at Lake
173 Son Kol, with a maximum in summer and a minimum in winter (Academy of Science of the
174 **Kyrgyz** SSR, 1987). The current precipitation regime in the Tian Shan Mountains
175 surrounding Son Kol is mainly regulated by the interaction between cyclonic activity driven

176 by the mid-troposphere Westerlies and the southwestern branch of the Siberian anticyclonic
177 circulation (Aizen et al., 1997). Atmospheric dust concentrations (and loess) are maximal
178 below 3000 m a.s.l in Central Asia, with thickest loess accumulations occurring between 1000
179 and 1300 m a.s.l in the piedmont regions of the Tian Shan (Machalett et al., 2008). Different
180 airflows govern convective surface winds in western Central Asia and thus, the regional
181 distribution of remote dust particles over the year (Bugayev et al., 1957; Shettler et al., 2014).
182 During the summer northwestern, western and northern air intrusions and northeast
183 anticyclonic flows control up to 83% of the local wind directions. Dust monitoring near
184 Bishkek reveals besides that wet deposition is a substantial process for aerosol deposition in
185 mountain areas (Shettler et al., 2014). Nonetheless, backwards trajectories calculated from
186 Son Kol document that remote air masses likely also come from southern (Afghanistan) and
187 further western (Mediterranean) and northern (North Atlantic) sources (Shettler et al., 2014).

188

189 **3. Material and methods**

190 *3.1. Lake coring and cross-core correlation*

191 Four sediment cores, including two sets of parallel cores, were extracted with a Uwitec piston
192 corer in July 2007 from two different sites in the southeastern part of Lake Son Kol (Fig. 1).
193 Core SK04 (133cm) was obtained at ca. 1.15 km from the southern shore (41°46'56.6"N,
194 75°08'18.2"E, 10.5 m water depth) while a second set of parallel cores, SK06 and SK07 was
195 retrieved at 1.6 km northeast of core SK04 (41°47'45.2"N, 75°08'22"E, 12.5m water depth)
196 (Fig. 1). Although cores SK04 and SK07 have been extensively analyzed recently (Mathis et
197 al., 2014; Huang et al., 2014; Pacton et al., 2015), no integrative study has been undertaken on
198 core SK06 yet, although it represents the longest lacustrine archive ever retrieved in Lake Son
199 Kol. In spite of the significant distance between the two coring sites, the cores exhibit very
200 similar sedimentary successions (and lithological marker layers), thus allowing a robust core-

201 to-core correlation (Fig. 2a), as also demonstrated by the good match between magnetic
202 susceptibility records for cores SK04, SK06 and SK07 (Fig. S1). Core SK06 was split in half
203 lengthwise and photographed directly after core opening. A detailed macroscopic description
204 established the different sedimentary facies (Units I-VI) as well as the stratigraphic
205 correlations between cores SK04, SK06 and SK07 (Fig. 2a). No compaction and/or
206 disturbance was observed in SK06.

207

208 *3.2. Age-depth model*

209 Owing to the robust core-to-core correlation between cores SK06, SK04 and SK07 (Figs. 2
210 and S1) and SK06 and SONK_11_D1/2 from Lauterbach et al. (2014) (Fig. S2), the
211 chronology of core SK06 (Fig. 2b) was established using available radiocarbon (AMS ^{14}C)
212 dating of both terrestrial plant macroremains and bulk sediment organic matter (OM),
213 complemented by one short-lived radionuclide measurement (^{241}Am). We selected and
214 reappraised 8 bulk sediment samples and one terrestrial leaf from cores SK04 and SK07
215 (Huang et al., 2014) along with one terrestrial plant remains from core SONK_11_D1/2
216 (Lauterbach et al., 2014) (Table 1). ^{14}C age anchor points were transferred to core SK06 using
217 a detailed correlation of both lithological marker layers and magnetic susceptibility between
218 core SK06 and SK04 and SK07 (Figs. 2a and S1) and between cores SK06 and
219 SONK_11_D1/2 (Fig. S2). For the latter, the correlation at 33 cm in core SK06 of Poz-44073
220 date (at 33 cm in SONK_11_D1/2; Lauterbach et al., 2014) is a robust anchor point for the
221 core SK06 chronology, making the Poz-26363 radiocarbon age at 30 cm too old, and so this
222 date was rejected from the age model. Note that not all the radiocarbon ages reported in
223 Huang et al. (2014) were applied to core SK06 by cross-core correlation, since we only rely
224 on age points that could be confidently transferred to core SK06. Poz-26363, Poz-26370 and
225 Poz-26372 dates were rejected because they imply the reworking of older material in younger

226 sediments. Poz-37470 date (bulk sediment organic matter) was not used either because a
227 terrestrial leaf dated in the same stratigraphic level was preferred (Poz-37469). Other
228 discarded ages correspond to depths that could not been matched properly between cores
229 SK04 and SK06 (Poz-26368).

230 A 150-yr reservoir age correction was further applied prior to calibration to each transferred
231 conventional radiocarbon age by using the 155 ± 30 ^{14}C yr BP reservoir age obtained on live
232 deep-water algae by Lauterbach et al. (2014) (Table 1). Such a small reservoir age offset
233 implies that the recent hardwater effect is small; this is further confirmed by nearly identical
234 radiocarbon ages (within error bars) obtained on one bulk organic sample (not used here) and
235 a terrestrial leaf from the same stratigraphical level in core SK04 (see Table 1 and Huang et
236 al., 2014). More detail regarding reservoir effects in Lake Son Kol is provided in Lauterbach
237 et al. (2014).

238 In order to constrain the age of the topmost sediments in core SK06, we used the close
239 agreement of lithological markers in cores SK06 and SONK_11D1/2 and assumed that the
240 notable increase in ^{241}Am activity at 5.5 cm in SONK_11_D1/2 occurs at 2 cm in core SK06
241 (Fig. S2), and corresponded to the Northern Hemisphere nuclear weapons testing maximum
242 in 1963 CE (Appleby et al., 1991). The topmost ca. 2 cm of SONK_11_D1/2 are missing in
243 core SK06. Finally, the age-depth model for core SK06 was established using 10 corrected
244 conventional radiocarbon ages (Table 1), as well as the ^{241}Am anchor point at 2 cm sediment
245 depth (most likely 1963 CE), as input parameters for a *P_Sequence* deposition model
246 implemented in Oxcal 4.4 (Ramsey, 2008) (Fig. 2b). The *k* parameter was set to 1 (as in
247 Huang et al., 2014) and the radiocarbon ages were converted to calendar years by using the
248 IntCal20 terrestrial calibration curve of Reimer et al. (2020). The high agreement index A_{model}
249 of 89.8% indicates that the chronology of core SK06 is reliable (Ramsey, 1995; 2001).

250

251 *3.3. Geochemistry*

252 Major elements in the sediments were measured by XRF at high-resolution (500- μm step
253 size) on the split-core surface with an Avaatech Core Scanner (EDYTEM Laboratory). Prior
254 to measuring, the core surface was cleaned and foiled to prevent contamination and core
255 desiccation. The X-ray beam was generated with a rhodium anode and a 125- μm beryllium
256 window, which allows a voltage range of 7–50 kV and a current range of 0–2 mA.
257 Geochemical data were obtained with different settings according to the elements analyzed.
258 They were adjusted at 10 kV and 1 mA for 10 s to detect Si, Ca, Al, Fe, Ti, K, Mn and S. For
259 heavier elements (i.e., Sr, Rb, Zr, Br and Pb), measurements were performed at 30 kV and
260 9.75 mA for 30 s. XRF data are displayed as element intensities in count per second (cps)
261 (Fig. S3). Because no major element concentration was conducted on core SK06, we rely here
262 on XRF core-scanner data available on core SK07 (unpublished data) in order to calibrate
263 XRF measurements in core SK06. Major element concentrations on core SK07 were
264 previously measured on 1.5-cm thick discrete samples using XRF at the University Lyon1.
265 The sampling was conducted at a step of about 3–5 cm to allow for one calibration
266 measurement every 3–5 core scanner measurement in core SK07.

267

268 *3.4. Hyperspectral imaging*

269 Hyperspectral imaging is a visible and near infra-red spectroscopic method measuring the
270 reflectance of the sediment surface exposed to an incident light, and thus the sediment
271 composition (Barranco et al., 1989; Balsam and Deaton, 1996; Balsam et al., 1997).
272 Prior to analysis, the thin, superficial layer was scraped from the sediment to ensure a uniform
273 surface. Reflectance spectroscopy was performed on a core logger bench equipped with two
274 hyperspectral cameras (VNIR-PFD and SWIR, SPECIM®), which offers a short sampling
275 time and allows for high spatial resolution (several dozens of μm ; Jacq et al., 2020). The

276 distance between the studied sample/core and the camera lens was 130 mm, with an
277 acquisition angle of 90°. The surface of the sediment is indirectly illuminated by 18 halogen
278 bulbs homogeneously distributed around the lens. The acquisition speed of the core logger
279 was 0.5 mm.s⁻¹. Pixels in the resulting Visible and Near Infrared (VNIR) image each cover an
280 area of 47×47 μm² and contain raw spectral data from 400 to 1000 nm at 6.5 nm spectral
281 resolution (given in digit number, DN), whereas the Short Wave-Infrared (SWIR) image
282 pixels have a resolution of 150×150 μm² and cover a spectral range from 1000 nm to 2500 nm
283 at 10 nm spectral resolution. Therefore, each pixel contains spectral information related to
284 sediment composition. The depth of field of the imagers is ca. 2 mm. A normalization step is
285 required to convert raw data (8 bits) to % reflectance. At each acquisition, a 100% reflective
286 SPECTRALON standard was measured and used as a reference to express the data as %
287 reflectance. Data processing was carried out with the ENVI (v. 5.3) and IDL 8.0 softwares.
288 Pigment composition and concentrations were inferred from area indices (Fig. 3), which
289 directly measure the surface area (here referred to as relative absorption band area or RABA;
290 Trachsel et al., 2010) between the continuum and the reflectance spectrum at the absorption
291 band targeted (for instance for the chlorophyll-a and its derivatives at 670 nm). The sum of
292 the reflectance bands located throughout the absorption band is subtracted from the area under
293 a line connecting the beginning and the end of the depression formed by the absorption band.
294 In this study, area indices are followed by wavelengths indicating the width of the absorption
295 band considered (e.g. RABA630-700). The main absorptions are determined by continuum
296 removal, which also allows estimating the most interesting wavelength interval. In this study,
297 we have defined four of them centered on 670 nm, 615 nm, 844 nm and 1670 nm (Fig. 3).

298

299 *3.5. Coprophilous fungal spore analysis*

300 Catchment erosion processes in high-altitude landscapes may be re-shaped by livestock
301 grazing related to pastoral activity (Giguet-Covex et al., 2011). The variability of grazing
302 pressure and associated soil erosion in a catchment area can be reconstructed from the
303 sedimentary abundance of coprophilous fungal ascospores in lake sediments, as a proxy for
304 the density of wild and domestic herbivores (Davies and Schaefer, 2006; Gauthier et al.,
305 2010; Etienne et al., 2013). Son Kol catchment is still an area of transhumance today;
306 therefore, domestic herbivores are assumed to be the primary source of coprophilous fungal
307 spores in lake sediments, and wild herbivores are regarded to be a trivial source. To evaluate
308 the impact of grazing activity on soil erosion in the catchment, abundances of coprophilous
309 fungi ascospores were measured in 37 samples of core SK06 (for the last 6000 years). The
310 resolution is approximately 100 years from 6000 to 5000 cal. yr BP, and about 250 years from
311 5000 to 1500 cal. yr BP. Samples were chemically prepared according to the procedure of
312 Faegri and Iversen (1989). *Lycopodium clavatum* tablets (averaging 20,848 spores per tablet)
313 were added to each sample (Stockmarr, 1971) to express the results as influx or accumulation
314 rates (number.cm⁻².yr⁻¹). Coprophilous ascospores were identified and counted using a
315 photonic microscope following pictures and descriptions in van Geel and Andersen (1988),
316 van Geel et al. (2003), Van Geel and Aptroot, 2006, Feeser and O'Connell (2009), Cugny et
317 al. (2010) and Gelorini et al. (2012). Total coprophilous fungi *sensu stricto* include *Delitschia*
318 (UG-1066), *Neurospora* sp. (HdV-1), *Podospora*-type (HdV-368), *Sordaria*-type (HdV-55A),
319 *Sporormiella* (HdV-113), and *Trichodelitschia* (HdV-546).

320

321 3.6. Data analysis

322 Linear dimension reduction was performed by applying a correlation matrix-based principal
323 component analysis (PCA) using the PAST v. 4.02 software (Hammer et al., 2001). The PCA
324 was supplied with XRF (major elements) and hyperspectral data from core SK06 (Fig. 4).

325 Due to very high levels of non-normality and heteroscedasticity of the original descriptors,
326 data were first log-transformed and then separately centered on a per-sample basis in order to
327 remove overall-count effects and to focus on relative variations among samples. Also using
328 the PAST software, time series corresponding to the resulting PC1 and PC2 were further
329 investigated using Gallagher et al.'s (2011) Bayesian "changepoint modeling" algorithm for
330 identifying abrupt regime shifts in time series, often referred as tipping points in geosystems
331 (Lenton et al., 2008; Lenton, 2011; Scheffer et al., 2012). Number of MCMC iterations
332 required by Gallagher et al.'s method was set to 1,000,000 with a burn-in phase of 20,000
333 iterations; up to 300 changepoints were allowed a priori in order to generate fully-
334 unconstrained changepoint models, resulting in an average number of changepoints of 74 and
335 61 for the PC1 and PC2 time series, respectively.

336

337 **4. Results and interpretation**

338 *4.1. Sedimentology*

339 Sediments mostly consist of alternating yellowish and light-brownish calcareous muds in the
340 upper 1.22 m of core SK06, whereas dark organic-rich muds alternating with yellow-brown
341 calcareous muds and white carbonate layers predominate between 1.22 and 1.91 m (Fig. 2).
342 The transition between the basal organic-rich sediments and the upper 1.22 m alternating
343 calcareous muds is characterized by homogeneous greyish calcareous clays between 0.94 and
344 1.22 m. Apart from this transition, the sediments are finely laminated in core SK06, and
345 especially in dark organic sediments. XRF core scanning data allows deciphering the distinct
346 geochemical composition of the three main lithologies (Fig. S3). Dark organic sediments are
347 typified by their high abundances in Br and S (Figs. S3 and 5), whereas the intervening
348 carbonate layers exhibit an enrichment in Ca and Sr (Fig. S3). In contrast, the transition
349 interval (greyish homogenous clays) is best described by high abundances of Ti, Rb, Zr, Fe

350 and Si, emphasizing their high mineral content. The upper alternations between yellowish and
351 light-brownish sediments are characterized by a mixed geochemical composition and exhibit
352 fluctuating values of Ti, Rb and Zr, but also highly variable abundances of Ca and Sr (Fig.
353 S3).

354

355 *4.2. VNIR-SWIR proxy data*

356 Chlorophyll *a* (Chl *a*) and its derivative products (Rein and Sirocko, 2002; Das et al., 2005;
357 Wolfe et al., 2006; Michelutti et al., 2010) was measured as the area index between 630 and
358 700 nm (RABA630–700) normalized by the absorption maximum at 670 nm (R_{670}) (Fig. 3) as
359 defined by Wolfe et al. (2006). Although Chl *b* may contribute in the absorption range
360 between 630 and 700 nm (i.e., at 652 nm, Bidigare et al., 1990; Yacobi et al., 2015), it is not
361 observed in our spectral data (Fig. 3). We therefore assume that the measured absorption peak
362 of particles at 670 nm almost exclusively represents the optical activity of Chl *a*, apart from
363 the slight interference by the accessory chlorophylls Chl *b* and Chl *c*.

364 Phycocyanin is a water-soluble accessory pigment to chlorophyll exhibiting a notable
365 absorbance trough around 610–620 nm for C-phycocyanin (Glazer et al., 1973; Yacobi et al.,
366 2015). For this study, we propose to screen the phycocyanin content by measuring the area
367 index between 600 and 630 nm (RABA600–630) normalized by the absorption peak at 615
368 nm (R_{615}) as described by Yacobi et al. (2015) (Fig. 3). It must be recalled, however, that the
369 absorption of Chl *a* and accessory chlorophylls Chl *b* and Chl *c* partly overlap the range of
370 optical activity of phycocyanins, although weakly (Yacobi et al., 2015). However, the
371 RABA630-700 and RABA600-630 curves exhibit different patterns (Fig. 5), implying that a
372 genuine signal does exist apart from the Chl *a* plus derivatives. This also suggests that
373 phycocyanin predominates in some intervals where Chl *a* plus derivatives do not (Fig. 5), thus
374 allowing their respective trends to be distinguished and analyzed separately in core SK06.

375 Phycocyanins are common in cyanobacteria (Williams et al., 1980), and **their** concentration in
376 shallow lacustrine systems is used to estimate the presence and relative abundance of
377 cyanobacteria (Horváth et al., 2013). Phycocyanin has been reported in lacustrine sediments
378 of Lake Qinghai, Central China (Ji et al., 2009) and used to indicate cyanobacterial blooms in
379 the Baltic Sea (Woźniak et al., 2016). Chl *a* plus derivatives and phycocyanin contents are
380 thus interpreted here as a proxy for oxygenic phototrophs, which usually indicate aerobic
381 conditions. Total Bacteriopheophytin *a* (Bphe *a*) is a stable degradation product of
382 Bacteriochlorophyll *a* or Bchl *a* with a characteristic spectral absorbance in the infrared range
383 around 750–770 nm in ethanol extracts (Coolen and Overmann, 1998; Rogozin et al., 2012)
384 and 800–900 nm in sediments (Scheer, 2006; Ji et al., 2009; Blankenship, 2013). Bphe *a* was
385 determined from the area index between 750 and 900 nm (RABA750–900) normalized by the
386 absorption maximum at 844 nm (R_{844}) as defined by Butz et al. (2015) (Fig. 3). Because no
387 other known bacterial pigment absorbs at this wavelength, the relative abundance of Bchl *a*
388 (and its derivative by-products Bphe *a*) can be evaluated by measuring the Bphe *a* absorption
389 trough between 750 and 900 nm normalized by the absorption peak at R_{844} . Although
390 Bacteriopheophytin *b* (Bphe *b*) may also contribute at 820 nm (Oren, 2011), no absorption
391 peak is observed at that wavelength in our data (Fig. 3). We therefore assume that the
392 measured absorption peak of particles at 844 nm represents exclusively the optical activity of
393 Bphe *a*. Bphe *a* is produced by planktonic anoxygenic phototrophic bacteria (APBs) such as
394 purple sulphur bacteria (Chromatiales, Rhodospirillales; Schmidt et al., 2002; Rogozin et al.,
395 2012) which commonly live in the chemocline of stratified meromictic lakes, where reduced
396 sulphur species are **very abundant** and oxygen levels are low, if sufficient light is available for
397 photosynthesis (Butz et al., 2016). Hence Bphe *a* is a diagnostic pigment for photosynthesis
398 in anaerobic conditions and the presence of that pigment in the sediment may help infer long-
399 term lake stratification, and the presence of a chemocline (Fig. 5). Aromatic organic matter

400 content (ARO) has a characteristic absorption near 1670 nm (Wheeler, 1959) and it seems
401 that no other compound absorbs at this wavelength, so we assume that an index in this
402 spectral range can infer this property (Fig. 5). Nevertheless, we are aware that SWIR
403 measurements may be influenced by sediment moistness and grain size, for which no known
404 correction can be applied yet. For this study we propose the area index between 1660 and
405 1690 nm (RABA1660–1690), normalized by the absorption maximum at 1670 nm (R_{1670}).
406 ARO values are interpreted as a proxy for tracking the inputs of terrestrial organic matter
407 from the shore (e.g., soils) and related catchment, with high (low) ARO corresponding to
408 increased (decreased) inputs of detrital organic matter.

409

410 *4.3. PCA analysis*

411 XRF and hyperspectral (VNIR, SWIR) data, with a temporal resolution ranging between 3
412 yr.mm⁻¹ and 7–8 yr.mm⁻¹, were jointly analyzed by PCA in order to characterize sediment
413 end-members and detect abrupt shifts in the multivariate time series (Fig. 4a). The XRF core-
414 scanner of Ca, Fe and Ti was previously calibrated using XRF conducted on discrete samples
415 in core SK07 (28 samples), exhibiting an excellent match between the two methods (Fig. 4b).
416 PC1, PC2 and PC3 together explain about half of the total variance (Figs. 4a and S4),
417 allowing for the identification of different end-members in the lake system: (i) high Ti, Rb, Fe
418 and RABA1660–1690 values for PC1 > 0 (which are related to organic and minerogenic
419 inputs from the catchment), while Sr, Ca (which indicate endogenic carbonate precipitation)
420 and to a lesser extent Zr (coarse detrital input) drive PC1 < 0, making PC1 a proxy for
421 allochthonous vs autochthonous products in the lake system; (ii) high Mn, RABA600–630
422 and RABA630–700 values for PC2 > 0 (which are related to oxic conditions in the water
423 column) vs. high Br, S (and to a lesser extent Zr) and RABA750–900 values (which are
424 linked to organic matter content and infer reducing and sulphidic conditions in bottom waters)

425 for PC2 < 0, making PC2 a proxy for oxygenated vs. euxinic bottom water conditions in Lake
426 Son Kol. Based on this result, prominent environmental change occurred around 8400, 8200–
427 7800, 7700–7500, 7300–7000, 6500–6100, 6000–5700 and 5500–5250 cal. yr BP (Fig. S5),
428 when increased influx of detritus (PC1 > 0) is in close correspondence with the development
429 of euxinic conditions in the hypolimnion (and thus of a chemocline; PC2 < 0). The Si content
430 appears to slightly differ from other terrigenous (river-borne) elements (PC1 and also PC3
431 [14.4%], whose palaeoenvironmental signature is more difficult to interpret and is not
432 discussed here) (Fig. S4). Si-value is high in the silt fraction and is, in addition to its weak
433 riverine contribution, mostly linked with aeolian inputs into the lake (Huang et al., 2014).
434 Judging from the lack of any conspicuous link to the other elements (Fig. 4a) and given the
435 biochemical reactivity of this element, Mn is used to decipher bottom-water redox processes,
436 as already reported in other lacustrine settings (Wirth et al., 2013; Davies et al., 2015).

437

438 *4.4. Time series of hyperspectral and geochemical data*

439 Between 8500 and 7000 cal. yr BP (191–164 cm), phycocyanin (from cyanobacteria) has the
440 highest abundances downcore, probably related to the development of benthic cyanobacterial
441 mats in a shallow water body (Fig. 5). Chl *a* plus derivatives and Bphe *a* also oscillate at very
442 **high** frequency. Bphe *a* and ARO maxima occur repeatedly around 8500, 8400, 8200–7800,
443 7700–7500 and 7300–7000 cal. yr BP, corresponding to dark organic-rich sediments along
444 with high Br and S content. No match was found, however, between phycocyanin and Chl *a*.
445 Subsequently, Chl *a* plus derivatives and phycocyanin exhibit a pronounced decline between
446 7000 and 6500 cal. yr BP. Phycocyanin abundances drop from 7000 cal. yr BP onwards and
447 never recover to former levels. Bphe *a* and ARO values also exhibit a drastic decrease in this
448 time interval, as also do Br and S content.

449 From 6500–5250 cal. yr BP, Chl *a* plus derivatives values increase and reach their highest
450 abundances downcore, with however considerable fluctuations. Bphe *a* content recovers to
451 high levels and even dominates between 5750 and 5250 cal. yr BP, but dramatically decreases
452 and readily disappears from 5250 cal. yr BP onwards (Fig. 5). Coevally, Br, S and ARO
453 values are also high between 5750 and 5250 cal. yr BP (corresponding to the deposition of
454 dark organic sediments), but rapidly drop after 5250 cal. yr BP. Chl *a* plus derivatives and
455 phycocyanin exhibit a prominent decline in their abundances between 5000 and 3850 cal. yr
456 BP. Judging from XRF core-scanner data, enhanced detrital inputs occurred in this interval
457 (Fig. S3). Subsequently, Chl *a* plus derivatives and phycocyanin increase from 3850–3100
458 cal. yr BP, whereas Bphe *a* still oscillates at very low levels. Higher Br, along with enhanced
459 ARO, also parallels the occurrence of dark organic-rich sediments in this interval. Over the
460 course of the last 3000 years, oxygenic phototrophs fluctuate at lower levels, whereas Bphe *a*,
461 Br and ARO remain low (Fig. 5).

462

463 *4.5. Abrupt regime shifts*

464 Abrupt regime shifts were identified based on slope variations of the average changepoint
465 model calculated separately for PC1 and PC2. For each principal component, average
466 changepoint model's slope (absolute value) is plotted against time, with 99% and 95%
467 percentiles taken as cut-off values (Figs. 6 and 7). Time intervals with average changepoint
468 model's slope values above these cut-offs indicate abrupt changes in the time series, and can
469 therefore be regarded as tipping points. For PC1, tipping points are recognized as 10–20 yr
470 intervals (beyond the 99% percentile value) at ca. 3670, 5050–5080 (highest slope value),
471 5250, 5370, 5850, 6150, 6220, 6900, 7030, 7980, 8230, 8280 and 8340 cal. yr BP (Fig. 6).
472 For PC2, tipping points are similarly identified as 10–20 yr intervals (beyond the 99%
473 percentile value) around 5250 (highest slope value), 5950, 6040, 6230, 6280, 6380, 6670,

474 6700, 6900, 7060, 7750, 7800, 8320 and 8380 cal. yr BP (Fig. 7). Hence the time intervals
475 around 5050–5250, 6220–6240 and 8200–8400 cal. yr BP are detected for both PC1 and PC2,
476 and probably correspond to abrupt regime shifts in the lacustrine system. Interestingly, low
477 PC2 values match with a strong peak of magnetic susceptibility (Figs. S1 and S5) around
478 8200–8400 cal. yr BP, suggesting the occurrence of abrupt changes in the lake and
479 surrounding catchment that are tied with the 8200 cal. yr BP cold event widely recognized in
480 the Northern Hemisphere (Alley et al., 1997; Rohling and Pälike, 2005).

481

482 **5. Discussion**

483 Between ca. 8500 and 5250 cal. yr BP anaerobic conditions (and the penetration of light deep
484 in the lake to support photosynthesis for APB) are inferred from high Bphe *a* and Br (PC2 <
485 0), along with the occurrence of dark organic layers (Fig. 5). Coevally, high ARO and
486 minerogenics content (PC1 > 0) suggest enhanced inputs of detrital organic matter and a link
487 between terrestrial influx, sedimentary dynamics and the advent of hypolimnetic anoxia in the
488 earlier part of the record. After 5250 cal. yr BP, the near absence of Bphe *a*, lower ARO
489 values but persistently high abundances of oxygenic phototrophs suggest prolonged aerobic
490 conditions within a well-mixed water body (Fig. 5).

491

492 *5.1 Episodes of increased snowmelt and indicators of lake-level changes*

493 Grain size and geochemical/isotopic (TC, TN, TOC, C/N, $\delta^{13}\text{C}_{\text{org}}$ and $\delta^{18}\text{O}$) data can be useful
494 in supporting interpretations from hyperspectral and XRF data. Here we compare previously
495 unpublished and available data from SK07 (Huang et al., 2014) to help interpret the 8500–
496 5000 cal. yr BP time interval during which dark organic intervals are recurrently interbedded
497 with yellow-brown calcareous muds (Fig. 7). Altogether, these parameters help to distinguish

498 changes in onshore and in-lake environmental conditions such as sheet-wash from the shore,
499 productivity and nutrient availability, and the possible mechanisms at work.

500 Generally, the timing of changes in geochemical and isotopic proxies in core SK07 agree with
501 those of XRF and hyperspectral data from core SK06 (Fig. 8). Dark organic intervals with Br
502 and ARO in core SK06 have elevated TOC, TC, TN and C/N values but strongly depleted
503 $\delta^{13}\text{C}_{\text{org}}$ and $\delta^{18}\text{O}$ values in core SK07. Such prominent $\delta^{13}\text{C}_{\text{org}}$ depletions might indicate the
504 predominance of APB during phases of lake stratification because pronounced carbon isotope
505 fractionation towards lighter $\delta^{13}\text{C}$ has been associated with purple sulphur bacteria in
506 meromictic lakes (Zyakun et al., 2009). Alternatively, short-lasting depletions in $\delta^{18}\text{O}$ can
507 indicate seasonal shifts in the hydrological regime; i.e., increased winter precipitation and/or
508 increased meltwater discharges linked to erosive runoff during spring snow thaw, but also
509 increased temperatures (Huang et al., 2014; Lauterbach et al., 2014) that would have fueled
510 meltwater runoff. The dark organic-rich and $\delta^{18}\text{O}$ -depleted intervals co-occur with coarser
511 (e.g., silt- and sand-sized) minerogenic and organic (e.g. phytoclasts) inputs into the lake
512 (Figs. 8 and S6), whereas the intervening yellow-brown calcareous muds are enriched in finer
513 (clay-sized) detritus with low sand content (Fig. 8). Erosive runoff after thawing during spring
514 (Huang et al., 2014), and/or higher winter snowfall (Lauterbach et al., 2014), can account for
515 coarse detritus discharge into the lake. Interestingly, the PCA analysis reveals that higher
516 detrital inputs ($\text{PC1} > 0$) are not associated with elevated Zr content in core SK06 (Fig. 4);
517 instead, dark organic layers exhibit low Zr content (Figs. 8 and S3). Zr is usually more
518 concentrated in sand-size lacustrine sediments (e.g., Cuven et al., 2010; Davies et al., 2015)
519 and so, this Zr depletion in a few coarse dark organic intervals is puzzling, since otherwise all
520 detritics (Ti, Rb and Zr) exhibit similar trends in the intervening yellow-brown calcareous
521 muds (Fig. S3). Therefore, enhanced levels of Ti and Rb along with lower Zr (thus $\text{PC1} > 0$)
522 in the Br-enriched ($\text{PC2} < 0$) dark organic layers either suggest the dilution of Zr input by Ti-

523 and Rb-laden terrestrial organic matter delivered during snow thaw (as inferred from high
524 ARO values), and/or the dilution of Zr (associated with the coarse detrital fraction) during
525 relatively high lake water level phases. Zr mainly originates from erosion processes in the
526 upper catchment of Lake Son Kol because Zr is abundant in local Paleozoic intrusives
527 (granites and granodiorites), but these tend to be covered by Quaternary sediments and soils
528 (colluvial and glacifluvium deposits) on the shoreline belt and further up-valley. Although Zr
529 is linked to Br and S on PC2, Zr is also significantly associated with elevated Sr and Ca (PC1
530 < 0 ; Fig. 4), suggesting that Zr is enhanced during low lake water levels, which promotes
531 endogenic carbonate precipitation (Huang et al., 2014). Hence higher Zr infers the increasing
532 influence of riverine discharges at relatively low lake levels and the reduced distance between
533 the coring site and proximal detrital sources. Dilution of Zr most probably occurs during
534 short-lived increases in lake levels, which in turn enhance the formation of organic-rich
535 deposits in Lake Son Kol when elevated terrestrial inputs from snowmelt strengthened the
536 export of dissolved organic matter into the lake. The most probable sources of material during
537 snowmelt discharges (e.g., floods) are the abundant fan delta/fluviial material at the southern
538 edge of the lake, and/or superficial sediments on the slopes (e.g., colluvium) directly adjacent
539 to Lake Son Kol. In contrast, higher Ti, Rb and Zr occur coevally during lower lake level
540 phases, which correspond to the deposition of grey calcareous muds enriched in fine
541 siliciclastics (Fig. S3). Hence higher Zr, Sr and Ca values indicate lower lake water levels and
542 coincide with the deposition of yellow-brown calcareous muds. In contrast, dark organic
543 deposits enriched in Br, S, C/N and TOC are associated with the injection of large volumes of
544 sediment-laden sheet-wash from the shores (high ARO values) during intensified winter
545 precipitation and/or stronger snowmelt (Huang et al., 2014; Lauterbach et al., 2014), which
546 are likely to occur under warmer climate conditions.

547

548 *5.2. Redox conditions and oxygenation of the hypolimnion*

549 The geochemical behavior of Mn-enriched phases can be used to infer past redox conditions
550 and thus help diagnose oxic and sulphidic redox regimes in the water column (Fig. 9), as
551 reported from Lake Cadagno (Wirth et al., 2013). Based on elevated Mn content, oxic
552 conditions ($PC2 > 0$) prevailed around 8300–8200, 7400–7250, 5500–5400 and in particular
553 between 5000 and 3800 cal. yr BP in Son Kol when Mn was flushed to the lake with detrital
554 input, and probably occurred in the water column as dissolved unstable Mn^{2+} or as oxidized,
555 particulate Mn-(oxyhydr)oxides. However, no Mn-(oxyhydr)oxides were found in SK06,
556 possibly due to a reduction of Mn^{2+} (leading to a further recycling of Mn into the water
557 column) if the sediment-interface and/or the sediments were reducing. Oxic water column
558 conditions were accompanied by the precipitation of various carbonate minerals judging from
559 the regular occurrences of carbonate layers (Pacton et al., 2015), as well as high XRF-Ca
560 content in the yellow-brown calcareous muds separating dark organic-rich sediments, thus
561 reflecting the high alkalinity of lake waters during those time intervals. The regular
562 occurrence of sediment-dwelling ostracods and molluscs (which require well-oxygenated
563 bottom waters) further supports that a well-mixed (polymictic) water body, with oxic bottom
564 waters, has been established during the last 5000 years (Huang et al., 2014), in accordance
565 with recent monitoring data (Lauterbach et al., 2014).

566 In contrast, there is compelling evidence that the hypolimnion recurrently turned to a
567 sulphidic (euxinic) redox regime between 8500 and 5250 cal. yr BP, involving a stabilization
568 of the water-column stratification and the full development of a chemocline. Low sedimentary
569 Mn concentrations (Fig. 9), the absence of carbonate phases and the drastic increases in Br
570 and S content ($PC2 < 0$) characterize several phases of euxinic bottom waters between 8500
571 and 5250 cal. yr BP. Coevally, high Bphe *a* values (inferring that purple sulphur bacteria
572 thrived in the water column), and the rising TOC content in dark organic-rich sediments (Fig.

573 8), are good indicators of reducing bottom water conditions. Although such a hypothesis was
574 ruled out by Lauterbach et al. (2014) based on the absence of varves and/or vivianite crystals
575 (the latter indicating anoxic conditions in a non-sulphidic environment during sediment
576 deposition, which is obviously not the case here) in dark organic-rich sediments, several lines
577 of evidence advocate for the recurrence of hypolimnetic euxinia (and thus the development of
578 a chemocline) in Lake Son Kol before 5250 cal. yr BP. Prolonged lake stratification and
579 reducing oxygen conditions around 8500, 8400, 8200–7800, 7700–7500, 7300–7000, 6500–
580 6100, 6000–5700 and 5500–5250 cal. yr BP (Fig. 9) are inferred from (1) high TOC contents
581 (up to 16%) in dark organic-rich layers (Fig. 8) that imply both reducing bottom water
582 conditions which prevents organic matter degradation and especially of photosynthetic
583 pigments (Figs. 7 and 9); (2) the unusual preservation of numerous colonies of coccoid
584 prokaryotes altogether with the preserved intracellular content of cyanobacteria that requires
585 anoxic conditions (Pacton et al., 2015) and in particular, (3) the regular occurrence of Bphe *a*
586 pigments that indicate photosynthesis in anaerobic conditions and thus hypolimnetic anoxia
587 between 8500 and 5250 cal. yr BP.

588

589

590 *5.3. Origin and driving mechanisms of detrital-rich organic layers in Lake Son Kol*

591 As developed previously, the onset of euxinic bottom water conditions in Lake Son Kol
592 between 8500 and 5250 cal. yr BP most likely occurred under warmer climate conditions that
593 fostered meltwater runoff. Climate warming possibly fostered permafrost melting, which
594 activated the subaquatic springs that in turn fed the lake with solute-rich waters. Such
595 conditions are highly favorable for the sustainment of stable lake stratification and the
596 development of a chemocline, as observed in other high-altitude lake settings (Wirth et al.,
597 2013). However, this seems to be inconsistent with the regular occurrence of coarse material
598 in dark organic-rich sediments, which implies the transport of organic and minerogenic

599 detritus over long distances in Lake Son Kol. In meromictic lakes, the deposition of coarse
600 detritics in a coring site today located more than 3 km from the shore could be attributed to
601 hyperpycnal or mesopycnal currents (Mulder and Chapron, 2011), depending on the density
602 contrast between inflowing and lake waters. Hypopycnal flows are not considered here
603 because hypopycnal flows in lakes generally supply small quantities of sediments and would
604 require further reconcentration processes in the lake (over a 3-km distance in the case of Lake
605 Son Kol). Aeolian deposition of coarse-grained detritics (including large phytoclasts, Fig. S6)
606 is unlikely as we would expect much finer grain sizes in the case of loess deposits (i.e., silts)
607 (Huang et al., 2014), and loess deposits are generally restricted to lower altitudes in the
608 piedmont of the Tian Shan Mountains (Machalett et al., 2006; Lauterbach et al., 2014).
609 Finally, increased coastal wave erosion driven by winds can be rejected as well because
610 higher wave activity would also have broken down the chemocline. Moreover, hyperpycnal
611 deposits are often typified by sharp-based lag rich in organic debris (Mulder and Chapron,
612 2011), which is never encountered in core SK06 where coarse detritics are rather
613 homogenously distributed in the organic-rich layers (apart from the occurrence of a few
614 normal graded beds; Fig. S6). Hence, the most reasonable candidates for the deposition of
615 coarse minerogenic and organic detritus at the coring site are homopycnal or mesopycnal
616 flows. Homopycnal flows result from a small density contrast between the entering flow and
617 the surrounding water (Bates, 1953) and can export suspended sediment load either in
618 stratified or non-stratified lake basins (Mulder and Chapron, 2011). Alternatively, mesopycnal
619 flows are partially buoyant mesopycnal flows (interflows) of turbid plumes forming along
620 each pycnocline (that separates each water mass) (Van Rensbergen et al., 1998, 1999;
621 Schröder et al., 1998; Chapron et al., 2004, 2005; Støren et al., 2010; Mulder and Chapron,
622 2011).

623

624 Formation of mesopycnal currents is promoted when the inflowing waters enter more saline
625 lake waters, as occurred between (at least) 6000 and 5000 cal. yr BP when higher
626 conductivities (very low lake levels) were inferred in Lake Son Kol (Schwarz et al., 2017).
627 Modern conductivity of Lake Son Kol (515–530 mS.cm⁻¹) is about twice that of meromictic
628 Lake Bourget in France (300–350 mS.cm⁻¹). Therefore, mesopycnal (or homopycnal) flow
629 deposits occurred during periods of strong snowmelt (and/or increased temperatures),
630 indicated by the strong depletion of $\delta^{18}\text{O}$ in dark organic layers (Fig. 8), thus implying the
631 remobilization of sediments from the lake catchment whenever sufficient stream discharge
632 occurred to form a turbid plume. Two main factors control floods triggered by snowmelt: the
633 snowmelt potential and the actual melting. Snowmelt depends on the amount of seasonal solid
634 winter precipitation (e.g., snow) in the catchment while the intensity of melting is mostly
635 controlled by spring/summer temperature (Støren et al., 2010). During the warm Mid-
636 Holocene in Central Asia (Chen et al., 2008), periods of higher solid winter precipitation
637 coupled with relatively cool springs and warm summers may have favored increased
638 availability of snow when early summer temperatures started to rise in the mountains,
639 intensifying spring melt floods and the amount of terrestrial material washed into the lake.
640 Convective summer rainstorms during warm summers could also lead to lower lake water
641 $\delta^{18}\text{O}$ values during the Holocene thermal maximum. Intensive rainstorms in the summer
642 would increase hillslope erosion and stream discharge at a time when the lake was probably
643 more saline, stratified and shallow due to more intensive evaporation. Increased intensities of
644 rainstorms and snowmelt are therefore both possible drivers of mesopycnal or homopycnal
645 flows (and coarse sediment deposition) in Lake Son Kol. Alternatively, hypolimnetic anoxia
646 could have developed seasonally, by occurring under ice during the winter when organic
647 matter decomposes (if there is a limited snow cover that enables photosynthesis) and breaking
648 down towards the end of summer. However, this hypothesis is unlikely as (1) very few graded

649 beds related to ice melt are observed (i.e., resulting from the settling of fine-grained aeolian
650 particles deposited on ice to the lake bottom), (2) the regular occurrence of large phytoclasts
651 in dark organic layers with coarse minerogenic detritus (Fig. S6) implies a soil erosion from
652 the catchment, and thus is not compatible with aeolian depositional processes and, (3) Bphe *a*
653 data exhibit multi-annual periods of stratification (i.e., meromixis), rather than a seasonal
654 signal (Fig. 10).

655

656 *5.4. Holocene chemocline development*

657 The prominent changes in oxygenation conditions in the lake water column led to distinctive
658 changes in Lake Son Kol autotrophs (Figs. 9 and 10). Bphe *a* is produced by anoxygenic
659 phototrophic bacteria (APBs) which commonly live in the chemocline (oxic/anoxic transition)
660 of meromictic lakes (Overmann et al., 1991; Rogozin et al., 2012; Blankenship, 2013; Butz et
661 al., 2016). APBs require nutrients, sulphides and light to flourish, as provided by stable
662 chemocline (Parkin and Brock, 1980; Van Gemerden and Mas, 1995, Hodgson et al., 1998;
663 Madigan, 2003), and thus APBs may form dense populations in meromictic lakes (Van
664 Gemerden and Mas, 1995; McGowan et al, 2008). Such conditions usually develop when
665 lakes permanently stratify, but oxygen drawdown **could** also develop under ice with
666 respiration of organic matter, potentially leading to anoxic conditions during certain times of
667 the year when light is available and APB could flourish. Since dark organic-rich sediments
668 represent more than a single year in duration (Figs. 5 and 9), Bphe *a* suggest that the stability
669 of stratification increased and that there may have been periods of prolonged hypolimnetic
670 anoxia in Lake Kol during the Mid-Holocene. Recurring Bphe *a*, Br, S and ARO positive
671 anomalies (i.e., PC2 < 0) at 8500, 8400, 8200–7800, 7700–7500, 7300–7000, 6500–6100,
672 6000–5700 and 5500–5250 cal. yr BP thus imply periods characterized by the development of
673 a chemocline, which are tied to increased terrestrial inputs into the lake (PC1 > 0) (Fig. 9). A

674 very similar assemblage of spectral signatures (i.e., Bphe *a* and phycocyanin) was reported
675 from Lake Qinghai (China), where higher Bphe *a* concentrations are tied to elevated
676 freshwater inputs and increased warmth between ca. 8200 and 5500 cal. yr BP (Ji et al., 2009)
677 (Fig. 10). This also coincides well with higher (alkenone-inferred) summer temperatures in
678 Lake Balikun and Lake Qinghai at 8500–5000 cal. yr BP (Hou et al., 2016; Zhai et al., 2017).
679 Warmer summers in Lake Jaczno (Poland) were also linked to the onset of lake meromixis
680 (Butz et al., 2016). Besides, vegetation expansion in lake catchments during warm periods
681 (and more specifically the expansion of **broad-leaved** trees, such as birch) is known to
682 increase dissolved organic matter (DOM) fluxes to lakes and **decrease** UV light penetration
683 (Sommaruga and Psenner, 1997; Schmidt et al., 2002). At Lake Son Kol, the presence of trees
684 in the catchment and around the lake probably influenced lake trophic status and mixing
685 conditions by (1) stimulating pedogenesis, which fueled oxygen consumption rates into the
686 water body due to enhanced organic compounds exported to the lake and (2) sheltering the
687 lake surface from wind-induced mixing (i.e., reduced wind exposure) thus favoring a
688 prolonged time of stratification. Interestingly, the initiation of euxinic conditions in the
689 hypolimnion (coeval to the deposition of dark organic-rich layers) followed the mass
690 expansion of birch and arboreal **taxa** in the catchment between 8500 and 5000 cal. yr BP (Fig.
691 10), which enhanced terrestrial inputs of DOM and thus organic matter decomposition in the
692 lake. Therefore pollen, geochemical and sedimentological proxies all suggest that
693 hypolimnetic euxinia developed during warmer summers, and probably milder
694 winters/springs. As proposed by Mathis et al. (2014), centennial-scale episodes of weakened
695 seasonality indeed occurred between 8500 and 5000 cal. yr BP, alongside increased winter
696 temperatures and warm summers hence promoting a longer growing season for the trees. Our
697 findings also agree with other speleothem records from Central Asia which indicate increased
698 moisture transport by the Westerlies, and higher winter/spring precipitation in the early part of

699 the Holocene (Fig. 10; Chen et al., 2016; Cai et al., 2017; Liu et al., 2019). In contrast, the
700 breakdown of lake stratification at 7500–7300 and 7000–6500 cal. yr BP probably coincides
701 with cooler climate conditions (and longer and thicker ice cover) linked to the strengthening
702 and southward position of the Siberian High in arid Central Asia. Increased influence of the
703 Siberian High would weaken the mid-latitude Westerlies and/or push their tracks further to
704 the south, resulting in comparably drier conditions in the Tian Shan (Aichner et al., 2015;
705 Wolff et al., 2017).

706

707 Enhanced nutrient concentrations (linked with higher soil development DOM supply to the
708 lake) are associated with a decrease in lake water transparency between 8500 and 7000 cal. yr
709 BP, judging from unstable values of oxygen phototrophs (Fig. 9). At roughly the same time,
710 phycocyanin abundances rapidly oscillate at high levels, suggesting that cyanobacteria
711 predominated over Chl *a* and derivatives. A similar pattern was reported from shallow lakes
712 in southern Sweden (Randsalu-Wendrup et al., 2014) and northern Germany (Dressler et al.,
713 2007). Therefore, cyanobacteria, which have the ability to sustain relatively high growth rates
714 at low light intensities (Mur et al., 1999), likely overtook oxygen phototrophs between 8500
715 and 7000 cal. yr BP, concurrently with the occurrence of APB blooms at the chemocline (Fig.
716 9). Before 5000 cal. yr BP, Lake Son Kol was a shallow and closed lacustrine system (Fig. 9;
717 Schwarz et al., 2017) and it is likely that a substantial proportion of the oxygen phototrophs
718 could derive from benthic algae (e.g., benthic diatoms and cyanobacteria). This also agrees
719 well with the occurrence of biofilms and/or microbial mat structures between 8500 and 5000
720 cal. yr BP (Pacton et al., 2015).

721

722 A major and abrupt change in lake regime is recorded around 5250 cal. yr BP in Lake Son
723 Kol, with a drastic decline of Bphe *a* concentrations that occurred within a decade or so (Figs.

724 9, 10). The abrupt decline in APB pigments indicates a rapid collapse of stratified conditions
725 and the breakdown of anoxic conditions in the hypolimnion after 5250 cal. yr BP, although
726 high Br and S are still recorded until ca. 5050 cal. yr BP. Because this event tipped the system
727 over a threshold (as observed in both PC1 and PC2 average model slopes; Figs. 6 and 7) into
728 another stable state **from** which former trophic and mixing conditions never recovered, we
729 consider this time interval as a tipping point in Lake Son Kol. A similar abrupt ecosystem
730 threshold was reported at 4900 cal. yr BP based on chironomid remains, and attributed to the
731 increased influence of the Westerlies causing enhanced precipitation and higher windiness at
732 Lake Son Kol (Schwarz et al., 2017; Laug et al., 2020). Drastic environmental change in the
733 lake may also be triggered by land-use processes in the catchment, which **are** mainly driven
734 by changes in grazing pressure at high altitudes. Abundance of coprophilous fungal
735 ascospores is assumed to reflect changes in grazing activity (Gauthier et al., 2010; Etienne et
736 al., 2013). Influxes of coprophilous fungi *sensu stricto* fluctuated between 0 and 10
737 spores.cm⁻².yr⁻¹ between 6000 and 4000 cal. yr BP and during the last 2500 years (Fig. 10),
738 suggesting the predominance of wild animals in the Lake Son Kol catchment. Nevertheless
739 the presence of domestic herbivores cannot be excluded during 6000–4000 cal. yr BP, which
740 would match the occurrence of human fecal biomarkers in Lake Chatyr Kol sediments around
741 6000–5000 cal. yr BP (Schroeter et al., 2020). The imprint of domesticated livestock and
742 human occupation was likely highest around 3700 and 2500 cal. yr BP, where influxes
743 reached their maxima. Interestingly, influx values were generally stable during the period
744 5500–5000 cal. yr BP, when an abrupt ecosystem shift occurred in Lake Son Kol (Figs. 9,
745 10), thus implying that **pastoralism** apparently did not have a significant impact in the
746 catchment and that climate is the main driver of the abrupt regime shift identified around
747 5250 cal. yr BP.

748 After 5050 cal. yr BP, the lake became persistently well-mixed and the chemocline no longer
749 existed in the lake water column, as indicated by pervasive PC2 > 0 values (Fig. 9) and
750 elevated Mn content between 5000 and 3850 cal. yr BP that argue for oxic conditions in the
751 lake. Surprisingly, the advent of well-mixed conditions is closely related to a strong decline of
752 oxygen phototrophs (including cyanobacteria) between 5050 and 3850 cal. yr BP, with strong
753 negative anomalies around 4400–3800 cal. yr BP, coeval to high Mn anomalies (Fig. 9).
754 Because this phase corresponds to a noticeable decline in summer humidity and low lake
755 levels (PC1 < 0; Fig. 9), we posit that cooler summer temperatures and longer lasting and
756 thicker snow cover may have prevented the growth of oxygen phototrophs in the water
757 column (as well as the establishment of a stable stratification in summer after ice cover).
758 Alternatively, regular wind-driven lake mixing resulting from strengthened winds during the
759 spring ice break-up season (inducing stronger vertical water circulation) may have **caused** low
760 pigment deposition and preservation in the sediments (Cuddington and Leavitt, 1999). The
761 disappearance of the chemocline after 5050 cal. yr BP coincides with a decrease in terrestrial
762 DOM inputs into the lake, as catchment vegetation opened up with a decline of birch and
763 other arboreal taxa and a rise in *Artemisia* (Fig. 10; Mathis et al., 2014), similarly as in Lake
764 Qinghai (Ji et al., 2009). Tree removal around the lake thus probably led to higher wind
765 exposure of the open-water area (due to increased wind fetch), inducing in turn a better
766 mixing of the water column. Lake levels **increased** around 3700 cal. yr BP, corresponding to
767 another main tipping point based on high PC1 values (Fig. 6), in line with increased
768 planktonic diatom abundances and lower conductivities (Fig. 9; Schwarz et al., 2017). High
769 PC2 (and low Bphe *a*) values from 3800 cal. yr BP onwards infer that repetitive mixing
770 conditions prevented the establishment of a chemocline by oxygenating the hypolimnion
771 (Butz et al., 2016), which hampers the onset of stable temperature stratification during
772 summer. Such conditions promoted nutrient cycling in the photic zone to support oxygenic

773 phototrophs in the epilimnion, which diminished light penetration to the chemocline and led
774 to the **destruction** of a suitable lake habitat for APB growth. Hence, all proxy data suggest that
775 Lake Son Kol has had persistently high lake levels with well-mixed and oxygenated
776 conditions for the last 5000 years, in accordance with its modern polymictic regime.

777

778 **6. Conclusions and perspectives**

779 Ultra-high temporal resolution geochemical and hyperspectral palaeolimnology data can yield
780 a more complete understanding of changing ecological patterns and redox processes in
781 shallow lacustrine systems. As a biomarker of planktonic APB at the chemocline of lakes, the
782 presence/absence and variability of Bphe *a* in Lake Son Kol sediments provides crucial
783 information about the timing and forcing mechanisms that control the advent and vanishing of
784 lake stratification over the last thousands of years. Increased Bphe *a*, Br, S abundances along
785 with higher terrestrial inputs from the catchment are recorded at 8500, 8400, 8200–7800,
786 7700–7500, 7300–7000, 6500–6100, 6000–5700 and 5500–5250 cal. yr BP, and involve
787 recurring episodes of hypolimnetic anoxia and euxinic bottom water conditions coeval to the
788 deposition of dark organic sediments on the lake floor. The presence of a chemocline and the
789 associated deposition of coarse minerogenic detritus transported by concentrated flows
790 appears to be dependent on recurrent seasonal snowmelt episodes, and thus on the quantity of
791 snow and spring/summer temperatures that regulate the potential for meltwater events. Our
792 data thus suggest that low lake levels and warmer climate conditions regularly prevailed
793 during 8500–5250 cal. yr BP, and promoted massive APB blooms in a well-stratified lake
794 system. After 5250 cal. yr BP, Son Kol water level **recovered** and the lake became persistently
795 well mixed, leading to the disappearance of the chemocline. This abrupt ecosystem shift was
796 associated with cooler conditions and strengthened wind forcing, when the influence of the
797 Westerlies increased from 5250 cal. yr BP onwards. Intensified wind forcing also most

798 probably coincided with a significant increase of wave activity at the lake shoreline acting as
799 an erosive agent. The imprint of wind forcing on coastal sedimentary dynamics (e.g., littoral
800 sand drift) and geomorphological processes can be observed based on the diagnostic
801 sedimentary features along the northeastern and southeastern shores of the lake, where sand
802 spits, wave-dominated beaches, barrier spits and associated open lagoons have developed,
803 hence suggesting that Lake Son Kol is presently a wind-driven water body (as defined by
804 Nutz et al., 2018). Further studies are required in order to properly date such
805 geomorphological features and assess the relationships between wind forcing, past lake levels
806 and climatic trends during the Mid- to Late Holocene.

807

808 **Acknowledgments**

809 We are grateful to Anne-Lise Develle (EDYTEM) for conducting XRF core-scanner analyses
810 on core SK06. The hyperspectral imaging acquisition was conducted at the University of
811 Normandie-Rouen. This research was funded by the Laboratory of Geology of Lyon (LGL-
812 TPE). All data from Son Kol core SK06 are available as supplementary data via the QSR
813 network.

814

815 **Figure captions:**

816

817 **Figure 1:** Location of Lake Son Kol: (a) global map with focus on western central Asia; (b)
818 central Tian Shan including south Kazakhstan and Lake Balkash (image courtesy of NASA
819 GSFC); and (c) bathymetric map of the lake, with isobaths given in meters below lake level
820 (modified after Academy of Science of the Kyrgyz SSR, 1987). Orange points refer to
821 sediment core SK06, while the two yellow and the white points indicate cores SK07 and
822 SK04 (Mathis et al., 2014; Huang et al., 2014; Pacton et al., 2015) and SONK_11_D1/2

823 (Lauterbach et al., 2014), respectively.

824

825 **Figure 2:** (a) Detailed stratigraphic correlation between cores SK06 (this study) and cores
826 SK04 and SK07 (Huang et al., 2014; Mathis et al., 2014) from Lake Son Kol using distinct
827 lithological marker layers. Studied cores are located on the map provided in Fig. 1. (b) Age
828 model for core SK06 based on the *P_Sequence* depositional model (model parameter $k=1$)
829 established using Oxcal 4.4 (Ramsey, 1995, 1998, 2001). The blue envelope corresponds to
830 the 2σ probability range. See text for detail.

831

832 **Figure 3:** (a) Visible signatures and (b) continuum removal spectra with associated main
833 absorptions for Chl *a* (and derivatives), phycocyanin and Bphe *a* (Bacteriopheophytin *a*) from
834 three selected intervals in which these pigments are abundant in core SK06.

835

836 **Figure 4:** (a) PCA biplot of major elements obtained by XRF core scanner and hyperspectral
837 analyses (PC1 vs PC2). PC1 vs PC3 and PC2 vs PC3 are shown in Fig. S4. (b) Comparison
838 between Ca, Fe and Ti content from XRF core-scanner and discrete XRF analyses in core
839 SK07.

840

841 **Figure 5:** Hyperspectral image and spectral indices (relative absorption band area or RABA)
842 for core SK06. From left to right: 500- μm resolution Br and S (XRF core-scanner),
843 distribution maps and plots of Chl *a* plus derivatives (RABA630–700), phycocyanin
844 (RABA600–630), Bacteriopheophytin (Bphe *a*) (RABA750–900) and aromatic organic
845 matter (ARO; RABA1660–1690), respectively. Grey shadings outline intervals with higher
846 Br, S, phycocyanin and Bphe *a* contents, which are tied to the occurrence of dark organic

847 sediments in core SK06. Yellow shadings outline intervals with elevated content in Chl *a* and
848 phycocyanin in core SK06.

849

850 **Figure 6:** Average changepoint model's slope calculated for PC1. The green curve in the
851 individual plots outlines the most pronounced regime shift identified around 5050 cal. yr BP.
852 Purple-filled curves correspond to abrupt shifts identified in both PC1 and PC2.

853

854 **Figure 7:** Average changepoint model's slope calculated for PC2. The green curve in the
855 individual plots outlines the most pronounced regime shift identified around 5250 cal. yr BP.
856 Purple-filled curves correspond to abrupt shifts identified in both PC1 and PC2.

857

858 **Figure 8:** Close-up on XRF and hyperspectral data for the interval 131.5–151.5 cm in core
859 SK06 (ca. 6500–5600 cal. yr BP), plotted against grain size (clay, sand), stable isotope ($\delta^{18}\text{O}$,
860 $\delta^{13}\text{C}_{\text{org}}$), total organic matter (TOC), total carbon (TC), total nitrogen (TN) and C/N data from
861 core SK07 in Lake Son Kol (Huang et al. 2014; unpublished data). Yellow shadings refer to
862 intervals characterized by dark organic layers with coarser minerogenic detritus, high Br, S,
863 Bphe *a*, ARO in core SK06, as well as high TOC (and TC, TN and C/N) and depleted $\delta^{18}\text{O}$
864 and $\delta^{13}\text{C}_{\text{org}}$ values in core SK07.

865

866 **Figure 9:** Time series of hyperspectral, geochemical and PCA data for Lake Son Kol (core
867 SK06), plotted with proxy data from core SONK_11_D1/2 from Lake Son Kol (Lauterbach et
868 al., 2014; Schwarz et al., 2017). XRF (9-point running means) and hyperspectral data (15-
869 point running means) for core SK06 are represented as standardized values (mean = 0,
870 standard deviation = 1). Yellow shadings outline periods for which hypolimnetic euxinia
871 prevailed in Lake Son Kol. See text for detail.

872

873 **Figure 10:** Comparison of geochemical and palynological proxies from Lake Son Kol to
874 other regional records in Central Asia. Shaded areas indicate periods characterized by
875 relatively warm summers and milder winters/springs with intensified precipitation (rain or
876 snow) at Lake Son Kol, favoring enhanced winter/spring snow thaw episodes; (a) *Bphe a*
877 record from Lake Qinghai, China (Ji et al., 2009); (b) alkenone-inferred temperature record at
878 Lake Qinghai (Hou et al., 2016); (c) speleothem $\delta^{18}\text{O}$ record from Tonnel'naya (Ton) Cave,
879 Uzbekistan (Chen et al., 2016); (d) speleothem $\delta^{18}\text{O}$ record from Baluk Cave, China (Liu et
880 al., 2019); (e) influx of coprophilous fungi *sensu stricto* (core SK06, this study); (f) arboreal
881 pollen (%) record from Lake Son Kol, Kyrgyzstan (Mathis et al., 2014); (g) total content of
882 Betulaceae (%) from Lake Son Kol, Kyrgyzstan (Mathis et al., 2014) and (h) *Bphe a* record
883 (15-point running means) from Lake Son Kol (core SK06, this study).

884

885 **Figure S1:** Core-to-core correlation between sediment cores SK06, SK04 and SK06 using
886 magnetic susceptibility.

887

888 **Figure S2:** Correlation of core SK06 (this study) with core SONK_11_D1/2 from Lauterbach
889 et al., (2014) using distinct lithological marker layers.

890

891 **Figure S3:** 500- μm resolution (grey curves) XRF core-scanner data exhibiting Br, S, Ti, Rb,
892 Zr, Fe, Si, Ca, Sr and Mn. Red curves represent 31-point running means. Yellow shadings
893 outline intervals with high Br, S, Ti, Rb and low Zr, Ca, Sr and Mn, which are tied to the
894 occurrence of dark organic sediments in core SK06.

895

896 **Figure S4:** PCA biplots of major elements obtained by XRF core scanner and hyperspectral
897 analyses (PC1 vs PC3 and PC2 vs PC3).

898

899 **Figure S5:** Average changepoint models of PC1 and PC2 during the last 8500 years in core
900 SK06.

901

902 **Figure S6:** Thin-section images (cross-polarized light) of the detrital layer microfacies types
903 observed in dark organic-rich sediments; (a) minerogenic detrital layer characterized by rare
904 normal graded beds in a non-graded matrix-supported (i.e., fine silt) layer. Note the absence
905 of a sharp erosive basal layer; (b) a typical detrital layer exhibiting mixed organic (e.g.,
906 phytoclasts), bioclasts (e.g., ostracod valves) and minerogenic detritus. Note the absence of
907 grading into the detrital layer; (c, d) organic detrital layer mainly consisting of phytoclasts,
908 interpreted as reworked terrestrial material (sheet-wash) from the catchment.

909

910 **Table 1:** AMS ^{14}C dates obtained on terrestrial plant remains and bulk sediment OM (with
911 unknown proportion of aquatic OM) in cores SK07 and SK04 (Huang et al., 2014) and core
912 SONK_11_D1/2 (Lauterbach et al., 2014). Published AMS ^{14}C dates were transposed to core
913 SK06 using lithological and magnetic susceptibility-inferred correlations (Figs. 2a, S1 and
914 S2). Conventional ^{14}C ages were calibrated using Oxcal 4.4 with the IntCal20 calibration
915 dataset (Reimer et al., 2020). A reservoir correction of 150 years was applied before
916 calibration to conventional ^{14}C ages. Only the samples with asterisks (*) were considered for
917 age modeling (see text for detail).

918

919 **References**

920

921 Abramowski U, Bergau A, Seebach D et al. (2006) Pleistocene glaciations of Central Asia:
922 results from ^{10}Be surface exposure ages of erratic boulders from the Pamir (Tajikistan) and
923 the Alay-Turkestan range (Kyrgyzstan). *Quat Sci Rev* 25:1080–1096
924
925 Academy of Science of the Kyrgyz SSR (1987) Atlas of the Kyrgyz Soviet Socialist Republic.
926 Volume 1: Natural conditions and resources. Moscow: State Agency for Cartography and
927 Geodesy, Central Directorate for Geodesy and Cartography, Council of Ministers of the
928 USSR
929
930 Adrian R, Deneke R, Mischke U et al. (1995) A long-term study of the Heiligensee (1975–
931 1992). Evidence for effects of climatic change on the dynamics of eutrophied lake
932 ecosystems. *Arch Hydrobiol* 133(3):315–337
933
934 Aichner B, Feakins SJ, Lee JE et al. (2015) High-resolution leaf wax carbon and hydrogen
935 isotopic record of the late Holocene paleoclimate in arid Central Asia. *Clim Past* 11:619–633
936
937 Aizen VB, Aizen EM, Melack JM et al. (1997) Climatic and hydrologic changes in the Tien
938 Shan, Central Asia. *Journal of Climate* 10:1393–1404
939
940 Alley RB, Mayewski PA, Sowers T et al. (1997) Holocene climate instability: a large event
941 8000–8400 years ago. *Geology* 25:482–486
942
943 Appleby PG, Richardson N, Nolan PJ (1991) ^{241}Am dating of lake sediments. *Hydrobiologia*
944 214:35–42
945

946 Balsam WL, Damuth JE, Schneider RR (1997) Comparison of shipboard vs. shorebased
947 spectral data from Amazon Fan cores: implications for interpreting sediment composition.
948 Ocean Drill Prog Sci 155:193–215
949

950 Balsam WL, Deaton BC (1996) Determining the composition of Late Quaternary marine
951 sediments from NUV, VIS, and NIR diffuse reflectance spectra. Marine Geol 134:31–55
952

953 Barranco FT, Balsam WL, Deaton BC (1989) Quantitative reassessment of brick red lutites—
954 evidence from reflectance spectrophotometry. Marine Geol 89:299–314
955

956 Bates CC (1953) Rational theory of delta formation: AAPG Bulletin 37(9):2119–2162
957

958 Beer R, Heiri O, Tinner W (2007) Vegetation history, fire history and lake development
959 recorded for 6300 years by pollen, charcoal, loss on ignition and chironomids at a small lake
960 in southern Kyrgyzstan (Alay Range, Central Asia). The Holocene 17:977–985
961

962 Beloglasova VN, Smirnova NB (1987). Atlas Kirgizskoj SSR. GUGK SSSR, Bishkek (in
963 Russian)
964

965 Blankenship RE (2013) Molecular mechanisms of photosynthesis. John Wiley & Sons.
966

967 Bugaev VA, Dzhordzhio VA, Kozik EM et al. (1957) Sinopticheskie Protsessy Sredney Azii
968 (Synoptic processes in Central Asia). Izd Akad Nauk Uzbek, SSR, Tashkent, 447 pp
969

970 Butz C, Grosjean M, Poraj-Górska A et al. (2016) Bacteriopheophytin *a* as an indicator of

971 meromixis in sediments of Lake Jaczno, north-east Poland, CE 1891–2010. *Glob Planet*
972 *Change* 144:109–118
973
974 Butz C, Grosjean M, Fischer D et al. (2015) Hyperspectral imaging spectroscopy: a promising
975 method for the biogeochemical analysis of lake sediments. *J Appl Remote Sens* 9:1–20
976
977 Cai Y, Chiang JCH, Breitenbach SFM et al. (2017) Holocene moisture changes in western
978 China, Central Asia, inferred from stalagmites. *Quat Sci Rev* 158:15–28
979
980 Chapron E, Arnaud F, Noel H et al. (2005) Rhône River flood deposits in Lake Le Bourget: a
981 proxy for Holocene environmental changes in the NW Alps, France. *Boreas* 31:1–13
982
983 Chapron E, Van Rensbergen P, De Batist M et al. (2004) Fluid-escape features as a precursor
984 of a large sublacustrine sediment slide in Lake Le Bourget, NWAlps, France. *Terra Nova*
985 16:305–311
986
987 Chen Y, Li W, Deng H et al. (2016) Changes in Central Asia’s water tower: Past, present and
988 future. *Scientific Reports* 6, <https://doi.org/10.1038/srep35458>
989
990 Chen FH, Chen JH, Holmes JA et al. (2010) Moisture changes over the last millennium in the
991 arid central Asia: A review, synthesis and comparison with monsoon region. *Quat Sci Rev*
992 29:1055–1068
993
994 Chen FH, Yu ZC, Yang ML et al. (2008) Holocene moisture evolution in arid central Asia
995 and its out-of-phase relationship with Asian monsoon history. *Quat Sci Rev* 27:351–364

996

997 Christensen JH et al (2017) Regional Climate Projections. *Climate Change 2007: the Physical*
998 *basis*, 847–940

999

1000 Coolen MJL, Overmann J (1998) Analysis of subfossil molecular remains of purple sulphur
1001 bacteria in a lake sediment. *App Env Microbiol* 64(11):4513–4521

1002

1003 Cuddington K, Leavitt PR (1999) An individual-based mode of pigment flux in lakes:
1004 implications for organic biogeochemistry and paleoecology. *Can J Fish Aqua Sci* 56(10):
1005 1964–1977

1006

1007 Cugny C, Mazier F, Galop D (2010) Modern and fossil non-pollen palynomorphs from the
1008 Basque mountains (western Pyrenees, France): the use of coprophilous fungi to reconstruct
1009 pastoral activity. *Veget Hist Archaeobot* 19:391–408

1010

1011 Cuven S, Francus P, Lamoureux SF (2010) Estimation of grain size variability with micro X-
1012 ray fluorescence in laminated lacustrine sediments, Cape Bounty, Canadian High Arctic. *J*
1013 *Paleolimnol*: doi10.1007/s10933-010-9453-1

1014

1015 Das B, Vinebrooke RD, Sanchez-Azofeifa A et al. (2005) Inferring lake sediment chlorophyll
1016 concentrations with reflectance spectroscopy: a novel approach to
1017 reconstructing trophic status changes in alpine lakes. *Can J Fish Aquat Sci* 62:1067–1078

1018

1019 Davies SJ, Lamb HF, Roberts S (2015) Micro-XRF core scanning in palaeolimnology: recent
1020 developments. In: Croudace I, Rothwell R (eds) Micro-XRF Studies of Sediment Cores.
1021 Developments in Paleoenvironmental Research, vol 17. Springer, Dordrecht
1022

1023 Davies OK, Schafer D (2006) *Sporormiella* fungal spores, a palynological means of detecting
1024 herbivore density. *Palaeogeog, Palaeoclimato., Palaeoecol.* 237:40–50
1025

1026 Davison W (1993) Iron and Manganese in lakes. *Earth Sci Rev* 34(2):119–163
1027

1028 Dressler M, Hübener T, Görs S et al. (2007) Multi-proxy reconstruction of trophic state,
1029 hypolimnetic anoxia and phototrophic sulphur bacteria abundance in a dimictic lake in
1030 Northern Germany over the past 80 years. *J Paleolimnol* 37:205–219
1031

1032 Etienne D, Wilhelm B, Sabatier P et al. (2013) Influence of sample location and livestock
1033 numbers on *Sporormiella* concentrations and accumulation rates in surface sediments of Lake
1034 Allos, French Alps. *J Paleolimnol* 49:117–127
1035

1036 Faegri K, Iversen J (1989) Textbook of pollen analysis. John Wiley and Sons, Chichester-
1037 New York-Brisbane-Toronto-Singapore.
1038

1039 Feeser I, O’Connell M (2009) Fresh insights into long-term changes in flora, vegetation, land
1040 use and soil erosion in the karstic environment of the Burren, western Ireland. *Journal of*
1041 *Ecology* 97(5):1083–1100
1042

1043 Foley B, Jones ID, Maberly SC et al. (2012) Long term changes in oxygen depletion in a

1044 small temperate lake: effects of climate change and eutrophication. *Freshw Biol* 57(2):278–
1045 289
1046
1047 Gallagher K, Bodin T, Sambridge M, Weiss D, Kylander M, Large D (2011) Inference of
1048 abrupt changes in noisy geochemical records using transdimensional change point models.
1049 *Earth and Planetary Science Letters* 311 :182–194
1050
1051 Gauthier E, Bichet V, Massa C, Petit C et al. (2010) Pollen and non-pollen palynomorph
1052 evidence of medieval farming activities in southwestern Greenland. *Veget Hist Archaeobot*
1053 19:427–438
1054
1055 Gelorini V, Verbeken A, van Geel B et al. (2011) Modern non-pollen palynomorphs from
1056 East African lake sediments. *Rev Palaeobot Palynol* 164(3–4):143–173
1057
1058 Giguet-Covex C, Arnaud F, Poulenard J et al. (2011) Changes in erosion patterns during the
1059 Holocene in a currently treeless subalpine catchment inferred from lake sediment
1060 geochemistry (Lake Anterne, 2063 m.a.s.l, NW France): the role of climate and human
1061 activities. *The Holocene* 21:651–665
1062
1063 Glazer AN, Fang S, Brown DM (1973) Spectroscopic properties of C-Phycocyanin and of its
1064 α and β subunits. *The Journal of Biological Chemistry* 248(16):5679–5685
1065
1066 Håkanson L, Jansson M (1983) Principles of lake sedimentology. Heidelberg, Springer
1067 Verlag, 312 pp
1068

1069 Hammer Ø, Harper DAT, Ryan PD (2001) PAST: Paleontological Statistics Software
1070 Package for Education and Data Analysis. *Palaeontologia Electronica* 4(1), 9p.
1071
1072 Heinecke L, Fletcher WJ, Mischke S et al. (2018) Vegetation change in the eastern Pamir
1073 Mountains, Tajikistan, inferred from Lake Karakul pollen spectra of the last 28 kyr.
1074 *Palaeogeogr Palaeoclimatol Palaeocol* 511(15):232–242
1075
1076 Heinecke L, Mischke S, Adler K et al. (2017) Climatic and limnological changes at Lake
1077 Karakul (Tajikistan) during the last ~29 cal ka. *J Paleolimnol* 58:317–334
1078
1079 Heinecke L, Mischke S, Adler K et al. (2016) Late Pleistocene to Holocene climate and
1080 limnological changes at Lake Karakul (Pamir Mountains, Tajikistan). *Clim Past Discuss*,
1081 doi:10.5194/cp-2016-34
1082
1083 Heymann C, Nelle O, Dörfler W et al. (2013) Late Glacial to Mid-Holocene palaeoclimates
1084 development of Southern Greece inferred from the sediment sequence of Lake Stymphalia
1085 (NE Peloponnese). *Quat Int* 302:42–60
1086
1087 Hodgson D, Wright S, Tyler P et al. (1998) Analysis of fossil pigments from algae and
1088 bacteria in meromictic Lake Fidler, Tasmania, and its application to lake management. *J*
1089 *Paleolimnol* 19(1):1–22
1090
1091 Horváth H, Kovacs AW, Riddick C et al. (2013) Extraction methods for phycocyanin
1092 determination in freshwater filamentous cyanobacteria and their application in a shallow lake.
1093 *Eur J Phycol* 48(3):278–286

1094

1095 Hou J, Huang Y, Zhao J et al. (2016) Large Holocene summer temperature oscillations and
1096 impact on the peopling of the northeastern Tibetan Plateau. *Geophys res Lett* 43 :1323–1330
1097

1098 Huang X, Oberhänsli H, von Suchodoletz H et al. (2014) Hydrological changes in western
1099 Central Asia (Kyrgyzstan) during the Holocene: Results of a paleolimnological study from
1100 Son Kul. *Quat Sci Rev* 103:134–152
1101

1102 IPCC (2013) *Climate Change 2013: The Physical Science Basis. Contribution of Working*
1103 *Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.*
1104 Cambridge; New York: Cambridge University Press, 1535 pp
1105

1106 Jacq K, Martinez-Lamas R, Van Exem A, Debret M (2020) Hyperspectral core-logger image
1107 acquisition. *Protocols* io, [https://protocols.io/view/hyperspectral-core-logger-image-](https://protocols.io/view/hyperspectral-core-logger-image-acquisition-bikckcsw)
1108 [acquisition-bikckcsw](https://protocols.io/view/hyperspectral-core-logger-image-acquisition-bikckcsw)
1109

1110 Jenny JP, Arnaud F, Dorioz JM et al. (2013) A spatiotemporal investigation of varved
1111 sediments highlights the dynamics of hypolimnetic hypoxia in a large hard-water lake over
1112 the last 150 years. *Limnol Oceanogr* 58(4):1395–1408
1113

1114 Ji J, Balsam W, Shen J et al. (2009) Centennial blooming of anoxygenic phototrophic bacteria
1115 in Qinghai Lake linked to solar and monsoon activities during the last 18,000 years. *Quat Sci*
1116 *Rev* 28:1304–1308
1117

1118 Kalff J (2002) *Limnology: Inland Water Ecosystems.* Prentice Hall, 592 pp

1119

1120 Koppes M, Gillespie AR, Burke RM et al. (2008) Late Quaternary glaciation in the Kyrgyz
1121 Tien Shan. *Quat Sci Rev* 27:846–866

1122

1123 Laug A, Schwarz A, Lauterbach S et al. (2020) Ecosystem shifts at two Mid-Holocene tipping
1124 points in the alpine Lake Son Kol (Kyrgyzstan, Central Asia). *The Holocene*,
1125 doi:10.1177/0959683620932973

1126

1127 Lauterbach S, Witt R, Plessen B et al. (2014) Climatic imprint of the mid-latitude Westerlies
1128 in the Central Tien Shan of Kyrgyzstan and teleconnections to North Atlantic climate
1129 variability during the last 6000 years. *The Holocene* 24(8):970–984

1130

1131 Lenton TM (2011) Early warming of climate tipping points. *Nat Clim Change* 1:201–209

1132

1133 Lenton TM, Held H, Kriegler E et al. (2008) Tipping elements in the Earth’s climate system.
1134 *Proc Nat Acad Sci USA* 105:1786–1793

1135

1136 Li X, Zhao K, Dodson J et al. (2011) Moisture dynamics in central Asia for the last 15 kyr:
1137 New evidence from Yili Valley, Xinjiang, NW China. *Quat Sci Rev* 30:3457–3466

1138

1139 Liu, X, Rao, Z, Shen CC et al. (2019) Holocene solar activity imprint on centennial- to
1140 multidecadal-scale hydroclimatic oscillations in Arid Central Asia. *JGR Atmospheres*
1141 124:2562–2573

1142

1143 Livingstone DM (2003) Impact of secular climate change on the thermal structure of a large

1144 temperate central European lake. *Clim Change* 57(1–2):205–225

1145

1146 Machalett B, Oches EA, Frechen M et al. (2008) Aeolian dust dynamics in Central Asia
1147 during the Pleistocene: driven by the long-term migration, seasonality and permanency of the
1148 Asiatic polar front. *Geochemistry, Geophysics, Geosystems* 9:Q08Q09

1149

1150 Machalett B, Frechen B, Hambach U et al (2006) The loess sequence from Remisowka
1151 (northern boundary of the Tien Shan Mountains (Kazakhstan). Part I: luminescence dating.
1152 *Quat Int* 152–153:192–201

1153

1154 Madigan MT (2003) Anoxygenic phototrophic bacteria from extreme environments.
1155 *Photosynth Res* 76:157–171

1156

1157 Mathis M, Sorrel P, Klotz S et al. (2014) Regional vegetation patterns at lake Son Kul reveal
1158 Holocene climatic variability in central Tien Shan (Kyrgyzstan, Central Asia). *Quat Sci Rev*
1159 89:169–185

1160

1161 McGowan S, Juhler RK, Anderson NJ (2008) Autotrophic response to lake age, conductivity
1162 and temperature in two West Greenland lakes. *J Paleolimnol* 39:301–317

1163

1164 Michelutti N, Blais JM, Cumming BF et al. (2010) Do spectrally inferred determinations of
1165 chlorophyll *a* reflect trends in lake trophic status? *J Paleolimnol* 43:205–217

1166

1167 Mischke S, Rajabov I, Mustaeva N et al. (2010) Modern hydrology and late Holocene history
1168 of Lake Karakul, eastern Pamirs (Tajikistan): A reconnaissance study. *Palaeogeogr*

1169 Palaeoclimatol Palaeocol 289:10–24
1170
1171 Mulder T, Chapron E (2011) Flood deposits in continental and marine environments:
1172 Character and significance, in R. M. Slatt and C. Zavala (eds) Sediment transfer from shelf to
1173 deep water–Revisiting the delivery system: AAPG Studies in Geology 61:1–30
1174
1175 Mur LR, Skulberg OM, Utkilen H (1999) Cyanobacteria in the environment. In: Chorus I,
1176 Bartram J (eds) Toxic cyanobacteria in water: a guide to their public health consequences,
1177 monitoring and management. E & FN Spon, London
1178
1179 Narama C, Kondo R, Tsukamoto S et al. (2007) OSL dating of glacial deposits during the last
1180 glacial in the Terskey-Alatoo range, Kyrgyz Republic. Quat Geochronol 2:249–254
1181
1182 Narama C, Okuno M (2006) Record of a glacier variations during the last glacial in the
1183 Turkestan Range of Pamir-Alay, Kyrgyz Republic. Ann Glaciol 43:397–404
1184
1185 Oren A (2011) Characterization of pigments of prokaryotes and their use in taxonomy and
1186 classification. In : Methods in Microbiology, Academic Press, pp. 261–282.
1187
1188 Overmann J, Beatty JT, Hall KJ et al. (1991) Characterization of a dense, purple sulfur
1189 bacteria layer in a meromictic salt lake. Limnol Oceanogr 36(5):846–859
1190
1191 Ozturk T, Altinsoy H, Türkes M et al. (2012) Simulation of temperature and precipitation
1192 climatology for the Central Asia CORDEX domain using RegCM 4.0. Climate Research
1193 52:63–76

1194

1195 Pacton M, Sorrel P, Bevillard B et al. (2015) Sedimentary facies analyses from nano- to
1196 millimetre scale exploring past microbial activity in a high-altitude lake (Lake Son Kul,
1197 Central Asia). *Geol Mag* 152(5):902–922

1198

1199 Parkin T, Brock T (1980) Photosynthetic bacterial production in lakes: the effects of light
1200 intensity. *Limnol Oceanogr* 25(4):711–718

1201

1202 Ramsey CB (2008) Deposition models for chronological records. *Quat Sci Rev* 27:42–60

1203

1204 Ramsey CB (2001) Development of the radiocarbon calibration program. *Radiocarbon*
1205 43:355–363

1206

1207 Ramsey CB (1995) Radiocarbon calibration and analysis of stratigraphy: The OxCal program.
1208 *Radiocarbon* 37:425–430

1209

1210 Randsalu-Wendrup L, Conley D, Carstensen J et al. (2014) Combining limnology and
1211 palaeolimnology to investigate recent regime shifts in a shallow, eutrophic lake. *J Paleolimnol*
1212 51:437–448

1213

1214 Reimer PJ, Austin EN, Bard E et al. (2020) The IntCal20 Northern Hemisphere radiocarbon
1215 age calibration curve (0–55 cal kBP). *Radiocarbon* 62(4):725–757

1216

1217 Rein B, Sirocko F (2002) In-situ reflectance spectroscopy— analysing techniques for high-
1218 resolution pigment logging in sediment cores. *Int J Earth Sci* 91:950–954

1219

1220 Rogozin DY, Zykov VV, Degetmendzhi AG (2012) Ecology of purple sulfur bacteria in the
1221 highly stratified meromictic Lake Shunet (Siberia, Khakassia) in 2002–2009. *Microbiology*
1222 81(6) :727–735

1223

1224 Rohling EJ, Pälike H (2005) Centennial-scale climate cooling with a sudden cold event
1225 around 8,200 years ago. *Nature* 434:975–979

1226

1227 Rousseau M, Demory F, Miramont C et al. (2020) Palaeoenvironmental change and glacier
1228 fluctuations in the high Tien Shan Mountains during the last millennium based on sediments
1229 from Lake Ala Kol, Kyrgyzstan. *Pal Pal Pal* 558, 109987.

1230

1231 Savoskul OS, Solomina ON (1996) Late Holocene glacier variations in the frontal and inner
1232 ranges of the Tien Shan, central Asia. *The Holocene* 6(1):25–35

1233

1234 Scheer H (2006) An overview of chlorophylls and bacteriochlorophylls : biochemistry,
1235 biophysics, functions and applications. In : Grimm B, Porra R, Rüdiger W, Scheer H (Eds),
1236 *Chlorophylls and bacteriochlorophylls Advances in Photosynthesis and Respiration*. Springer,
1237 Netherlands:1–26

1238

1239 Scheffer M, Carpenter SR, Lenton TM et al. (2012) Anticipating critical transitions. *Science*
1240 338:344–348

1241

1242 Schmidt R, Psenner R, Müller J et al. (2002) Impact of late glacial climate variations on
1243 stratification and trophic state of the meromictic lake Längsee (Austria): validation of a

1244 conceptual model by multi-proxy studies. *J Limnol* 61(1):49–60
1245
1246 Schröder HG, Wessels M, Niessen F (1998) Acoustic facies and depositional structures of
1247 Lake Constance. *Archiv fur Hydrobiologie Special Issues Advance Limnology* 53:351–368
1248
1249 Schroeter N, Lauterbach S, Stebich M et al. (2020) Biomolecular evidence of early human
1250 occupation of a high-altitude site in Western Central Asia during the Holocene. *Front Earth*
1251 *Sci* 8, doi:10.3389/feart.2020.00020
1252
1253 Schwarz A, Turner F, Lauterbach S et al. (2017) Mid- to Late Holocene climate-driven
1254 regime shifts inferred from diatom, ostracod and stable isotope records from Lake Son Kol
1255 (Central Tien Shan, Kyrgyzstan). *Quat Sci Rev* 177:340–356
1256
1257 Shnitnikov AV (1980) *Ozera Tian-Shanya i ikh istoriya* [Lakes of the Tian Shan and their
1258 history]. Leningrad: Nauka.
1259
1260 Solomon S, Qin D, Manning M et al. (eds) 2007 *Climate Change 2007: The Physical Science*
1261 *Basis (Contribution of Working Group I to the Fourth Assessment Report of the*
1262 *Intergovernmental Panel on Climate Change)*. Cambridge: Cambridge University Press
1263
1264 Sommaruga R, Psenner R (1997) Ultraviolet radiation in a high mountain lake of the Austrian
1265 Alps: Air and underwater measurements. *Photochemistry and Photobiology* 65:957–963
1266
1267 Stockmarr J (1971) Tablets with spores used in absolute pollen analysis. *Pollen spores*
1268 13:615–621

1269

1270 Støren EN, Dahl SO, Nesje A et al. (2010) Identifying the sedimentary imprint of high-
1271 frequency Holocene river floods in lake sediments: development and application of a new
1272 method. *Quat Sci Rev* 29:3021–3033

1273

1274 Trachsel M, Grosjean M, Schnyder D et al. (2010) Scanning reflectance spectroscopy (380–
1275 730 nm): a novel method for quantitative high-resolution climate reconstructions from
1276 minerogenic lake sediments. *J Paleolimnol* 44:979–994

1277

1278 van Geel B, Andersen ST (1988) Fossil ascospores of the parasitic fungus *Ustilina deusta* in
1279 Eemian deposits in Denmark. *Rev Palaeobot Palynol* 56:89–93

1280

1281 van Geel B, Aptroot A (2006) Fossil ascomycetes in Quaternary deposits. *Nova Hedwigia*
1282 82:313–329

1283

1284 van Geel B, Buuman J, Brinkkemper O et al. (2003) Environmental reconstruction of a
1285 Roman Period settlement site in Uitgeest (The Netherlands), with special reference to
1286 coprophilous fungi. *J Archaeol Sci* 30:873–883

1287

1288 Van Gemerden H, Mas J (1995) Ecology of phototrophic bacteria. In : Blankenship R,
1289 Madigan M, Bauer C (Eds) *Anoxygenic photosynthetic bacteria. Advances in photosynthesis*
1290 *and respiration*. Springer, Netherlands : pp 49–85

1291

1292 Van Rensbergen P, De Batist M, Beck C et al. (1999) High-resolution seismic stratigraphy of
1293 glacial to interglacial fill of a deep glacigenic lake: Lake Le Bourget, Northwestern Alps,

1294 France. *Sed Geol* 128:99–129

1295

1296 Van Rensbergen P, De Batist M, Beck C et al. (1998) High-resolution seismic stratigraphy of
1297 late Quaternary fill of Lake Annecy (Northwestern Alps): evolution from glacial to
1298 interglacial sedimentary processes. *Sed Geol* 117:71–96

1299

1300 Wheeler OH (1959) Near infrared spectra of organic compounds. *Chem Rev* 59:629–666

1301

1302 Williams RC, Gingrich JC, Glazer AN (1980) Cyanobacterial phycobilisomes. *J Cell Biol*
1303 85:558–566

1304

1305 Wirth S, Gili A, Niemann H et al. (2013) Combining sedimentological, trace metal (Mn, Mo)
1306 and molecular evidence for reconstructing past water-column redox conditions: The example
1307 of meromictic Lake Cadagno (Swiss Alps). *Geoch Cosmoch Acta* 120:220–238

1308

1309 Wolfe AP, Vinebrooke RD, Michelutti N et al. (2006) Experimental calibration of lake-
1310 sediment spectral reflectance to Chlorophyll a concentrations: methodology and
1311 paleolimnological validation. *J Paleolimnol* 36:91–100

1312

1313 Wolff C, Plessen B, Dudashvili AS et al. (2017) Precipitation evolution of Central Asia
1314 during the last 5000 years. *The Holocene*, DOI: 10.1177/0959683616652711

1315

1316 Woźniak M, Bradtke K, Darecki M et al. (2016) Empirical model for phycocyanin
1317 concentration estimation as an indicator of cyanobacterial bloom in the optically complex
1318 coastal waters of the Baltic Sea. *Remote Sens* 8(3): 212, <https://doi.org/10.3390/rs8030212>

1319

1320 Yacobi YZ, Köhler J, Leunert F et al. (2015) Phycocyanin-specific absorption coefficient:

1321 eliminating the effect of chlorophyll absorption. *Limnol and Oceanogr Methods* 13(4):157–

1322 1168

1323

1324 Zhao J, An C-B, Huang Y et al. (2017) Contrasting early Holocene temperature variations


1325 between monsoonal East Asia and westerly dominated Central Asia. *Quat Sci Rev* 178:14–23

1326

1327 Zyakun AM, Lunina ON, Prusakov TS et al. (2009) Fractionation of stable carbon isotopes by

1328 photoautotrophically growing anoxygenic purple and green sulfur bacteria. *Microbiology*

1329 78(6):757–768






Figure 2




Figure 3





Figure 4




Figure 5





Figure 7




Figure 8




Figure 9




Figure 10

Lab. ID	Sampling interval	Dated material	Correlation LITHO in Core SK6 (cm)	AMS ¹⁴ C age (yr BP + σ)	Corr. AMS ¹⁴ C age (yr BP + σ)	Calibrated age (cal. yr BP + 2 σ)
	SONK_11_D1/2 (Lauterbach et al., 2014)	²⁴¹ Am peak (5,5 cm)	2			-13
Poz-26363	Sonkul 4A 0-75 0-1cm	Bulk sediment	30	2000 ± 30	1850 ± 30	2139**
*Poz-44073	SONK_11_D1/2 (Lauterbach et al., 2014)	Terrestrial plant remains	33	1575 ± 35		
*Poz-26364	Sonkul 4A 0-75 30-31cm	Bulk sediment	63,5	2770 ± 40	2620 ± 40	
*Poz-26366	Sonkul 4A 75-150 0-1cm	Bulk sediment	100	3750 ± 35	3600 ± 35	
*Poz-26367	Sonkul 4A 75-150 20-21cm	Bulk sediment	127	4730 ± 40	4580 ± 40	
*Poz-26371	Sonkul 7A 120-180 0-1cm	Bulk sediment	135	5270 ± 40	5120 ± 40	
Poz-26368	Sonkul 4A 75-150 40-41cm	Bulk sediment	137	5240 ± 40	5090 ± 40	5817**
*Poz-26375	Sonkul 7A 120-180 5cm	Bulk sediment	139	5250 ± 40	5100 ± 40	
Poz-37470	Sonkul 7A 120-180 cm 7.2 cm- 8.6cm	Bulk sediment	141	5410 ± 40	5260 ± 40	6052**
*Poz-37469	Sonkul 7A 120-180 cm 7.2 cm- 8.6cm	Terrestrial plant remains	141	5460 ± 40	5310 ± 40	
Poz-26372	Sonkul 7A 120-180 15.5-16.5cm	Bulk sediment	149	6200 ± 50	6050 ± 50	6900**
Poz-26370	Sonkul 4A 75-150 65-66cm	Bulk sediment	167?	6790 ± 40	6640 ± 40	7521**
*Poz-26373	Sonkul 7A 120-180 26.5-27.5cm	Bulk sediment	166	6330 ± 50	6180 ± 50	
*Poz-37471	Sonkul 7A 120-180 cm 37.5-38.5cm	Bulk sediment	172	6880 ± 50	6730 ± 50	
*Poz-26374	Sonkul 7A 120-180 45.5-46.3cm	Bulk sediment	184	7570 ± 50	7420 ± 50	

The asterisks (*) denotes the radiocarbon dates applied to establish the age modelling. Bold samples correspond to excluded dates from age-dep
Double asteriks (**) correspond to median probability ages as calculated online with Oxcal8.2