



HAL
open science

Editorial: The Oral Microbiome Is a Key Factor in Oral and Systemic Health

Denis Bourgeois, Lucio Souza Gonçalves, Josué da Costa Lima-Junior,
Florence Carrouel

► **To cite this version:**

Denis Bourgeois, Lucio Souza Gonçalves, Josué da Costa Lima-Junior, Florence Carrouel. Editorial: The Oral Microbiome Is a Key Factor in Oral and Systemic Health. *Frontiers in Microbiology*, 2022, 10.3389/fmicb.2022.855668 . hal-03571799

HAL Id: hal-03571799

<https://univ-lyon1.hal.science/hal-03571799v1>

Submitted on 14 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Editorial: The Oral Microbiome Is a Key Factor in Oral and Systemic Health

Denis Bourgeois¹, Lucio Souza Gonçalves², Josué da Costa Lima-Junior^{2,3} and Florence Carrouel^{1*}

¹ Health, Systemic, Process, UR4129 Research Unit, University Claude Bernard Lyon 1, University of Lyon, Lyon, France, ² Laboratory of Molecular Biology and Immunology, Postgraduate Program in Dentistry, Faculty of Dentistry, Estácio de Sá University, Rio de Janeiro, Brazil, ³ Immunoparasitology Laboratory, Oswaldo Cruz Institute, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil

Keywords: inflammation, saliva, microbiota, virus, biomarker, periodontitis, chronic disease, cancer

Editorial on the Research Topic

The Oral Microbiome Is a Key Factor in Oral and Systemic Health

The oral cavity is a dynamic ecosystem comprising an assemblage of microbial communities, including many pathogenic or opportunistic species (Proctor and Relman, 2017). After the gut microbiome, the human oral microbiome (HOM) is the largest microbial community in the human body. The HOM plays a role in the onset and progression of several localized and systemic diseases including those of bacterial, viral and fungal origin (Soffritti et al.). In this Special Issue, there is a consensus: the oral microbiome is a key factor in oral and systemic health. Also, considering HOM as biomarkers for diseases is a significant emerging orientation. Saliva, the cornerstone of the HOM and systemic health relationship, has gained popularity as a readily available source of biomarkers useful for diagnosing specific oral and systemic conditions (Ferrari et al., 2021).

In oral health, biomarkers in saliva (e.g., enzymes, antibodies, protein markers, or oxidative stress markers) can be used for activity determination and for periodontal disease prognosis (Podzimek et al., 2016). The presence of key pathobionts and ongoing gingival inflammation are critical to the progression of periodontal disease. Mediators of periodontal disease initiation, progression, and recurrence are related to dysbiosis within the subgingival biofilm microbial community and the host immune response generated (Martínez et al., 2021). Abnormal changes in bacterial correlations, community structures, and local stability are linked to the dysbiosis observed in periodontal or peri-implant disease (Zhang et al.). Current findings indicate that salivary markers of oxidative stress are indicative of other clinical disease indices such as the papillary bleeding index and the caries index. Fungal species *Candida dubliniensis* and *Candida tropicalis* more abundant in the saliva of children with severe early childhood caries should play a role as caries risk markers (de Jesus et al.).

Recent studies have also demonstrated the role of periodontal disease as a risk factor or potentiator of distant systemic pathologies such as diabetes, inflammatory bowel diseases, Alzheimer's disease and oral cancer, further highlighting the importance of the oral cavity in systemic health (Kapila, 2021). This allowed for further characterization of oral microbial dysbiosis, especially putative bacterial periodontopathogens and changes in the composition of the oral virome during disease. New microorganisms (viruses, phages and bacteria) have recently been identified for their role in disease progression (Sedghi et al., 2021).

OPEN ACCESS

Edited and Reviewed by:

M. Pilar Francino,
Fundación para el Fomento de la
Investigación Sanitaria y Biomédica de
la Comunitat Valenciana
(FISABIO), Spain

*Correspondence:

Florence Carrouel
florence.carrouel@univ.lyon1.fr

Specialty section:

This article was submitted to
Microbial Symbioses,
a section of the journal
Frontiers in Microbiology

Received: 15 January 2022

Accepted: 24 January 2022

Published: 14 February 2022

Citation:

Bourgeois D, Gonçalves LS,
Lima-Junior JdC and Carrouel F
(2022) Editorial: The Oral Microbiome
Is a Key Factor in Oral and Systemic
Health. *Front. Microbiol.* 13:855668.
doi: 10.3389/fmicb.2022.855668

Thus, oral bacteria could be biomarkers for diseases such as specific cancer types. Some specific bacteria such as *Capnocytophaga gingivalis*, *Peptostreptococcus sp.*, *Porphyromonas gingivalis*, *Prevotella sp.* and *Streptococcus sp.* are strongly associated with oral cancer (Karpiński, 2019). Bacteria belonging to genera *Actinomyces*, *Clostridium*, *Enterobacteriaceae*, *Fusobacterium*, *Haemophilus*, and *Veillonella* are linked to epithelial precursor lesions and oral cancer (La Rosa et al., 2020). Oral bacteria are detected in tumors outside the oral cavity. Every day, about 10^{11} bacteria from the oral cavity migrate into the lower gastrointestinal tract (DeClercq et al., 2021). Microorganisms act as chemical converters and metabolize nutrients from the host and from the diet (Anand et al., 2016). HOM can generate an ectopic colonization and produce numerous microbial metabolites capable of promoting tumorigenesis through the modulation of pathways related to energy homeostasis, immunological balance and nutritional intake (Zhang et al., 2016).

Oral pathobionts are essential in the development of colorectal and pancreatic cancer with current evidence showing differences in oral microbiota composition between patients with and without digestive cancers (Reitano et al., 2021). In cases of colorectal cancers, two periopathogenic species in particular have been frequently mentioned: *Fusobacterium nucleatum* and *Porphyromonas gingivalis*. In pancreatic cancers, in addition to previously mentioned bacteria, strains of *Aggregatibacter actinomycetemcomitans*, *Neisseria elongata*, and *Streptococcus mitis* have been described (Fan et al., 2018). Oral *Prevotella* species play an important role as commensals in health but can also be involved in diseases of the lower airways and upper gastrointestinal tract (Könönen and Gursoy). Oral bacteria from genera *Capnocytophaga* and *Veillonella* are apparently present in increased amounts in lung cancer patients (Yan et al., 2015). The study of complex interaction between the oral and gut microbiome in the pathogenesis of type 1 diabetes is advanced, and suggests the use of saliva microbiome composition for early diagnosis (Moskovitz et al.). Also, bacterial dysbiosis plays an important role in the esophageal carcinogenesis process through microbial metabolism, inflammation and genotoxicity (Dan et al.) and *Capnocytophaga* and *Veillonella* are reportedly present in increased amounts in lung cancer patients (Najafi et al., 2021). At least, a predictive salivary microbiome signature is associated with a high risk of developing cardiovascular diseases (Murugesan et al.).

It is important for prevention of viral infection to draw a perspective on the role of the oral cavity in the virus infection (Tada and Senpuku, 2021). Salivary markers for viral infections involve direct detection of specific viral antigens, such as proteins and nucleic acids or host antibodies to viral infections and may provide a high accuracy point-of-care platform for detection of viral infections. HOM dysbiosis may facilitate inflammation and virus replication, limiting the development of a protective IgA response (Soffritti et al.). Immunity in saliva is, in particular, thought to have considerable impacts on the incidence and progression of respiratory viral infection. Parts of

antiviral mechanisms against influenza virus and SARS-CoV-2 by immunity in saliva are similar.

Saliva based biomarkers are useful in diagnosis of several viral infections such as hepatitis A virus, hepatitis B virus, hepatitis C virus, Human immunodeficiency virus (HIV) 1, etc. (Zhang et al., 2016). Of course, several viruses have previously been isolated from saliva such as cytomegalovirus, Ebola virus, human herpes virus, herpes simplex virus, Influenza virus A, etc. (Corstjens et al., 2016). More recently, Zika and SARS-CoV-2 have been identified (To et al., 2020). In people living with HIV saliva might be used as a diagnostic tool for antioxidant changes in the future (Amjad et al., 2019). Oral bacterial species (e.g., *Leptotrichia spp.*), possessing unique niches and invasive properties, coexist with Human Papilloma Virus (HPV) within HPV-induced oral lesions in head and neck cancer patients (Mougeot et al.).

The fact that oral cavity is an important site for SARS-CoV-2 infection implicates saliva as a potential route of SARS-CoV-2 transmission (Carrouel et al.). The susceptibility of each individual to SARS-CoV-2 infection could therefore be characterized by HOM profile, which could facilitate virus replication and inflammation or conversely induce a protective IgA response (Soffritti et al.). Several routes of SARS-CoV-2 viral entry into the saliva have been suggested. There is direct entry to the oral cavity from upper and lower respiratory tract secretions, while circulatory viruses in the blood enter the gingival crevicular fluid. Studies reported a high yield of virus particles in the gingival sulcus and crevicular fluid, which are suspected to provide favorable conditions for virus replication and maintenance (Sri Santosh et al., 2020). Informations from the 2019 coronavirus pandemic highlight the link between oral and systemic health in a setting of viremia/bacteremia/microbemia, systemic inflammation, and/or immune system disruption in a susceptible host (Martínez et al., 2021).

Overall, the 12 contributions that make up this Special Issue highlight the fundamental relationship between HOM and systemic health providing potential microbiome-based clinical applications improving prevention, diagnosis, or drug response, which is of great significance. Characterization of microbial biomarkers is of great interest for precision medicine and represents a simple method to transfer microbiome research into clinical practice (Gilbert et al., 2018). If during the last years, an extraordinary effort has been made to identify biomarkers, today, HOM investigations have reached a critical inflection point. With the deepening understanding of the association between the HOM and other human microbiomes certain pathogens may be utilized as potential diagnostic biomarkers, screening tools, and prognostic indicators and interventions related to an altered human microbial composition may become the new adjuvant treatment in oral and systemic health.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

REFERENCES

- Amjad, S. V., Davoodi, P., Goodarzi, M. T., Abdolsamadi, H., Poorolajal, J., Parsa, S., et al. (2019). Salivary antioxidant and oxidative stress marker levels in HIV-positive individuals. *Comb. Chem. High Throughput Screen* 22, 59–64. doi: 10.2174/1386207322666190306144629
- Anand, S., Kaur, H., and Mande, S. S. (2016). Comparative *in silico* analysis of butyrate production pathways in gut commensals and pathogens. *Front. Microbiol.* 7:1945. doi: 10.3389/fmicb.2016.01945
- Corstjens, P. L. A. M., Abrams, W. R., and Malamud, D. (2016). Saliva and viral infections. *Periodontology* 70, 93–110. doi: 10.1111/prd.12112
- DeClercq, V., Nearing, J. T., and Langille, M. G. I. (2021). Investigation of the impact of commonly used medications on the oral microbiome of individuals living without major chronic conditions. *PLoS ONE* 16:e0261032. doi: 10.1371/journal.pone.0261032
- Fan, X., Alekseyenko, A. V., Wu, J., Peters, B. A., Jacobs, E. J., Gapstur, S. M., et al. (2018). Human oral microbiome and prospective risk for pancreatic cancer: a population-based nested case-control study. *Gut* 67, 120–127. doi: 10.1136/gutjnl-2016-312580
- Ferrari, E., Pezzi, M. E., Cassi, D., Pertinhez, T. A., Spisni, A., and Meleti, M. (2021). Salivary cytokines as biomarkers for oral squamous cell carcinoma: a systematic review. *Int. J. Mol. Sci.* 22:6795. doi: 10.3390/ijms22136795
- Gilbert, J. A., Blaser, M. J., Caporaso, J. G., Jansson, J. K., Lynch, S. V., and Knight, R. (2018). Current understanding of the human microbiome. *Nat. Med.* 24, 392–400. doi: 10.1038/nm.4517
- Kapila, Y. L. (2021). Oral health's inextricable connection to systemic health: Special populations bring to bear multimodal relationships and factors connecting periodontal disease to systemic diseases and conditions. *Periodontology* 87, 11–16. doi: 10.1111/prd.12398
- Karpiński, T. M. (2019). Role of oral microbiota in cancer development. *Microorganisms* 7:20. doi: 10.3390/microorganisms7010020
- La Rosa, G. R. M., Gattuso, G., Pedullà, E., Rapisarda, E., Nicolosi, D., and Salmeri, M. (2020). Association of oral dysbiosis with oral cancer development. *Oncol. Lett.* 19, 3045–3058. doi: 10.3892/ol.2020.11441
- Martínez, A., Kuraji, R., and Kapila, Y. L. (2021). The human oral virome: shedding light on the dark matter. *Periodontology* 87, 282–298. doi: 10.1111/prd.12396
- Najafi, S., Abedini, F., Azimzadeh Jamalkandi, S., Shariati, P., Ahmadi, A., and Gholami Fesharaki, M. (2021). The composition of lung microbiome in lung cancer: a systematic review and meta-analysis. *BMC Microbiol.* 21:315. doi: 10.1186/s12866-021-02375-z
- Podzimek, S., Vondrackova, L., Duskova, J., Janatova, T., and Broukal, Z. (2016). Salivary markers for periodontal and general diseases. *Dis. Markers* 2016:9179632. doi: 10.1155/2016/9179632
- Proctor, D. M., and Relman, D. A. (2017). The landscape ecology and microbiota of the human nose, mouth, and throat. *Cell Host Microbe* 21, 421–432. doi: 10.1016/j.chom.2017.03.011
- Reitano, E., de'Angelis, N., Gavriilidis, P., Gaiani, F., Memeo, R., Inchingolo, R., et al. (2021). Oral bacterial microbiota in digestive cancer patients: a systematic review. *Microorganisms* 9:2585. doi: 10.3390/microorganisms9122585
- Sedghi, L. M., Bacino, M., and Kapila, Y. L. (2021). Periodontal disease: the good, the bad, and the unknown. *Front. Cell Infect. Microbiol.* 11:766944. doi: 10.3389/fcimb.2021.766944
- Sri Santosh, T., Parmar, R., Anand, H., Srikanth, K., and Saritha, M. (2020). A review of salivary diagnostics and its potential implication in detection of Covid-19. *Cureus* 12:e7708. doi: 10.7759/cureus.7708
- Tada, A., and Senpuku, H. (2021). The impact of oral health on respiratory viral infection. *Dent. J.* 9:43. doi: 10.3390/dj9040043
- To, K. K.-W., Tsang, O. T.-Y., Chik-Yan Yip, C., Chan, K.-H., Wu, T.-C., Chan, J. M. C., et al. (2020). Consistent detection of 2019 novel coronavirus in saliva. *Clin. Infect. Dis.* 71, 841–843. doi: 10.1093/cid/ciaa149
- Yan, X., Yang, M., Liu, J., Gao, R., Hu, J., Li, J., et al. (2015). Discovery and validation of potential bacterial biomarkers for lung cancer. *Am. J. Cancer Res.* 5, 3111–3122.
- Zhang, C.-Z., Cheng, X.-Q., Li, J.-Y., Zhang, P., Yi, P., Xu, X., et al. (2016). Saliva in the diagnosis of diseases. *Int. J. Oral Sci.* 8, 133–137. doi: 10.1038/ijos.2016.38

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Bourgeois, Gonçalves, Lima-Junior and Carrouel. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.