



# The scaffold protein menin is essential for activating the MYC locus and MYC-mediated androgen receptor transcription in androgen receptor-dependent prostate cancer cells

Yakun Luo, Virginie Vlaeminck-guillem, Romain Teinturier, Razan Abou Ziki, Philippe Bertolino, Muriel Le Romancer, Chang Xian Zhang

## ► To cite this version:

Yakun Luo, Virginie Vlaeminck-guillem, Romain Teinturier, Razan Abou Ziki, Philippe Bertolino, et al.. The scaffold protein menin is essential for activating the MYC locus and MYC-mediated androgen receptor transcription in androgen receptor-dependent prostate cancer cells. *Cancer Communications*, 2021, 41 (12), pp.1427-1430. 10.1002/cac2.12217 . hal-03666677

**HAL Id: hal-03666677**

**<https://univ-lyon1.hal.science/hal-03666677>**

Submitted on 1 Dec 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## LETTER TO THE EDITOR

# The scaffold protein menin is essential for activating the *MYC* locus and *MYC*-mediated androgen receptor transcription in androgen receptor-dependent prostate cancer cells

Dear Editor,

The proto-oncogene *MYC* has a well-documented role in driving androgen receptor (AR) oncogenic functions in prostate cancer (PCa cells) [1] and was recently reported as a key activator in the regulation of AR transcription [2]. Dissecting the mechanisms underlying *MYC*-mediated AR regulation may be crucial for better understanding PCa development and improving its management. We recently demonstrated that menin, encoded by the *MEN1* gene, plays an oncosuppressive role in the generation of AR<sup>low</sup>/CD44<sup>+</sup> microinvasive prostate adenocarcinoma in mice but exerts oncogenic effects specifically in AR-dependent human PCa cells, both likely by modulating AR transcription and AR targets [3]. We herein wondered whether this oncogenic function of menin would occur through its interaction with other factors known to play a crucial role for AR transcription in PCa, in particular, *MYC*.

We initially examined the expression of *MYC* upon *MEN1* knockdown (KD) in PCa cells. We found that *MEN1* silencing led to a decrease in *MYC* expression at transcriptional (Fig. 1a, Supplementary Fig. S1A) and protein levels (Fig. 1b, Supplementary Fig. S1B-D) in AR-dependent (VCaP, LNCaP, and CWR22Rv1), but not AR-independent (PC3 and DU145), PCa cells. We also examined *MYC* expression in mouse prostate and found no altered *MYC* expression in menin-deficient, compared to menin-expressing, luminal cells (Fig. S1E). Therefore, we hereafter focused on AR-dependent PCa cells. We detected


the interaction between menin and *MYC* in AR-dependent PCa cells (Supplementary Fig. S2A-B), similar to HeLa cells [4], and found that *MYC*-KD triggered reduced cell proliferation in these cells (Supplementary Fig. S2C-D). Recently, several distal regulatory sequences in the *MYC* locus have been identified, including 2 enhancers situated 67 kb upstream [5] and 20 kb downstream [6] of the transcription start site, respectively named *MYC* enhancer 1 and 2 for the current study. Importantly, ChIP analyses demonstrated that menin bounded not only to the *MYC* promoter but also to these two enhancers (Fig. 1C, Supplementary Fig. S3A). Moreover, menin was necessary for major components of the MLL complex (KMT2A, KMT2B, ASH2L), to which menin belongs [7], to bind to and activate the *MYC* promoter and enhancer 2 likely via H3K4me3 marks (Fig. 1D-E, Supplementary Fig. S3B-E). *MEN1* silencing resulted in decreased expression of known *MYC* target genes, enhanced cell cycling alterations (Supplementary Fig. S4A-C), and decreased expression of AR variants (Supplementary Fig. S4D-E), reminiscent of previous findings in other organs [8]. Notably, the expression of menin and *MYC*, their interaction, and the binding of menin to the *MYC* promoter and enhancer 2, were augmented upon stimulation with the androgen, dihydrotestosterone (Fig. 1F-H, Supplementary Fig. S5). Hence, these data revealed that menin is a crucial factor positively regulating *MYC* transcription specifically in AR-dependent PCa cells.

Next, we investigated whether menin was involved in *MYC*-mediated activation of AR transcription [2]. ChIP analysis and Luciferase reporter assays revealed that menin bounded to and positively regulated several sites on the AR promoter, whereas *MYC* had a greater binding affinity and regulatory activity for an E-box site located -300 kb of the AR promoter (Fig. 1I, Supplementary Fig. S6). Menin is also bounded to the *KLK3* enhancer (Supplementary Fig. S6D). Importantly, deletion of the

**Abbreviations:** AR, Androgen Receptor; AR<sup>low</sup>/CD44<sup>+</sup>, with low AR expression and positive CD44 expression; ASH2L, ASH2 like; ChIP, Chromatin Immunoprecipitation; Chr 1, Chromosome 1; EZH2, Enhancer of zeste homolog 2; KMT2A, Lysine-Specific Methyltransferase 2A; KMT2B, Lysine-Specific Methyltransferase 2B; lncRNA, Long non-coding RNAs; MEN1, Multiple endocrine neoplasia type 1; mCRPC, metastatic castration-resistant PCa; PCa, prostate cancer; PCAT1, Prostate Cancer Associated Transcript 1; siRNA, Small interfering RNA

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Cancer Communications* published by John Wiley & Sons Australia, Ltd. on behalf of Sun Yat-sen University Cancer Center



E-box on the *AR* promoter significantly reduced the transactivation activity of MYC and that of menin in Luciferase reporter assays whereas this latter activity remained unaffected on other parts of the *AR* promoter (Fig. 1J, Supplementary Fig. S6D-G, Supplementary Fig. S7A-C). Remarkably, we found that MYC silencing resulted in less menin binding to the E-box on the *AR* promoter (Fig. 1K, Supplementary Fig. S7D-E). Our data indicate, for the first time, that menin exerts a regulatory role on the MYC-mediated *AR* transcription.

In parallel, we assessed whether menin is involved in the regulation of lncRNA *PCAT1*, situated upstream of the *MYC* locus and known to play an oncogenic role through *MYC* in PCa [9]. Markedly, *MEN1* silencing led to reduced *PCAT1* expression in AR-dependent PCa cell lines (Fig. 1L, Supplementary Fig. S8A). Since EZH2 has been reported to directly regulate *PCAT1* expression and that the interaction between EZH2 and menin has been described in different contexts [10], we investigated whether menin controls *PCAT1* transcription through EZH2 in PCa cells. Here, menin was crucial for the expression of EZH2 (Fig. 1M, Supplementary Fig. S8B), via its physical interaction with EZH2 (Supplementary Fig. S8C-D) and its binding to the *EZH2* promoter (Fig. 1N-O, Supplementary Fig. S8E-F). Finally, ChIP analysis showed that menin bound to the same binding sites as EZH2 on the regulatory sequences of *PCAT1* (Fig. 1P, Supplementary Fig. S8G), the binding being weakened upon *EZH2*-KD (Fig. 1Q, Supplementary Fig. S8H). Taken together, our work unveiled that menin is critically involved in the regulation of MYC-related *PCAT1*, through its interaction with and regulation of EZH2.

Finally, through data mining, significant positive correlations were observed between *MEN1* and *MYC* mRNA expression (Fig. 1R) and between *MEN1* and *EZH2* mRNA expression in both primary PCa and mCRPC (Supplementary Fig. S9). Lastly, the combined use of siMEN1 and the MYC/Max dimerization inhibitor 10058-F4 had a synergistic effect on reducing the proliferation of VCaP, LNCaP, and CWR22Rv1 cells (Fig. 1S, Supplementary Fig. S10).

Collectively, we uncovered a critical crosstalk between menin, AR, and MYC, involving *PCAT1* regulation via EZH2, specifically in AR-dependent PCa cells (Fig. 1T), providing novel clues as to the mechanisms underlying the regulation of *AR* transcription.

## DECLARATIONS

### FUNDING

This study was supported by Epigenetics & Cancer Program (ASC14092CSA), the Ligue Inter-régionale contre le Cancer (R19040CC), Institut National du Cancer (PRTK2017) and the Association: «Le Cancer du sein, parlons-en». R.T. was the recipients of a PhD-fellowship from Ministère de l'Enseignement Supérieur et de la Recherche, France. Y.L. was the recipient of a PhD-fellowship from the China Scholarship Council. R.A.Z. was the recipient of a PhD-fellowship from the Association "G04MEDIA S.A.R.L.", Lebanon.

### ACKNOWLEDGMENTS

We are grateful to Brigitte Manship for her assistance in editing and proofreading the manuscript, Zhi Chong Wu for his scientific and technical input, and Silvere Baron for his scientific discussion and advice.

**FIGURE 1 Menin exerts a regulatory role on the MYC locus and is essential for AR transcription mediated by MYC in AR-dependent PCa cells.** MYC mRNA and protein levels were analyzed by RT-qPCR (A), Western blotting analysis (B) in siCtrl or siMEN1 #1 and siMEN1 #3-co-transfected VCaP and PC3 cells. C-K The experiments were conducted exclusively in AR-dependent VCaP cells. ChIP-qPCR assessing the binding of menin to (C) and H3K4me3 marks (D) on the MYC promoter and enhancers in siMEN1 #1 and siMEN1 #3-co-transfected cells. (E) ChIP-qPCR assessing the binding of menin to the MYC promoter in siCtrl or siASH2L-treated cells. (F) Proximity ligation assay (PLA) showing menin and MYC interaction upon DHT treatment for 24 h. Scale bar, 25  $\mu$ m. Quantification of PLA signals per cell is shown on the right. (G) Western blotting analysis of AR, menin, and MYC levels upon treatment with DHT for 12 h and 24 h. (H) ChIP-qPCR showing the binding of menin to the MYC promoter and enhancer 2 following treatment with DHT for 24 h. Chr 1 was used as the negative control. (I) ChIP-qPCR of the binding of menin to the AR promoter. (J) Relative luciferase activity after transfection of siCtrl- or siMEN1 #1 and siMEN1 #3-co-transfected cells with AR-Luc-2 or AR-Luc-2- $\Delta$ E. (K) ChIP assessing menin occupancy on the AR promoter in siCtrl or siMYC #1 and #2-co-transfected VCaP cells as indicated. (L) RT-qPCR detecting *PCAT1* expression in siCtrl or siMEN1 #1 and siMEN1 #3-transfected cells. (M) RT-qPCR showing EZH2 expression in siCtrl or siMEN1 #1 and siMEN1 #3-co-transfected cells. (N) ChIP evaluating menin occupancy on the *EZH2* promoter. (O) RT-qPCR detecting *PCAT1* expression in siCtrl or siEZH2 #1 and #2-co-transfected cells. (P) ChIP showing menin occupancy on the *PCAT1* promoter. (Q) ChIP showing the effect of siEZH2 #1 and 2 on the levels of menin recruitment to the *PCAT1* promoter. (R) Data mining analysis of the correlation between *MEN1* and *MYC* mRNA expression in primary PCa (left panel) and mCRPC (right panel). (S) Combined use of MYC inhibitor (10058-F4) and siMEN1 #1 and 3 showing effects on cell proliferation. (T) Schematic summary of uncovered functions of menin in AR-dependent PCa cells. For all the RT-qPCR and ChIP-PCR analyses, shown is the mean  $\pm$  s.d. ( $n = 3$ ). t-test, ns, non-significant, \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ .

Abbreviations: AR, androgen receptor; ChIP, chromatin immunoprecipitation; Chr 1, Chromosome 1; Ctrl, Control; DAPI, 4',6-diamidino-2-phenylindole; EZH2, Enhancer of zeste homolog 2; HPRT, Hypoxanthine-guanine phosphoribosyltransferase; MEN1, Multiple endocrine neoplasia type 1; PCa, prostate cancer; *PCAT1*, Prostate Cancer Associated Transcript 1; PCR, Polymerase chain reaction; RT-qPCR, quantitative real-time PCR; si, interfering RNA.



## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

All animal experiments were performed following the animal care guidelines of the European Union and were validated by the local Animal Ethics Evaluation Committee (CECCAPP: C2EA-15 agreed by the French Ministry of High School and Research, « Autorisation de projet » CLB-2012-053).

## AUTHORS' CONTRIBUTIONS

Y.K.L conducted the experiments, analyzed and interpreted the data, prepared figures, and manuscript; V.V.G, T.R, and R.A.Z provided technical and material support and participated in data interpretation; P.B participated in study design and provided critical revision of the manuscript; M.L.R and C.X.Z conceived and supervised the study and manuscript preparation, and obtained funding.

## CONSENT FOR PUBLICATION

All of the authors approved the contents and the data in this manuscript and agreed to its submission for publication.

## CONFLICT OF INTERESTS

The authors declare that they have no conflict of interest.

## AVAILABILITY OF DATA AND MATERIAL

All data generated or analyzed during this study are included in this article.

Yakun Luo<sup>1</sup>  
Virginie Vlaeminck-Guillem<sup>1,2</sup>  
Romain Teinturier<sup>1</sup>  
Razan Abou Ziki<sup>1</sup>  
Philippe Bertolino<sup>1</sup>  
Muriel Le Romancer<sup>1</sup>   
Chang Xian Zhang<sup>1</sup> 

<sup>1</sup> Department of Tumor Escape, Resistance and Immunity, Université Lyon, Université Claude Bernard Lyon 1, INSERM 1052, CNRS 5286, Centre Léon Bérard, Centre de Recherche en Cancérologie de Lyon, Lyon 69008, France

<sup>2</sup> Centre de Biologie Sud, Hôpital Lyon Sud, Hospices Civils de Lyon, Pierre-Bénite 69310, France

Corresponding Author: Muriel Le Romancer, Ph.D., Cancer Research Centre of Lyon, France, F-69008, France; E-mail: [muriel.leromancer-cherifi@lyon.unicancer.fr](mailto:muriel.leromancer-cherifi@lyon.unicancer.fr)

## ORCID

Muriel Le Romancer  <https://orcid.org/0000-0002-8491-4015>

Chang Xian Zhang  <https://orcid.org/0000-0002-5443-8951>

## REFERENCES

1. Grad JM, Dai JL, Wu S, Burnstein KL. Multiple androgen response elements and a Myc consensus site in the androgen receptor (AR) coding region are involved in androgen-mediated up-regulation of AR messenger RNA. *Mol Endocrinol* Baltim Md 1999;13:1896-911. <https://doi.org/10.1210/mend.13.11.0369>.
2. Bai S, Cao S, Jin L, Kobelski M, Schouest B, Wang X, *et al.* A positive role of c-Myc in regulating androgen receptor and its splice variants in prostate cancer. *Oncogene* 2019;38:4977-89. <https://doi.org/10.1038/s41388-019-0768-8>.
3. Teinturier R, Luo Y, Decaussin-Petrucci M, Vlaeminck-Guillem V, Vacherot F, Firlej V, *et al.* Men1 disruption in Nkx3.1-deficient mice results in ARlow/CD44+ microinvasive carcinoma development with the dysregulated AR pathway. *Oncogene* 2021;40:1118-27. <https://doi.org/10.1038/s41388-020-01589-1>.
4. Brès V, Yoshida T, Pickle L, Jones KA. SKIP interacts with c-Myc and Menin to promote HIV-1 Tat transactivation. *Mol Cell* 2009;36:75-87. <https://doi.org/10.1016/j.molcel.2009.08.015>.
5. Wang C, Mayer JA, Mazumdar A, Fertuck K, Kim H, Brown M, *et al.* Estrogen induces c-myc gene expression via an upstream enhancer activated by the estrogen receptor and the AP-1 transcription factor. *Mol Endocrinol* Baltim Md 2011;25:1527-38. <https://doi.org/10.1210/me.2011-1037>.
6. Gao L, Schwartzman J, Gibbs A, Lisac R, Kleinschmidt R, Wilmot B, *et al.* Androgen receptor promotes ligand-independent prostate cancer progression through c-Myc upregulation. *PloS One* 2013;8:e63563. <https://doi.org/10.1371/journal.pone.0063563>.
7. Huang J, Gurung B, Wan B, Matkar S, Veniaminova NA, Wan K, *et al.* The same pocket in menin binds both MLL and JUND but has opposite effects on transcription. *Nature* 2012;482:542-6. <https://doi.org/10.1038/nature10806>.
8. Wu G, Yuan M, Shen S, Ma X, Fang J, Zhu L, *et al.* Menin enhances c-Myc-mediated transcription to promote cancer progression. *Nat Commun* 2017;8:15278. <https://doi.org/10.1038/ncomms15278>.
9. Prensner JR, Chen W, Han S, Iyer MK, Cao Q, Kothari V, *et al.* The long non-coding RNA PCAT-1 promotes prostate cancer cell proliferation through cMyc. *Neoplasia* N Y N 2014;16:900-8. <https://doi.org/10.1016/j.neo.2014.09.001>.
10. Gherardi S, Ripoché D, Mikaelian I, Chanal M, Teinturier R, Goehrig D, *et al.* Menin regulates Inhbb expression through an Akt/Ezh2-mediated H3K27 histone modification. *Biochim Biophys Acta Gene Regul Mech* 2017;1860:427-37. <https://doi.org/10.1016/j.bbaggm.2017.02.003>.

## SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.