



HAL
open science

Determinants of genetic diversity and species richness of North American amphibians

Chloé Schmidt, Jason Munshi-South, Stéphane Dray, Colin J Garroway

► To cite this version:

Chloé Schmidt, Jason Munshi-South, Stéphane Dray, Colin J Garroway. Determinants of genetic diversity and species richness of North American amphibians. *Journal of Biogeography*, 2021, 49 (11), pp.2005-2015. 10.1101/2021.01.04.425301 . hal-03847577

HAL Id: hal-03847577

<https://univ-lyon1.hal.science/hal-03847577>

Submitted on 10 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Title: Determinants of genetic diversity and species richness of North**
2 **American amphibians**

3 **Running title:**

4 **Authors:** Chloé Schmidt^{1*}, Jason Munshi-South², Stéphane Dray³, Colin J Garroway^{1*}

5 **Affiliations:**

6 ¹Department of Biological Sciences, University of Manitoba, Winnipeg, MB, Canada

7 ²Department of Biological Sciences and Louis Calder Center, Fordham University, Armonk, NY,
8 USA

9 ³Univ Lyon, Université Claude Bernard Lyon 1, CNRS, Laboratoire de Biométrie et Biologie
10 Evolutive, F-69100, Villeurbanne, France

11

12 ***Correspondence to:**

13 Chloé Schmidt
14 Department of Biological Sciences
15 50 Sifton Rd
16 University of Manitoba
17 Winnipeg, MB R3T 2N2
18 email: schmid46@myumanitoba.ca

19

20 Colin J Garroway
21 Department of Biological Sciences
22 50 Sifton Rd
23 University of Manitoba
24 Winnipeg, MB R3T 2N2
25 email: colin.garroway@umanitoba.ca

26 **Abstract:** Ecological limits on the population sizes and number of species a region is capable of
27 supporting are thought to simultaneously produce spatial patterns in genetic diversity and species
28 richness. However, we do not know the extent to which resource-based environmental limits
29 jointly determine these patterns of biodiversity in ectotherms because of their low energy
30 requirements compared to endotherms. Here, we adapt a framework for the ways ecological
31 limits may shape genetic diversity and species richness previously tested in mammals for
32 amphibians to determine whether similar processes produce continental patterns of biodiversity
33 across both taxa. Repurposing open, raw microsatellite data from 19 species sampled at 554 sites
34 in North America we found that spatial patterns of genetic diversity run opposite to patterns of
35 species richness and genetic differentiation. However, while measures of resource availability
36 and niche heterogeneity predict 89% of the variation in species richness, these landscape metrics
37 are poor predictors of genetic diversity. Although heterogeneity appears to be an important driver
38 of genetic and species biodiversity patterns in both amphibians and mammals, our results suggest
39 that variation in genetic diversity both within and across species makes it difficult to infer
40 general processes producing spatial patterns of amphibian genetic diversity.

41

42 **Keywords:** latitudinal diversity gradient, biogeography, heterozygosity, frogs, salamanders,
43 more individuals hypothesis, macrogenetics

44

45 **Introduction**

46 Although species richness is higher in the tropics for most taxa, the details of diversity patterns
47 differ among species groups. In North America for instance, vertebrate richness generally
48 increases with resource availability, but mammals and birds tend to have higher species richness
49 in dry, mountainous areas, and reptiles and amphibians are more diverse in wet, lower elevation
50 regions (Currie 1991). This pattern suggests that while richness increases with resource
51 availability, taxon-specific traits may cause richness patterns to diverge from a strictly latitudinal
52 gradient. Broad-scale patterns of biodiversity at the genetic level have only recently begun to be
53 mapped thanks to the accumulation of open data in public repositories (Miraldo et al. 2016;
54 Manel et al. 2020; Schmidt et al. 2020; Theodoridis et al. 2020). Genetic diversity is typically
55 thought of as the most fundamental level of biodiversity because it bears on populations'
56 capacities to evolve adaptively in response to environmental change (Frankham 1995). Recent
57 analyses of mammals suggest that environments can simultaneously shape species richness and
58 genetic diversity on continental scales (Schmidt et al. 2020). Whether this is also true in
59 ectothermic taxa is unknown. Understanding whether common processes underlie variation in
60 biogeographic patterns of diversity across taxa with different environmental requirements can
61 help us move toward a general understanding of the drivers of biodiversity at multiple levels.
62 In mammals, continental scale multi-species patterns of nuclear genetic diversity and species
63 richness could be inferred with estimates of resource and niche availability, or ecological
64 opportunity (Schmidt et al. 2020). This suggested that ecological limits placed on the number of
65 individuals and species an environment can support are important drivers of broad-scale
66 biodiversity patterns. Resource-rich environments supported larger populations with higher
67 genetic diversity and species richness, while niche availability in heterogeneous habitats

68 promoted species coexistence but reduced population sizes and genetic diversity because
69 specialization restricts available resources. These processes are related to two well-supported
70 hypotheses for the production of biogeographic patterns of species richness: the more-individuals
71 hypothesis (Wright 1983), and the effects of environmental and resource heterogeneity (Allouche
72 et al. 2012; Stein et al. 2014). The more individuals hypothesis posits that resource-rich regions
73 near the equator are capable of supporting larger populations and communities, and thus more
74 species than temperate regions. Heterogeneity is thought to increase species richness due to
75 greater niche availability in more complex heterogeneous environments which allows more
76 species to coexist, but with smaller population sizes because resources are partitioned.

77 The effects of resource heterogeneity on species richness and genetic diversity seem likely to be
78 generally applicable across taxa (Stein et al. 2014; Schmidt et al. 2020), however the general
79 relevance of the more-individuals hypothesis for ectotherms is uncertain (Buckley and Jetz
80 2010). This is because compared to endotherms, ectotherms have lower energy requirements and
81 can behaviorally thermoregulate, meaning their abundances are less likely to be limited by
82 resource-related ecological limits (Pough 1980; Buckley and Jetz 2010). Instead, ectotherm
83 distributions, and therefore species richness, seem more directly constrained by environmental
84 temperature because fewer species have evolved thermal adaptations required for expanding into
85 cooler regions (Buckley and Jetz 2007, 2010). Further, the evolution of traits associated with
86 better survival in temperate regions may have additional effects on speciation dynamics. For
87 example, species turnover tends to be higher among viviparous squamate reptiles, which
88 typically occupy cooler regions (Pyron and Burbrink 2014). Reaching a more comprehensive
89 understanding of how biodiversity patterns emerge requires knowledge about whether

90 relationships between genetic diversity, species richness, and environments are consistent across
91 endothermic and ectothermic taxa.

92 The determinants of species richness across all terrestrial vertebrates are generally related to
93 resource availability as estimated by energy (e.g., potential evapotranspiration, primary
94 productivity), water-energy balance (e.g., actual evapotranspiration, precipitation), and
95 heterogeneity (e.g., elevation variability, land cover diversity) (Currie 1991; Kerr and Packer
96 1997; Hawkins et al. 2003; Rodríguez et al. 2005; Buckley and Jetz 2007; Stein et al. 2014;
97 Jiménez-Alfaro et al. 2016). Amphibians are interesting because they are constrained both by
98 water availability and temperature and water availability is consistently identified as an
99 important driver of diversity in amphibians (Rodríguez et al. 2005; Buckley and Jetz 2007).

100 Indeed in Europe, the best predictors of species richness in mammals and birds shift from energy
101 to water availability at decreasing latitudes, but amphibian species richness remains strongly
102 related to water-energy balance regardless of latitude (Whittaker et al. 2007).

103 The causes of population genetic diversity are rarely studied at the same time or scale as patterns
104 of species richness (but see Marshall and Camp 2006; Schmidt et al. 2020), yet the presumed
105 mechanisms related to the more-individuals hypothesis and heterogeneity are closely related to
106 carrying capacity and population-level processes (Schmidt et al. 2020). The more-individuals
107 mechanism predicts a positive relationship between species richness and population genetic
108 diversity because bigger populations and communities tend to have higher levels of genetic and
109 species diversity (Kimura 1983; Hubbell 2001; Schmidt et al. 2020). With higher carrying
110 capacities, more species persist because they can reach minimal viable population sizes. On the
111 other hand, heterogeneity is predicted to cause negative correlations between genetic diversity
112 and species richness by increasing the number of species a given area can support which in turn

113 reduces population size and limits gene flow due to increased niche specialization.

114 Heterogeneous environments also facilitate population differentiation due to spatially varying
115 selection. In mammals, evolutionary processes acting on the population level scaled up and
116 interacted with resource availability and heterogeneity to produce genetic diversity and species
117 richness patterns (Schmidt et al. 2020).

118 Whether mechanisms related to ecological limits and niche availability predict patterns of
119 species richness and genetic diversity in ectotherms is unclear. To test this prediction, we
120 analyzed previously-published microsatellite genotype data from 19 North American amphibian
121 species (8 frogs, 11 salamanders), with >13000 individuals sampled at 554 sites. Our first
122 objective was to identify existing spatial patterns in genetic diversity and differentiation and
123 quantify the extent to which genetic diversity and species richness covary. We then tested
124 whether limits on resources and niche availability jointly determined genetic diversity and
125 species richness using structural equation models, which allowed us to evaluate multiple
126 hypotheses at genetic and species level biodiversity simultaneously. We based our conceptual
127 framework on previous findings in mammals (Schmidt et al. 2020) and adapted it for
128 amphibians. First, we excluded human presence because previous investigation shows it did not
129 have a clear effect on amphibian genetic diversity (Schmidt and Garroway 2021a). Second, we
130 included a measure of water availability as an additional indicator of resource availability. Third,
131 body size was used as a proxy for whole species census size in mammals, however we do not
132 know the extent to which body size is correlated with effective population size or genetic
133 diversity within amphibians. We therefore first tested for a relationship between body size and
134 genetic diversity for species in our sample to determine whether it should be incorporated into
135 the model. Finally, we compare our results to previous results in mammals (Schmidt et al. 2020)

136 to infer whether similar environmental features contribute to diversity gradients across
137 endothermic and ectothermic taxa in North America.

138

139 **Methods**

140 **Biodiversity data**

141 *Genetic diversity and differentiation.* We used raw genotypes (i.e. called allele sizes) of North
142 American amphibians compiled by (Schmidt and Garroway 2021a). This data set was assembled
143 from raw microsatellite datasets publicly archived in Dryad (DataDryad.org). To identify data
144 sets we conducted a systematic search of the Dryad data repository with the following keywords:
145 species name (e.g. *Plethodon cinereus*), “microsat*”, “short tandem*”, and “single tandem*”.
146 We used the IUCN Red List database to obtain a list of amphibian species native to North
147 America for the search. We excluded datasets that lacked spatial reference, were not located in
148 North America, did not sample neutral microsatellite loci, or had study designs that may have
149 affected genetic diversity (including sampling island populations, or captive or managed
150 populations). In total the data sets include genotypes for 13680 individuals spanning 19 species
151 sampled at 554 locations in the contiguous United States and Canada. We used gene diversity as
152 a measure of genetic diversity because it is minimally affected by sample size (Nei 1973;
153 Charlesworth and Charlesworth 2010). Our measure of differentiation was population-specific
154 F_{ST} (Weir and Goudet 2017). Population-specific F_{ST} differs from pairwise F_{ST} in that it
155 measures how far single populations in a sample have diverged from a common ancestor—this
156 means it is comparable across studies and species. Population-specific F_{ST} was not estimable at 2
157 sites because it requires a minimum of 2 sample sites in the original dataset ($n = 552$).

158 *Species richness.* We estimated species richness at each of our genetic diversity sample sites
159 using amphibian range extent data from the IUCN RedList (IUCN 2019), applying filters for
160 native, extant species ranges. We measured species richness as the number of species' ranges
161 overlapping each genetic sample site.

162

163 **Diversity maps and spatial variation partitioning**

164 We used distance-based Moran's eigenvector maps (MEMs) to detect spatial patterns in genetic
165 diversity and differentiation and compare these to patterns of species richness. MEMs are
166 orthogonal spatial eigenvectors with eigenvalues that are directly proportional to Moran's *I*.
167 They measure spatial autocorrelation at all scales present in the data. We computed MEMs in the
168 R package *adespatial* (Dray et al. 2017). We used the forward selection procedure described in
169 Blanchet et al. (2008) to select two sets of MEMs describing important patterns in genetic
170 diversity, differentiation, and species richness. We then selected MEMs which explained the
171 broadest spatial patterns in these data (Moran's $I > 0.25$). To create maps of genetic diversity,
172 differentiation, and species richness (Fig. 1), we used the fitted values for gene diversity,
173 population-specific F_{ST} , and species richness regressed on broad-scale MEMs. This approach
174 allowed us to visualize broad purely spatial patterns in genetic diversity, differentiation, and
175 species richness without variation due to other sources, such as local environments, species
176 identity, or population history. Maps of raw values are presented in Fig. S1.

177 Next, we determined the extent to which spatial patterns in genetic diversity and species richness
178 were shared using variation partitioning. Because our MEM analysis for both levels of
179 biodiversity had the same input distance matrix, the resulting spatial MEMs were directly

180 comparable. This was not the case for genetic differentiation, which had fewer sample sites. We
181 therefore did not partition variation in genetic differentiation because these MEMs are not the
182 same as those computed for genetic diversity and species richness.

183 We determined the fraction of total variation explained by spatial structure, shared spatial
184 structure, and non-spatial variation using variation partitioning as follows. We ran a series of
185 linear regressions with either species richness (y_{SR}) or gene diversity (y_{GD}) as the response
186 variable using all MEMs selected for that variable (Equations 1 and 2), or only MEMs shared by
187 both variables as predictors (Equations 3 and 4):

$$188 \quad y_{SR} \sim \alpha + \beta_{1S}(\text{MEM}_{1S}) + \beta_{2S}(\text{MEM}_{2S}) + \dots + \beta_{8S}(\text{MEM}_{8S}) + \epsilon \quad \text{Eq. 1}$$

$$189 \quad y_{GD} \sim \alpha + \beta_{1G}(\text{MEM}_{1G}) + \beta_{2G}(\text{MEM}_{2G}) + \dots + \beta_{5G}(\text{MEM}_{5G}) + \epsilon \quad \text{Eq. 2}$$

$$190 \quad y_{SR} \sim \alpha + \beta_{1SG}(\text{MEM}_{1SG}) + \beta_{2SG}(\text{MEM}_{2SG}) + \dots + \beta_{5SG}(\text{MEM}_{5SG}) + \epsilon \quad \text{Eq. 3}$$

$$191 \quad y_{GD} \sim \alpha + \beta_{1SG}(\text{MEM}_{1SG}) + \beta_{2SG}(\text{MEM}_{2SG}) + \dots + \beta_{5SG}(\text{MEM}_{5SG}) + \epsilon \quad \text{Eq. 4}$$

192

193 where α is the grand mean, and MEM_{iS} and MEM_{iG} are the set of MEMs selected for species
194 richness (8 MEMs) and genetic diversity (5 MEMs), respectively. The coefficients of variation
195 (R^2) from Eqs. 1 and 2 give the total amount of variation explained by spatial patterns for species
196 richness and genetic diversity. Subtracting these values from 1 gives the amount of non-spatial
197 variation. MEM_{iSG} represents the set of MEMs shared by both species richness and genetic
198 diversity (5 MEMs). R^2 values from Eqs. 3 and 4 tell us the amount of variation in each response
199 variable which can be explained by spatial variation shared at both levels of diversity. When
200 subtracted from the total spatial variation in genetic diversity or species richness (Eqs. 1 and 2),
201 we get the proportion of non-shared spatial variation.

202

203 **Structural equation modeling**

204 *Environmental data.* We measured resource and niche heterogeneity by computing Simpson's
205 Diversity Index for landcover categories within buffers at each site. We obtained a 30 m
206 resolution map of North American landcover data from the Commission for Environmental
207 Cooperation (CEC et al. 2015). This map is based on 2015 satellite imagery and has 19 standard
208 land cover classifications which includes forests, shrubland, grassland, wetlands, cropland,
209 barren land, and built-up land. Because this variable is scale-dependent, we recorded
210 heterogeneity within 4 buffer sizes (10, 25, 40, and 80 km) around each site.

211 We used two measures of resource availability because amphibians are habitat-limited by both
212 temperature and water availability. Water availability can be measured by evapotranspiration, or
213 the amount of water removed from the Earth's surface through soil or open water evaporation
214 and plant transpiration processes. Potential evapotranspiration (PET) measures the atmospheric
215 demand for water, depending on factors such as temperature and wind (Peng et al. 2019). It is
216 strongly correlated with temperature. PET is the maximum amount of water that would be
217 removed in the absence of biophysical limitations (Peng et al. 2019). The amount of water
218 actually removed, actual evapotranspiration (AET), reflects water availability and soil moisture
219 levels. Actual evapotranspiration has also been shown to be one of the strongest predictors of
220 amphibian species richness (Buckley and Jetz 2007). PET can be viewed as a measure of energy
221 availability, and AET one of water-energy balance (see Currie 1991; Buckley and Jetz 2007;
222 Kreft and Jetz 2007). Together these variables represent total ecosystem resource availability at
223 sites. We measured mean PET and AET (mm/yr) at sites within 10, 25, 40, and 80 km buffers
224 using data from the CGIAR Consortium for Spatial Information (Trabucco and Zomer 2019).

225 *Population size.* Body size is a readily measured trait that is typically negatively correlated with
226 long-term effective population size across a diversity of species (Frankham 1996; Romiguier et
227 al. 2014; Mackintosh et al. 2019). To explore whether body size was a useful substitute for long-
228 term effective population size in amphibians, we obtained body length data from the AmphiBIO
229 v1 database (Oliveira et al. 2017). We then regressed it on gene diversity in a hierarchical model
230 with species as a random effect allowing intercepts to vary and predicted a negative relationship.
231 We found no relationship between body size and genetic diversity in our data (Fig. S2), thus we
232 excluded population size from our proposed causal framework. Without population size as an
233 intermediate variable, we predicted that resource availability would have direct positive effects
234 on both genetic diversity and species richness, and that heterogeneity would have direct negative
235 and positive effects on genetic diversity and species richness, respectively. We hypothesize that
236 these direct effects would act through population size were such data available.

237 *Analysis.* We used structural equation modeling (SEM) to determine whether genetic diversity
238 and species richness are shaped by differential ecological limits due to limits on resources and
239 niche availability. Structural equation modeling begins with a causal diagram, or conceptual
240 model, where paths between variables represent hypothesized causal relationships (Fig. 2a).
241 Hypotheses are envisioned as a network where variables are nodes, and paths connecting them
242 represent causal relationships. In SEM, the effects of multiple predictors are simultaneously
243 assessed for multiple response variables (Shipley 2016). We implemented structural equation
244 models using the piecewiseSEM package (version 2.0.2), which uses a local estimation approach
245 for models in the hypothesis network allowing for the incorporation of more complex model
246 types (Lefcheck et al. 2019). Model fit is evaluated using tests of directed separation (Shipley
247 2016), which determine whether an association exists between two variables in the network

248 conditional on each of their causes. If two variables are not conditionally independent, the model
249 is updated by adding a path between them to make the model more consistent with the data. *P*-
250 values from tests of directed separation are used to calculate Fisher's *C*, which is then used to
251 calculate an overall *p*-value for the model network. Models are a good fit to the data when *p* >
252 0.05, indicating the null hypothesis—the proposed hypothesis network—is not rejected.

253 Our model network consisted of two models with genetic diversity and species richness as
254 response variables. Our genetic data is hierarchical with multiple sites nested within species. To
255 account for variation in mean genetic diversity across species we modeled random intercepts for
256 species. Additionally, we allowed the responses of genetic diversity to resource availability and
257 heterogeneity to vary across species by including a random slope term for each environmental
258 predictor (PET, AET, and land cover diversity). With this random effect structure, we thus allow
259 relationships to vary among species and do not assume effects will be of the same magnitude or
260 direction for all species. We fit the hierarchical model for genetic diversity using the lme4
261 package (Bates et al. 2015) within piecewiseSEM. We scaled and centered all variables before
262 analysis so path coefficients were comparable. Finally, we checked model residuals for spatial
263 autocorrelation using Moran tests. We conducted SEM analyses in parallel for each of our 4
264 buffer sizes.

265

266 **Effect of heterogeneity on population differentiation**

267 Finally, we tested whether landscape heterogeneity was related to increased population
268 differentiation. We regressed heterogeneity on population-specific F_{ST} using a hierarchical model
269 with a random effect for species accounting for differences in mean F_{ST} (intercepts) while

270 allowing the strength and direction of the effect of heterogeneity (slopes) to vary across species.
271 To account for spatial variation we included MEMs describing spatial patterns in F_{ST} as
272 covariates. We performed these analyses in parallel across all four heterogeneity buffers.

273

274 **Results**

275 *Spatial patterns in genetic variation.* We detected spatial patterns across genetic diversity,
276 genetic differentiation, and species richness (Fig. 1). The major axis of broad-scale variation is
277 longitudinal, and across all three biodiversity metrics the western sample sites show little
278 variation whereas spatial patterns were more complex in the east and appear to vary latitudinally.
279 This broad longitudinal pattern is consistent with environmental variation in North America (Fig.
280 1) as well as genetic diversity and species richness patterns in mammals (Schmidt et al. 2020).
281 We recovered known patterns of amphibian species richness with MEMs, where richness was
282 highest in the southeastern United States (Fig. 1) (Currie 1991). Western USA, which is hotter
283 and drier, had a comparatively low number of species. In eastern sample sites, genetic diversity
284 increased with latitude, while differentiation and species richness increase towards the tropics.
285 Genetic diversity and differentiation in the western samples were in the mid-range of values
286 across sample sites (Fig. 1).

287 In general, species richness was more spatially structured than genetic diversity, with 85% and
288 23% of variation explained by spatial patterns, respectively (Fig. 1). We detected shared spatial
289 patterns between both levels of biodiversity, however, while shared patterns accounted for the
290 entirety of the spatial variation in genetic diversity, they explained less of the variation in species
291 richness (18%).

292 *Common causes of genetic diversity and species richness.* Our conceptual model (Fig 2a) fit the
293 data well (Fisher's $C = 0.49$, $p = 0.78$, 2 degrees of freedom) with no additional links suggested
294 at any scale. Note that for SEM, $p > 0.05$ means that our conceptual model is not rejected. We
295 present results from the 40 km buffer in the main text; results from all models can be found in
296 tables S1 – 4. Species richness was well explained ($R^2 = 0.89$) and increased with water
297 availability, environmental heterogeneity, and species body size (Fig 2, Table S3). Water
298 availability had the strongest effect on species richness. Genetic diversity was not well predicted
299 by any variables in our model ($R^2 = 0.04$; Fig. 2). Residuals from genetic diversity models did
300 not exhibit spatial autocorrelation. Species richness residuals were spatially autocorrelated at
301 local scales (Moran's $I = 0.06$). In general, the environmental covariates in our models captured
302 broad spatial patterns well, and we did not incorporate fine-scale spatial structure into our
303 models as this was likely due to clustered sampling of some species. Lastly, genetic
304 differentiation within species decreased with heterogeneity ($\beta = -0.32 \pm 0.10$ SE) at the most
305 local spatial scale we tested (10 km buffer), but this relationship disappeared at larger buffer
306 sizes.

307

308 **Discussion**

309 We found spatial variation shared across genetic diversity, differentiation, and species richness at
310 broad spatial scales (Fig. 1). In general, areas with high species richness tended to have
311 genetically differentiated populations with relatively low genetic diversity. There was a
312 latitudinal gradient in genetic diversity across eastern sites which varied in the opposite direction
313 of the gradient in species richness and genetic differentiation. These patterns are consistent with
314 our predicted effects of heterogeneity, however, these relationships were not well-reflected by

315 our structural equation model (Fig. 2). Our variation partitioning suggests that all spatial
316 variation in genetic diversity was shared with species richness, but the environments that
317 predicted species richness did not predict genetic diversity well. This finding suggests that other
318 processes drive the genetic diversity patterns we detect. These processes are likely species-
319 specific, and not generalizable at continental scales as they were in mammals (Schmidt et al.
320 2020).

321 We suspect the general lack of relationship between genetic diversity and species richness and
322 climate we find in our structural equation model may come down to amphibian population
323 dynamics. Ecological limits hypotheses assume that communities are in equilibrium with respect
324 to speciation, colonization, and extinction dynamics (Storch et al. 2018)— extending this to the
325 genetic level, we also assume populations are in an equilibrium state with regards to gene flow,
326 mutation, and genetic drift. However, amphibians have variable local population sizes which can
327 sometimes fluctuate by orders of magnitude from year to year (Collins et al. 2009), high rates of
328 species turnover at local and regional scales (Werner et al. 2007; Buckley and Jetz 2008), and
329 relatively low occupancy within potential distributions (Munguía et al. 2012). Species turnover at
330 sites is associated with environmental heterogeneity in freshwater habitats, especially
331 hydroperiod variation in temporary ponds (Urban 2004). Varying population dynamics could
332 obscure general relationships between population genetic diversity and population size, species
333 richness, and the climatic factors we explore here.

334 Interestingly, although genetic diversity was not affected by environmental heterogeneity at any
335 scale, genetic differentiation decreased with heterogeneity at the most local scale we tested;
336 genetic differentiation was relatively low in the northeast where landscape heterogeneity was
337 higher (Fig. 1). If heterogeneity increases niche availability and creates opportunities for

338 specialization and divergence, we predicted that it would increase genetic differentiation.
339 However, the pattern we detect here may be expected if species that were capable of recolonizing
340 northern regions following glaciation tend to be widely distributed generalists that maintain
341 population connectivity over relatively large geographic distances (Smith et al. 2005; Zeisset and
342 Beebee 2008). Niche partitioning in amphibians may also occur at finer scales within suitable
343 habitats (Karlin et al. 1984; Cloyd and Eason 2017)—for example by modifying microhabitat
344 usage, diets, foraging strategies, or behaviors across species. Thus, the landscape level metric we
345 use here may not capture the varying ways that environmental heterogeneity affects genetic
346 diversity and population differentiation if species respond to environments in different ways.
347 These patterns point to the importance of species-specific responses to environmental conditions
348 and environmental instability in generating biogeographic patterns of genetic diversity and
349 species richness in amphibians (Urban 2004).

350 Previous exploration of the relationships between nuclear genetic diversity, species richness, and
351 environments in plethodontid salamanders produced similarly mixed results (Marshall and Camp
352 2006). Genetic diversity was positively associated with temperature and rainfall across all eight
353 species studied, but topographic heterogeneity had both positive and negative effects on genetic
354 diversity depending on species. Allelic richness was only correlated with species richness for
355 *Desmognathus* species and *Plethodon jordani*, but relationships were positive and negative
356 (Marshall and Camp 2006). In another example, Karlin et al. (1984) report a negative
357 relationship between species richness and genetic diversity across populations of *D. fuscus*.
358 Similar to our findings, it appears resource availability and heterogeneity simultaneously affect
359 biodiversity on genetic and species levels, but genetic diversity in general is less well predicted

360 by environments alone. Variation between species may reduce our ability to detect general
361 relationships across species.

362 Although we did not detect clear latitudinal or longitudinal gradients in genetic diversity across
363 North America, previous findings suggest mitochondrial genetic diversity in amphibians and
364 other ectotherms varies latitudinally and mirrors species richness patterns (Miraldo et al. 2016;
365 Manel et al. 2020). Miraldo et al. (2016) found that amphibian mitochondrial genetic diversity in
366 North America was highest in the species-rich southeastern United States. They suggested this
367 pattern may be related to the evolutionary speed hypothesis, where presumably high
368 environmental temperature increases rates of population divergence and speciation through its
369 effects on mutation rate and generation time (Miraldo et al. 2016). In addition to a negative
370 correlation between patterns of nuclear genetic diversity and species richness, we detected no
371 effect of temperature on nuclear genetic diversity in our SEM, casting doubt on this hypothesis
372 for amphibians. Furthermore, a lack of latitudinal gradient indicates that nuclear genetic diversity
373 is not related to temperature in a straightforward way. Environmental temperature has varied and
374 complex effects on metabolism and mitochondrial processes (Munro and Treberg 2017; Zhang
375 and Wong 2021), making relationships with nuclear and mitochondrial mutation rates unlikely to
376 be generalizable (Lanfear et al. 2007; Schmidt and Garroway 2021b). More generally
377 mitochondrial DNA alone is not a reliable marker for detecting intraspecific demographic
378 patterns (Bazin et al. 2006; Galtier et al. 2009; Schmidt and Garroway 2021b). Thus, marker
379 choice is very likely responsible for the diverging patterns we find here. This also appears to be
380 true in mammals, where patterns of genetic diversity measured using mitochondrial DNA and
381 nuclear DNA also trended in opposite directions (Schmidt et al. 2020).

382 Despite limitations modeling population size, it appears that spatial patterns in genetic diversity
383 and species richness in amphibians are driven by processes similar to mammals. Interestingly,
384 similar proportions of variation in genetic diversity (~ 25%) could be attributed to spatial
385 processes across both taxa. Additionally, it was the case in both groups that spatial variation in
386 genetic diversity was primarily due to factors that also shaped species richness, but other factors
387 contribute to unique spatial variation in species richness. The overall negative correlation
388 between spatial patterns of species richness and genetic diversity in amphibians and mammals
389 (Schmidt et al. 2020) suggests heterogeneity and niche partitioning are major contributors to
390 diversity across genetic and species levels in endo- and ectothermic vertebrates. Heterogeneity
391 has previously been put forth as a universal driver of species richness (Stein et al. 2014). In
392 general, similar environmental factors seem capable of generating an overall species richness
393 gradient across taxa, but slight deviations from this general pattern are mediated by differential
394 interactions between environments, species traits, and population processes.

395 Genetic diversity and species richness are two important metrics for biodiversity conservation
396 because they contribute to the resilience of populations and communities in rapidly changing
397 environments (Oliver et al. 2015). Neutral genetic diversity is indicative of population mean
398 fitness and the efficiency of selection in response to environmental change—although it is only
399 weakly correlated with additive genetic diversity, it is nevertheless informative for conservation
400 purposes because it reflects levels of inbreeding and the efficiency of selection due to its
401 relationship to the effective population size (Frankham 1995; Mittell et al. 2015). Amphibians
402 are among the most imperiled vertebrates (Stuart et al. 2004) and are especially susceptible to
403 environmental change. Macrogenetics approaches mapping multispecies patterns of genetic
404 diversity at broad scales have great potential for incorporation into conservation policies

405 targeting regional conservation of genetic diversity. However, complex ecophysiological
406 requirements, life histories, and population dynamics may render this approach impractical for
407 amphibians because the environmental factors affecting genetic diversity may differ depending
408 on species (Schmidt and Garroway 2021a). Species-specific measures of environmental
409 heterogeneity, resource availability, and habitat suitability may prove to be more reliable
410 predictors of genetic diversity, but may be less relevant for species richness. We are only
411 beginning to explore broad scale patterns of intraspecific genetic diversity across several species,
412 but it is already apparent that they are not as consistently clear as gradients in species richness
413 (Miraldo et al. 2016; Manel et al. 2020; Schmidt et al. 2020; Theodoridis et al. 2020). We look
414 forward to the continued exploration of these patterns in other taxonomic groups to help build a
415 comprehensive picture of the distribution of genetic biodiversity across the globe.

416

417 **Author contributions:** C.J.G., and C.S. conceived of the study. C.S., J.M.S, S.D. and C.J.G.
418 designed the study and C.S. conducted the analyses with input from S.D. and C.J.G. All authors
419 contributed to data interpretation. C.S. wrote the first draft of the manuscript and all authors
420 participated in editing subsequent manuscript drafts.

421

422 **Acknowledgements:** We would like to thank Mitchell Green for assistance with data
423 acquisition, as well as the authors of the original datasets for making their data public. C.S. and
424 C.J.G. were supported by a Natural Sciences and Engineering Research Council of Canada
425 Discovery Grant to C.J.G. C.S. was also supported by a U. Manitoba Graduate Fellowship, and a
426 U. Manitoba Graduate Enhancement of Tri-council funding grant to C.J.G.

427 **Data availability:** Synthesized genetic data is available from the Dryad Data Repository (DOI:
428 10.5061/dryad.qv9s4mwf0). Species range boundary files and environmental data are available
429 from open online sources (see Methods).

430

431 **References**

- 432 Allouche, O., Kalyuzhny, M., Moreno-Rueda, G., Pizarro, M., and Kadmon, R. 2012. Area-
433 heterogeneity tradeoff and the diversity of ecological communities. *Proc. Natl. Acad. Sci.*
434 *U. S. A.* **109**(43): 17495–17500. doi:10.1073/pnas.1208652109.
- 435 Bates, D., Mächler, M., Bolker, B., and Walker, S. 2015. Fitting linear mixed-effects models
436 using lme4. *J. Stat. Softw.* **67**(1). doi:10.18637/jss.v067.i01.
- 437 Bazin, E., Glémin, S., and Galtier, N. 2006. Population size does not influence mitochondrial
438 genetic diversity in animals. *Science* **312**(5773): 570–572. doi:10.1126/science.1122033.
- 439 Blanchet, G.F., Legendre, P., and Borcard. 2008. Forward selection of explanatory variables.
440 *Ecology* **89**(9): 2623–2632.
- 441 Buckley, L.B., and Jetz, W. 2007. Environmental and historical constraints on global patterns of
442 amphibian richness. *Proc. R. Soc. B Biol. Sci.* **274**(1614): 1167–1173.
443 doi:10.1098/rspb.2006.0436.
- 444 Buckley, L.B., and Jetz, W. 2008. Linking global turnover of species and environments. *Proc.*
445 *Natl. Acad. Sci. U. S. A.* **105**(46): 17836–17841. doi:10.1073/pnas.0803524105.
- 446 Buckley, L.B., and Jetz, W. 2010. Lizard community structure along environmental gradients. *J.*
447 *Anim. Ecol.* **79**(2): 358–365. doi:10.1111/j.1365-2656.2009.01612.x.
- 448 CEC, NRCan/CCMEO, USGS, INEGI, CONABIO, and CONAFOR. 2015. 2015 North
449 American Land Cover at 30 m spatial resolution. Available from <http://www.cec.org/north-american-environmental-atlas/land-cover-30m-2015-landsat-and-rapideye/>.
- 451 Charlesworth, B., and Charlesworth, D. 2010. *Elements of evolutionary genetics*. Roberts &
452 Company Publishers, Greenwood Village, Colorado, USA.
- 453 Cloyd, C.S., and Eason, P.K. 2017. Niche partitioning and the role of intraspecific niche
454 variation in structuring a guild of generalist anurans. *R. Soc. Open Sci.* **4**(3).
455 doi:10.1098/rsos.170060.
- 456 Collins, J.P., Crump, M.L., and Lovejoy III, T.E. 2009. *Extinction in Our Times: Global*
457 *Amphibian Decline*. Oxford University Press, Oxford.
- 458 Currie, D.J. 1991. Energy and large-scale patterns of animal- and plant-species richness. *Am.*
459 *Nat.* **137**(1): 27–49.
- 460 Dray, S., Blanchet, G., Borcard, D., Clappe, S., Guenard, G., Jombart, T., Larocque, G.,
461 Legendre, P., Madi, N., and Wagner, H.H. 2017. *adespatial: Multivariate Multiscale Spatial*

- 462 Analysis. Available from <https://cran.r-project.org/package=adespatial>.
- 463 Frankham, R. 1995. Conservation Genetics. *Annu. Rev. Genet.* **29**: 305–332.
- 464 Frankham, R. 1996. Relationship of Genetic Variation to Population Size in Wildlife. *Conserv.*
465 *Biol.* **10**(6): 1500–1508. doi:10.1046/j.1523-1739.1996.10061500.x.
- 466 Galtier, N., Nabholz, B., Glémin, S., and Hurst, G.D.D. 2009. Mitochondrial DNA as a marker
467 of molecular diversity: A reappraisal. *Mol. Ecol.* **18**(22): 4541–4550. doi:10.1111/j.1365-
468 294X.2009.04380.x.
- 469 Hawkins, B.A., Field, R., Cornell, H. V., Currie, D.J., Guégan, J.-F., Kaufman, D.M., Kerr, J.T.,
470 Mittelbach, G.G., Oberdorff, T., O'Brien, E.M., Porter, E.E., and Turner, J.R.G. 2003.
471 Energy, water, and broad-scale geographic patterns of species richness. *Ecology* **84**(12):
472 3105–3117. doi:10.1890/03-8006.
- 473 Hubbell, S.P. 2001. *The Unified Neutral Theory of Biodiversity and Biogeography*. Princeton
474 University Press, Princeton NJ.
- 475 IUCN. 2019. The IUCN Red List of Threatened Species. Version 2019-1. Available from
476 <https://www.iucnredlist.org>.
- 477 Jiménez-Alfaro, B., Chytrý, M., Mucina, L., Grace, J.B., and Rejmánek, M. 2016. Disentangling
478 vegetation diversity from climate-energy and habitat heterogeneity for explaining animal
479 geographic patterns. *Ecol. Evol.* **6**(5): 1515–1526. doi:10.1002/ece3.1972.
- 480 Karlin, A.A., Guttman, S.I., Rathbun, S.L., and Rathbun, S.L. 1984. Spatial autocorrelation
481 analysis of heterozygosity and geographic distribution in populations of *Desmognathus*
482 *fuscus* (Amphibia: Plethodontidae). *Copeia* **1984**(2): 343–356.
- 483 Kerr, J.T., and Packer, L. 1997. Habitat heterogeneity as a determinant of mammal species
484 richness. *Nature* **385**: 253–254.
- 485 Kimura, M. 1983. *The Neutral Theory of Molecular Evolution*. Cambridge University Press,
486 Cambridge.
- 487 Kreft, H., and Jetz, W. 2007. Global patterns and determinants of vascular plant diversity. *Proc.*
488 *Natl. Acad. Sci.* **104**(14): 5925–5930. doi:10.1073/pnas.0608361104.
- 489 Lanfear, R., Thomas, J.A., Welch, J.J., Brey, T., and Bromham, L. 2007. Metabolic rate does not
490 calibrate the molecular clock. *Proc. Natl. Acad. Sci. U. S. A.* **104**(39): 15388–15393.
491 doi:10.1073/pnas.0703359104.
- 492 Lefcheck, J., Byrnes, J., and Grace, J. 2019. *piecewiseSEM: Piecewise Structural Equation*
493 *Modeling*. Available from <https://cran.r-project.org/package=piecewiseSEM>.
- 494 Mackintosh, A., Laetsch, D.R., Hayward, A., Charlesworth, B., Waterfall, M., Vila, R., and
495 Lohse, K. 2019. The determinants of genetic diversity in butterflies. *Nat. Commun.* **10**(1):
496 1–9. doi:10.1038/s41467-019-11308-4.
- 497 Manel, S., Guerin, P.E., Mouillot, D., Blanchet, S., Velez, L., Albouy, C., and Pellissier, L. 2020.
498 Global determinants of freshwater and marine fish genetic diversity. *Nat. Commun.* **11**(1):
499 1–9. Springer US. doi:10.1038/s41467-020-14409-7.

- 500 Marshall, J.L., and Camp, C.D. 2006. Environmental correlates of species and genetic richness in
501 lungless salamanders (family plethodontidae). *Acta Oecologica* **29**(1): 33–44.
502 doi:10.1016/j.actao.2005.07.008.
- 503 Miraldo, A., Li, S., Borregaard, M.K., Florez-Rodriguez, A., Gopalakrishnan, S., Rizvanovic,
504 M., Wang, Z., Rahbek, C., Marske, K.A., and Nogues-Bravo, D. 2016. An Anthropocene
505 map of genetic diversity. *Science* **353**(6307): 1532–1535. doi:10.1126/science.aaf4381.
- 506 Mittell, E.A., Nakagawa, S., and Hadfield, J.D. 2015. Are molecular markers useful predictors of
507 adaptive potential? *Ecol. Lett.* **18**(8): 772–778. doi:10.1111/ele.12454.
- 508 Munguía, M., Rahbek, C., Rangel, T.F., Diniz-Filho, J.A.F., and Araújo, M.B. 2012. Equilibrium
509 of global amphibian species distributions with climate. *PLoS One* **7**(4): e34420.
510 doi:10.1371/journal.pone.0034420.
- 511 Munro, D., and Treberg, J.R. 2017. A radical shift in perspective: Mitochondria as regulators of
512 reactive oxygen species. *J. Exp. Biol.* **220**(7): 1170–1180. doi:10.1242/jeb.132142.
- 513 Nei, M. 1973. Analysis of gene diversity in subdivided populations. *Proc. Natl. Acad. Sci. U. S.*
514 *A.* **70**(12): 3321–3323. doi:10.1073/pnas.70.12.3321.
- 515 Oliveira, B.F., São-Pedro, V.A., Santos-Barrera, G., Penone, C., and Costa, G.C. 2017.
516 *AmphiBIO*, a global database for amphibian ecological traits. *Sci. Data* **4**: 1–7.
517 doi:10.1038/sdata.2017.123.
- 518 Oliver, T.H., Heard, M.S., Isaac, N.J.B., Roy, D.B., Procter, D., Eigenbrod, F., Freckleton, R.,
519 Hector, A., Orme, C.D.L., Petchey, O.L., Proença, V., Raffaelli, D., Suttle, K.B., Mace,
520 G.M., Martín-López, B., Woodcock, B.A., and Bullock, J.M. 2015. Biodiversity and
521 resilience of ecosystem functions. *Trends Ecol. Evol.* **30**(11): 673–684.
522 doi:10.1016/j.tree.2015.08.009.
- 523 Peng, L., Zeng, Z., Wei, Z., Chen, A., Wood, E.F., and Sheffield, J. 2019. Determinants of the
524 ratio of actual to potential evapotranspiration. *Glob. Chang. Biol.* **25**(4): 1326–1343.
525 doi:10.1111/gcb.14577.
- 526 Pough, F.H. 1980. The Advantages of Ectothermy for Tetrapods. *Am. Nat.* **115**(1): 92–112.
- 527 Pyron, R.A., and Burbrink, F.T. 2014. Early origin of viviparity and multiple reversions to
528 oviparity in squamate reptiles. *Ecol. Lett.* **17**(1): 13–21. doi:10.1111/ele.12168.
- 529 Rodríguez, M.Á., Belmontes, J.A., and Hawkins, B.A. 2005. Energy, water and large-scale
530 patterns of reptile and amphibian species richness in Europe. *Acta Oecologica* **28**(1): 65–70.
531 doi:10.1016/j.actao.2005.02.006.
- 532 Romiguier, J., Gayral, P., Ballenghien, M., Bernard, A., Cahais, V., Chenuil, A., Chiari, Y.,
533 Dernet, R., Duret, L., Faivre, N., Loire, E., Lourenco, J.M., Nabholz, B., Roux, C.,
534 Tsagkogeorga, G., Weber, A.A.T., Weinert, L.A., Belkhir, K., Bierne, N., Glémin, S., and
535 Galtier, N. 2014. Comparative population genomics in animals uncovers the determinants of
536 genetic diversity. *Nature* **515**(7526): 261–263. doi:10.1038/nature13685.
- 537 Schmidt, C., Dray, S., and Garroway, C.J. 2020. Genetic and species-level biodiversity patterns
538 are linked by demography and ecological opportunity. *bioRxiv*.

- 539 doi:10.1101/2020.06.03.132092.
- 540 Schmidt, C., and Garroway, C.J. 2021a. The population genetics of urban and rural amphibians
541 in North America. *Mol. Ecol.* doi:10.1111/mec.16005.
- 542 Schmidt, C., and Garroway, C.J. 2021b. The conservation utility of mitochondrial genetic
543 diversity in macrogenetic research. *Conserv. Genet.* **22**(3): 323–327. doi:10.1007/s10592-
544 021-01333-6.
- 545 Shipley, B. 2016. *Cause and Correlation in Biology*, 2nd edition. Cambridge University Press,
546 Cambridge.
- 547 Smith, S.A., Stephens, P.R., and Wiens, J.J. 2005. Replicate patterns of species richness,
548 historical biogeography, and phylogeny in holarctic treefrogs. *Evolution* **59**(11): 2433–
549 2450. doi:10.1111/j.0014-3820.2005.tb00953.x.
- 550 Stein, A., Gerstner, K., and Kreft, H. 2014. Environmental heterogeneity as a universal driver of
551 species richness across taxa, biomes and spatial scales. *Ecol. Lett.* **17**(7): 866–880.
552 doi:10.1111/ele.12277.
- 553 Storch, D., Bohdalková, E., and Okie, J. 2018. The more-individuals hypothesis revisited: the
554 role of community abundance in species richness regulation and the productivity–diversity
555 relationship. *Ecol. Lett.* **21**(6): 920–937. doi:10.1111/ele.12941.
- 556 Stuart, S.N., Chanson, J.S., Cox, N.A., Young, B.E., Rodrigues, A.S.L., Fischman, D.L., and
557 Waller, R.W. 2004. Status and trends of amphibian declines and extinctions worldwide.
558 *Science* **306**(5702): 1783–1786. doi:10.1126/science.1103538.
- 559 Theodoridis, S., Fordham, D.A., Brown, S.C., Li, S., Rahbek, C., and Nogués-Bravo, D. 2020.
560 Evolutionary history and past climate change shape the distribution of genetic diversity in
561 terrestrial mammals. *Nat. Commun.* **11**(1): 2557. doi:10.1038/s41467-020-16449-5.
- 562 Trabucco, A., and Zomer, R. 2019. Global Aridity Index and Potential Evapotranspiration (ET0)
563 Climate Database v2. doi:10.6084/m9.figshare.7504448.v3.
- 564 Urban, M.C. 2004. Disturbance heterogeneity determines freshwater metacommunity structure.
565 *Ecology* **85**(11): 2971–2978. doi:10.1890/03-0631.
- 566 Weir, B.S., and Goudet, J. 2017. A Unified Characterization of Population Structure and
567 Relatedness. *Genetics* **206**(4): 2085–2103. doi:10.1534/genetics.116.198424.
- 568 Werner, E.E., Yurewicz, K.L., Skelly, D.K., and Relyea, R.A. 2007. Turnover in an amphibian
569 metacommunity: the role of local and regional factors. *Oikos* **116**(10): 1713–1725.
570 doi:10.1111/j.2007.0030-1299.16039.x.
- 571 Whittaker, R.J., Nogués-Bravo, D., and Araújo, M.B. 2007. Geographical gradients of species
572 richness: a test of the water-energy conjecture of Hawkins et al. (2003) using European data
573 for five taxa. *Glob. Ecol. Biogeogr.* **16**: 76–89. doi:10.1111/j.1466-822x.2006.00268.x.
- 574 Wright, D.H. 1983. Species-Energy Theory: An Extension of Species-Area Theory. *Oikos* **41**(3):
575 496–506.
- 576 Zeisset, I., and Beebee, T.J.C. 2008. Amphibian phylogeography: a model for understanding


- 577 historical aspects of species distributions. *Heredity*. **101**: 109–119.
578 doi:10.1038/hdy.2008.30.
- 579 Zhang, Y., and Wong, H.S. 2021. Are mitochondria the main contributor of reactive oxygen
580 species in cells? *J. Exp. Biol.* **224**(5). doi:10.1242/jeb.221606.

581 **Table 1.** Data summary. Summary of aggregated raw genetic data: mean gene diversity, mean number of loci, median number of
 582 individuals at sites per species. Species richness is the mean species richness at sites. Environmental variables (40 km
 583 buffer): potential evapotranspiration (PET); actual evapotranspiration (AET); heterogeneity is the mean land cover diversity measured
 584 with Simpson's Index. Ranges of values are given in parentheses where applicable.

585

586

Species (# sites)	Gene diversity	Loci	Individuals	Species richness	PET	AET	Heterogeneity
Anura							
<i>Ascaphus montanus</i> (100)	0.74 (0.41–0.87)	13	20 (10–41)	7.35 (3–11)	1097.84 (1006.52–1327.35)	506.86 (341.66–601.4)	0.54 (0.31–0.72)
<i>Lithobates pipiens</i> (5)	0.86 (0.83–0.9)	9	40 (20–46)	18.4 (16–21)	961.02 (924.69–1041.02)	656.24 (614.39–711.78)	0.74 (0.69–0.78)
<i>Lithobates sylvaticus</i> (90)	0.8 (0.7–0.9)	10.57 (7–15)	24 (5–36)	21.4 (11–38)	1062.37 (843.19–1437.5)	759.82 (669.33–1050.57)	0.76 (0.63–0.84)
<i>Pseudacris crucifer</i> (11)	0.75 (0.65–0.81)	11	28 (5–114)	16.18 (15–17)	984.96 (913.86–1122.62)	656.36 (569.9–689.03)	0.53 (0.27–0.8)
<i>Pseudacris streckeri</i> (17)	0.59 (0.38–0.82)	14	8 (5–22)	24.71 (15–34)	1479.33 (1257.25–1925.78)	878.45 (521.88–1001.36)	0.63 (0.3–0.86)
<i>Rana draytonii</i> (17)	0.51 (0.27–0.66)	15	13 (7–83)	9.41 (7–12)	1776.46 (1586.25–2179.73)	405.28 (331.77–493.63)	0.71 (0.55–0.8)
<i>Rana luteiventris</i> (25)	0.55 (0.37–0.69)	8	15 (6–205)	9.16 (5–11)	1200.7 (1077.99–1563.67)	494.77 (254–579.38)	0.66 (0.41–0.74)
<i>Rana pretiosa</i> (23)	0.32 (0.16–0.55)	10.39 (7–13)	34 (26–299)	9.35 (5–18)	1195.73 (924.34–1341.85)	515.32 (424.91–631.07)	0.55 (0.37–0.81)
Caudata							
<i>Ambystoma barbouri</i> (76)	0.78 (0.54–0.88)	11	22 (8–43)	28.28 (21–34)	1274.78 (1189.33–1337.51)	889.87 (811.77–967.52)	0.68 (0.55–0.75)
<i>Ambystoma maculatum</i> (97)	0.69 (0.45–0.75)	9.16 (7–14)	24 (8–36)	20.74 (16–35)	1074.81 (961.91–1406.61)	752.9 (682.96–931.33)	0.74 (0.37–0.8)
<i>Desmognathus fuscus</i> (5)	0.39 (0.15–0.65)	5	26 (22–35)	26.4 (26–27)	1258.69 (1224.99–1271.15)	827 (813.24–849.54)	0.62 (0.58–0.67)
<i>Dicamptodon aterrimus</i> (3)	0.48 (0.43–0.53)	9	91 (90–180)	6.33 (5–7)	1039.28 (1029.16–1046.55)	527.8 (469.46–564.46)	0.36 (0.32–0.41)
<i>Dicamptodon copei</i> (29)	0.69 (0.48–0.87)	11	18 (6–81)	13.86 (13–15)	905.66 (826.03–1074.59)	652.43 (628.31–678.36)	0.57 (0.48–0.74)
<i>Ensatina eschscholtzii</i> (4)	0.8 (0.75–0.84)	10	11.5 (5–19)	10 (9–11)	2140.32 (2039.14–2308.24)	368.69 (341.99–380.48)	0.63 (0.6–0.72)
<i>Hydromantes brunus</i> (6)	0.52 (0.38–0.61)	10	10 (7–18)	9.5 (8–11)	2021.39 (2008.05–2037.4)	347.91 (291.31–424.01)	0.7 (0.63–0.75)
<i>Hydromantes platycephalus</i> (15)	0.44 (0.25–0.64)	10	13 (6–21)	6.93 (4–9)	1786.55 (1523.97–2159.97)	466.68 (326.23–576.76)	0.65 (0.55–0.73)
<i>Plethodon albagula</i> (21)	0.48 (0.44–0.55)	20	16 (10–24)	31 (31–31)	1349.69 (1348.91–1350.48)	833.13 (832.48–833.85)	0.66 (0.66–0.66)
<i>Plethodon cinereus</i> (1)	0.6	7	122	22	1348.41 (1348.41–1348.41)	843.96	0.77
<i>Taricha granulosa</i> (9)	0.4 (0.26–0.77)	6	16 (6–32)	4 (2–13)	746.76 (614.95–1612.18)	522.65 (480.12–609.21)	0.73 (0.69–0.82)



587

588 **Figure 1.** (Top row) Maps of predicted genetic diversity, species richness, and genetic differentiation at genetic sample sites (points)
 589 based on spatial MEMs. MEMs were able to recover known patterns of species richness, which are negatively correlated with spatial
 590 patterns of genetic diversity. Patterns of genetic differentiation mirror those of species richness. (Bottom row) Maps depicting the
 591 environmental variables predicted to have simultaneous effects on genetic diversity and species richness, and variation partitioning
 592 results. Note land cover heterogeneity is a categorical variable and this map represents different land cover classes.




Figure 2. Structural equation model results. (A) Our hypothesized conceptual model. Line color denotes predicted directions of effect. (B) Results for model using environments within 40 km of sites (results for all spatial scales are in Tables S1-S4). Regression coefficients with standard errors are shown along each path. Paths between variables where no effect was detected are colored in gray (see Table S3 for a complete summary of all paths). R^2 , the proportion of variation explained by the model, is given for genetic diversity and species richness. For genetic diversity, R^2_m is the variation explained by fixed effects and R^2_c is the variation explained by both fixed effects and the random species effect.

Determinants of genetic diversity and species richness of North American amphibians

Supplementary Information

Figures S1-3

Tables S1-4

Genetic diversity

Species richness




Figure S1. Maps of raw data for genetic diversity, species richness, and genetic differentiation. Genetic diversity is Nei's gene diversity; species richness is the number of amphibian species' ranges overlapping each site; genetic differentiation is site-specific F_{ST} .




Figure S2. Relationship between genetic diversity and body size (snout-vent length). We detected no relationship using a mixed-effects model including a random intercept for species ($\beta = -0.03 \pm 0.2$ SE).




Figure S3. Spatial patterns (MEMs) for genetic diversity and species richness are correlated with environments: energy availability (potential evapotranspiration; PET), water availability (actual evapotranspiration; AET), and heterogeneity (land cover diversity). MEMs are ordered along the x-axis according to spatial scale explained, from broadest (MEM1) to finest (MEM18). MEMs to the left of the dashed lines indicate the broadest-scale patterns with Moran's $I > 0.25$ used to produce maps of genetic diversity and species richness. Light blue bars are MEMs explaining spatial patterns of species richness, and dark blue bars are MEMs explaining spatial patterns shared by genetic diversity and species richness. No MEMs explained spatial patterns in genetic diversity only.

1 **Table S1.** Path coefficients and standard errors for SEM model with heterogeneity measured
2 within a 10 km buffer around sites (Fisher's $C = 0.28$, $p = 0.87$, 2 degrees of freedom). Genetic
3 diversity $R^2_m = 0.04$ (variance explained by fixed effects); $R^2_c = 0.92$ (variance explained by
4 fixed and random effects); species richness $R^2 = 0.87$.

5

Response	Predictor	Estimate \pm SE
gene diversity	potential evapotranspiration	0.16 \pm 0.16
gene diversity	actual evapotranspiration	0.13 \pm 0.14
gene diversity	heterogeneity	0.16 \pm 0.09
species richness	potential evapotranspiration	0.18 \pm 0.02
species richness	actual evapotranspiration	0.90 \pm 0.02
species richness	heterogeneity	0.09 \pm 0.02

6

7 **Table S2.** Path coefficients and standard errors for SEM model with heterogeneity measured
8 within a 25 km buffer around sites (Fisher's $C = 0.59$, $p = 0.74$, 2 degrees of freedom). Genetic
9 diversity $R^2_m = 0.04$ (variance explained by fixed effects); $R^2_c = 0.92$ (variance explained by
10 fixed and random effects); species richness $R^2 = 0.88$.

11

Response	Predictor	Estimate \pm SE
gene diversity	potential evapotranspiration	0.20 \pm 0.16
gene diversity	actual evapotranspiration	0.07 \pm 0.16
gene diversity	heterogeneity	0.19 \pm 0.12
species richness	potential evapotranspiration	0.19 \pm 0.01
species richness	actual evapotranspiration	0.90 \pm 0.02
species richness	heterogeneity	0.10 \pm 0.02

12

13 **Table S3.** Path coefficients and standard errors for SEM model with heterogeneity measured
14 within a 40 km buffer around sites (Fisher's $C = 0.49$, $p = 0.78$, 2 degrees of freedom). Genetic
15 diversity $R^2_m = 0.04$ (variance explained by fixed effects); $R^2_c = 0.93$ (variance explained by
16 fixed and random effects); species richness $R^2 = 0.89$.

17

Response	Predictor	Estimate \pm SE
gene diversity	potential evapotranspiration	0.22 \pm 0.17
gene diversity	actual evapotranspiration	0.10 \pm 0.17
gene diversity	heterogeneity	0.15 \pm 0.11
species richness	potential evapotranspiration	0.20 \pm 0.01
species richness	actual evapotranspiration	0.91 \pm 0.01
species richness	heterogeneity	0.09 \pm 0.01

18

19

20 **Table S4.** Path coefficients and standard errors for SEM model with heterogeneity measured
21 within an 80 km buffer around sites (Fisher's $C = 0.40$, $p = 0.82$, 2 degrees of freedom). Genetic
22 diversity $R^2_m = 0.03$ (variance explained by fixed effects); $R^2_c = 0.92$ (variance explained by
23 fixed and random effects); species richness $R^2 = 0.90$.

24

Response	Predictor	Estimate \pm SE
gene diversity	potential evapotranspiration	0.16 \pm 0.19
gene diversity	actual evapotranspiration	0.24 \pm 0.20
gene diversity	heterogeneity	0.09 \pm 0.09
species richness	potential evapotranspiration	0.22 \pm 0.01
species richness	actual evapotranspiration	0.95 \pm 0.01
species richness	heterogeneity	0.07 \pm 0.01

25

26

27